

Moebius

By filip

January 3, 2014

Contents

1 More about complex numbers	2
2 Systems of linear equations	35
3 Quadratic equations	37
4 Vectors, Matrices	41
4.1 Vectors	41
4.2 Matrices	42
4.3 Eigenvalues and eigenvectors	49
4.4 Bilinear and Quadratic forms; Congruence	51
5 Unitary matrices	54
6 Hermitean matrices	73
7 Elementary complex geometry	80
8 Homogeneous coordinates in extended complex plane	99
8.1 Ratio and crossratio	115
8.2 Distance	124
9 Riemann sphere	133
10 Moebius transformations	149
10.1 Action on points	151
10.2 Moebius group	153
10.3 Special kinds of Moebius transformations	157
10.4 Decomposition	162
10.5 Cross ratio and moebius existence	163
10.6 Fixed points and moebius uniqueness	165
10.7 Pole	170
10.8 Antihomographies	171

10.9 Classification	171
-------------------------------	-----

11 Circline	173
11.1 Circline definition	174
11.2 Connections with circles on the Riemann sphere	183
11.3 Some special circlines	189
11.4 Moebius action on circlines	190
11.5 Conjugation, reciprocation and inversion of circlines	195
11.6 Circline uniqueness	197
11.6.1 Zero type circline uniqueness	197
11.6.2 Negative type circline uniqueness	199
11.7 Circline set cardinality	201
11.7.1 Diagonal circlines	201
11.7.2 Zero type circline set cardinality	203
11.7.3 Negative type circline set cardinality	205
11.7.4 Positive type circline set cardinality	207
11.7.5 Cardinality determines type	208
11.7.6 Circline set is injective	209
11.8 Symmetric points wrt. circline	210
11.9 Oriented circlines; discs	211
11.10 Some special oriented circlines and discs	224
11.11 Moebius action on oriented circlines and discs	225
11.12 Oriented circline uniqueness	234
11.13 Disc automorphisms	256
11.14 Angle between circlines	263

1 More about complex numbers

```
theory MoreComplex
imports Complex-Main
begin
```

```
lemma mult-pow2-lt0:
assumes b ≠ 0
shows a < 0 ⟷ b² * a < (0::real)
using assms
by (metis mult.commute mult-eq-0-iff mult-neg-pos mult-pos-pos not-less-iff-gr-or-eq
not-real-square-gt-zero power2-eq-square)
```

```
lemma mult-pow2-gt0:
assumes b ≠ 0
shows a > 0 ⟷ b² * a > (0::real)
using assms
```

by (*metis mult.commute mult-eq-0-iff mult-neg-pos mult-pos-pos not-less-iff-gr-or-eq not-real-square-gt-zero power2-eq-square*)

lemma *square-cancel*:
assumes $a^2 \geq b^2$ $a \geq 0$ $b \geq (0::real)$
shows $a \geq b$
using *real-sqrt-le-iff*[*of* b^2 a^2]
using *assms*
by *auto*

lemmas *complex-cnj* = *complex-cnj-diff* *complex-cnj-mult* *complex-cnj-add* *complex-cnj-divide* *complex-cnj-minus*

abbreviation *cor* \equiv *complex-of-real*

lemma [*simp*]: $cor - 1 = -1$
by (*simp add: of-real-neg-numeral*)

lemma [*simp*]: $- cor - 1 = 1$
by *simp*

lemma *rcis-cnj*: $cnj a = rcis (cmod a) (- arg a)$
by (*subst rcis-cmod-arg*[*of* a , *symmetric*]) (*simp add: complex-cnj cis-def rcis-def*)

lemma *cmod-cis* [*simp*]:
assumes $a \neq 0$
shows $cor (cmod a) * cis (\arg a) = a$
using *assms*
by (*metis rcis-cmod-arg rcis-def*)

lemma *cis-cmod* [*simp*]:
assumes $a \neq 0$
shows $cis (\arg a) * cor (cmod a) = a$
using *assms cmod-cis*[*of* a]
by (*simp add: field-simps*)

lemma *cor-squared*: $(cor x)^2 = cor (x^2)$
by (*simp add: power2-eq-square*)

lemma *cor-add*: $cor (a + b) = cor a + cor b$
by *auto*

lemma *cor-mult*: $cor (a * b) = cor a * cor b$
by *auto*

lemma *cor-sqrt-mult-cor-sqrt* [*simp*]:
shows $cor (\sqrt{A}) * cor (\sqrt{A}) = cor |A|$

```

using assms
by (metis cor-mult real-sqrt-abs2 real-sqrt-mult-distrib2)

```

```

lemma [simp]: (Complex a b) * 2 = Complex (2*a) (2*b)
by (metis complex-add mult-2 mult-2-right)

```

```

lemma re-complex:
  Complex (Re z) 0 = (z + cnj z)/2
by (cases z) simp

```

```

lemma im-complex:
  Complex 0 (Im z) = (z - cnj z)/2
by (cases z) simp

```

```

lemma Complex-scale1: Complex (a * b) (a * c) = cor a * Complex b c
unfolding complex-of-real-def
by auto

```

```

lemma Complex-scale2: Complex (a * c) (b * c) = Complex a b * cor c
unfolding complex-of-real-def
by auto

```

```

lemma Complex-scale3: Complex (a / b) (a / c) = cor a * Complex (1 / b) (1
/ c)
unfolding complex-of-real-def
by auto

```

```

lemma Complex-scale4: c ≠ 0  $\implies$  Complex (a / c) (b / c) = Complex a b / cor
c
unfolding complex-of-real-def
by (auto simp add: field-simps)

```

```

lemma complex-mult-cnj-cmod:
  z * cnj z = cor ((cmod z)2)
by (cases z) (simp add: complex-of-real-def, simp add: power2-eq-square)

```

```

lemma
  cmod-square: (cmod z)2 = Re (z * cnj z)
using complex-mult-cnj-cmod[of z]
by (simp add: power2-eq-square)

```

```

lemma cnjE:
  assumes x ≠ 0
  shows cnj x = cor ((cmod x)2) / x
using complex-mult-cnj-cmod[of x] assms

```

```

by (auto simp add: field-simps)

lemma cmod-mult [simp]: cmod (a * b) = cmod a * cmod b
  unfolding cmod-def
  by (metis complex-norm-def norm-mult)

lemma cmod-divide [simp]: cmod (a / b) = cmod a / cmod b
  unfolding cmod-def
  by (metis complex-norm-def norm-divide)

lemma [simp]: cmod (z / cor k) = cmod z / |k|
  by auto

lemma [simp]: cmod (z*z1 - z*z2) = cmod z*cmod(z1 - z2)
  by (metis bounded-bilinear.diff-right bounded-bilinear-mult cmod-mult)

lemma cmod-eqI:
  assumes z1 * cnj z1 = z2 * cnj z2
  shows cmod z1 = cmod z2
  using assms
  by (subst complex-mod-sqrt-Re-mult-cnj)+ auto

lemma cmod-eqE:
  assumes cmod z1 = cmod z2
  shows z1 * cnj z1 = z2 * cnj z2
proof-
  from assms have cor ((cmod z1)^2) = cor ((cmod z2)^2)
    by auto
  thus ?thesis
    using complex-mult-cnj-cmod
    by auto
qed

lemma [simp]: cmod a = 1  $\longleftrightarrow$  a*cnj a = 1
  by (metis cmod-eqE cmod-eqI complex-cnj-one monoid-mult-class.mult.left-neutral norm-one)

```

```

abbreviation is-real where
  is-real z ≡ Im z = 0

lemma complex-eq-if-Re-eq:
  assumes is-real z1 is-real z2
  shows z1 = z2  $\longleftrightarrow$  Re z1 = Re z2
  using assms
  by (cases z1, cases z2) auto

lemma mult-reals:

```

```

assumes is-real a is-real b
shows is-real (a * b)
using assms
by auto

lemma div-reals:
assumes is-real a is-real b
shows is-real (a / b)
using assms
by (simp add: divide-inverse complex-inverse-def)

lemma complex-of-real-Re:
assumes is-real k
shows cor (Re k) = k
using assms
by (cases k) (auto simp add: complex-of-real-def)

lemma is-real-complex-of-real:
is-real (cor x)
by auto

lemma cor-cmod-real:
assumes is-real a
shows cor (cmod a) = a ∨ cor (cmod a) = -a
using assms
unfolding cmod-def
by (cases Re a > 0) (auto, (metis complex-of-real-Re)+)

lemma eq-cnj-iff-real:
z = cnj z ↔ is-real z
by (cases z) auto

lemma Re-divide-real:
assumes is-real b b ≠ 0
shows Re (a / b) = (Re a) / (Re b)
using assms
unfolding complex-divide-def
by (cases a, cases b) (auto simp add: field-simps power2-eq-square)

lemma Re-mult-real:
assumes is-real a
shows Re (a * b) = (Re a) * (Re b)
using assms
by auto

lemma Im-mult-real:
assumes is-real a
shows Im (a * b) = (Re a) * (Im b)
using assms

```

```

by auto

lemma Im-divide-real:
  assumes is-real b  $b \neq 0$ 
  shows  $\text{Im}(a / b) = (\text{Im } a) / (\text{Re } b)$ 
  using assms
  by (cases a, cases b) (auto simp add: complex-divide-def field-simps power2-eq-square)

lemma [simp]:  $\text{Re}(x / 2) = \text{Re } x / 2$ 
  using Re-divide-real[of 2 x]
  by simp

lemma [simp]:  $\text{Re}(2 * x) = 2 * \text{Re } x$ 
  using Re-mult-real[of 2 x]
  by simp

lemma Re-sgn:
  assumes is-real R
  shows  $\text{Re}(\text{sgn } R) = \text{sgn}(\text{Re } R)$ 
  using assms
  by (metis Re-sgn complex-of-real-Re norm-of-real real-sgn-eq)

```

abbreviation *rot90* **where**
 $\text{rot90 } z \equiv \text{Complex}(-\text{Im } z) (\text{Re } z)$

lemma *rot90-ii*: $\text{rot90 } z = z * ii$
 by (cases *z*) *simp*

abbreviation *cnj-mix* **where**
 $\text{cnj-mix } z1 z2 \equiv \text{cnj } z1 * z2 + z1 * \text{cnj } z2$

lemma *cnj-mix-minus*:
 shows $\text{cnj } z1 * z2 - z1 * \text{cnj } z2 = ii * \text{cnj-mix}(\text{rot90 } z1) z2$
 using *assms*
 by (cases *z1*, cases *z2*) *simp*

lemma *cnj-mix-minus'*:
 shows $\text{cnj } z1 * z2 - z1 * \text{cnj } z2 = \text{rot90}(\text{cnj-mix}(\text{rot90 } z1) z2)$
 using *assms*
 by (cases *z1*, cases *z2*) *simp*

lemma *cnj-mix-real*:
 is-real (*cnj-mix z1 z2*)
 by (cases *z1*, cases *z2*) *simp*

abbreviation *scalprod* **where**

scalprod z1 z2 ≡ cnj-mix z1 z2 / 2

```

lemma cos-periodic-pi2:  $\cos(pi + x) = -\cos x$ 
  using cos-periodic-pi[of x]
  by (simp add: field-simps)

lemma cos-periodic-pi3:  $\cos(x - pi) = -\cos x$ 
  by (smt cos-periodic-pi)

lemma cos-periodic-4 [simp]:  $\cos(pi - x) = -\cos x$ 
  by (metis cos-minus cos-periodic-pi2 minus-real-def)

lemma sin-periodic-pi3:  $\sin(x - pi) = -\sin x$ 
  by (smt sin-periodic-pi)

lemma cos-lt-zero:
  assumes  $x > pi/2 \wedge x \leq pi$ 
  shows  $\cos x < 0$ 
  using cos-gt-zero-pi[of pi - x] assms
  by simp

lemma sin-kpi:
  fixes k::int
  shows  $\sin(real k * pi) = 0$ 
  using sin-npi[of nat k]
  using sin-npi[of nat (-k)]
  by (cases k ≥ 0) auto

lemma cos-kpi-odd:
  fixes k::int
  assumes odd k
  shows  $\cos(real k * pi) = -1$ 
  proof (cases k ≥ 0)
    case True
    hence odd (nat k)
    using ⟨odd k⟩
    by (metis pos-int-even-equiv-nat-even)
    thus ?thesis
      using ⟨k ≥ 0⟩ cos-npi[of nat k]
      by auto
  next
    case False
    hence -k ≥ 0 odd (nat (-k))
    using ⟨odd k⟩
    by (auto, smt even-neg pos-int-even-equiv-nat-even)
    thus ?thesis
      using cos-npi[of nat (-k)]

```

```

    by auto
qed

lemma cos-kpi-even:
  fixes k::int
  assumes even k
  shows cos (real k * pi) = 1
proof (cases k ≥ 0)
  case True
  hence even (nat k)
  using ⟨even k⟩
  by (metis pos-int-even-equiv-nat-even)
thus ?thesis
  using ⟨k ≥ 0⟩ cos-npi[of nat k]
  by auto
next
  case False
  hence −k ≥ 0 even (nat (−k))
  using ⟨even k⟩
  by (auto, smt even-neg pos-int-even-equiv-nat-even)
thus ?thesis
  using cos-npi[of nat (−k)]
  by auto
qed

lemma sin-pi2-kpi-odd:
  fixes k::int
  assumes odd k
  shows sin (pi / 2 + real k * pi) = −1
using assms
by (simp add: sin-add cos-kpi-odd)

lemma sin-pi2-kpi-even:
  fixes k::int
  assumes even k
  shows sin (pi / 2 + real k * pi) = 1
using assms
by (simp add: sin-add cos-kpi-even)

lemma cos-zero-iff-int:
  shows cos x = 0 ↔ (exists k::int. odd k ∧ x = real k * (pi / 2))
proof
  assume cos x = 0
  then obtain n::nat where *: x = real n * (pi / 2) ∨ x = −(real n * (pi / 2))
  and odd n
  using cos-zero-iff[of x]
  by blast
  hence (odd (int n) ∧ x = real (int n) * (pi / 2)) ∨ (odd (−int n) ∧ x = real (−int n) * pi / 2)

```

```

    by (auto simp add: Parity.transfer-int-nat-relations)
  thus  $\exists k:\text{int}. \text{odd } k \wedge x = \text{real } k * (\pi / 2)$ 
    by (metis times-divide-eq-right)
next
  assume  $\exists k:\text{int}. \text{odd } k \wedge x = \text{real } k * (\pi / 2)$ 
  then obtain  $k:\text{int}$  where  $*: \text{odd } k \wedge x = \text{real } k * (\pi / 2)$ 
    by blast
  show  $\cos x = 0$ 
  proof (cases  $k \geq 0$ )
    case True
    hence  $\exists n:\text{nat}. \text{odd } n \wedge x = \text{real } n * (\pi / 2)$ 
      using *
      by (rule-tac  $x=\text{nat } k$  in exI) (auto simp add: pos-int-even-equiv-nat-even)
    thus ?thesis
      using cos-zero-iff[of x]
      by auto
  next
    case False
    hence  $\exists n:\text{nat}. \text{odd } n \wedge x = -(\text{real } n * (\pi / 2))$ 
      using *
      by (rule-tac  $x=\text{nat } (-k)$  in exI, auto) (smt even-neg pos-int-even-equiv-nat-even)
    thus ?thesis
      using cos-zero-iff[of x]
      by auto
  qed
qed

lemma sin-zero-iff-int:
   $\sin x = 0 \longleftrightarrow (\exists k:\text{int}. \text{even } k \wedge x = \text{real } k * (\pi / 2))$ 
proof-
  have  $\sin x = 0 \longleftrightarrow \cos(x - \pi/2) = 0$ 
    using cos-minus[of  $x - \pi/2$ ]
    by (simp add: sin-cos-eq)
  hence  $\sin x = 0 \longleftrightarrow (\exists k:\text{int}. \text{odd } k \wedge x - \pi/2 = \text{real } k * (\pi / 2))$ 
    using cos-zero-iff-int
    by simp
  thus ?thesis
    by auto (rule-tac  $x=k+1$  in exI, simp add: field-simps, rule-tac  $x=k-(1:\text{int})$ 
in exI, simp add: field-simps)
qed

lemma cos0-sin1:
  assumes  $\cos \varphi = 0 \sin \varphi = 1$ 
  shows  $\exists k:\text{int}. \varphi = \pi/2 + 2*k*\pi$ 
proof-
  from (cos  $\varphi = 0$ )
  obtain  $k:\text{int}$  where  $\text{odd } k \wedge \varphi = \text{real } k * (\pi / 2)$ 
    using cos-zero-iff-int[of  $\varphi$ ]
    by auto

```

```

then obtain k':int where k = 2*k' + 1
  by (metis odd-equiv-def)
hence φ = pi/2 + (real k' * pi)
  using ⟨φ = real k * (pi / 2)⟩
  by (auto simp add: field-simps)
hence even k'
  using ⟨sin φ = 1⟩ sin-pi2-kpi-odd[of k]
  by auto
thus ?thesis
  using ⟨φ = pi / 2 + (real k' * pi)⟩
  unfolding even-def
  by auto
qed

lemma cos-0-iff-normalized:
assumes cos φ = 0 –pi < φ φ ≤ pi
shows φ = pi/2 ∨ φ = –pi/2
proof –
  obtain k::int where odd k φ = real k * pi/2
    using cos-zero-iff-int[of φ] assms(1)
    by auto
  thus ?thesis
  proof (cases k > 1 ∨ k < –1)
    case True
    hence k ≥ 3 ∨ k ≤ –3
      using ⟨odd k⟩
    by auto (smt odd-one-int odd-plus-odd, smt odd-one-int odd-plus-even odd-plus-odd)
    hence φ ≥ 3*pi/2 ∨ φ ≤ –3*pi/2
      using ⟨φ = real k * pi/2⟩
      by auto
    thus ?thesis
      using ⟨– pi < φ⟩ ⟨φ ≤ pi⟩
      by auto
  next
    case False
    hence k = –1 ∨ k = 0 ∨ k = 1
      by auto
    hence k = –1 ∨ k = 1
      using ⟨odd k⟩
      by auto
    thus ?thesis
      using ⟨φ = real k * pi/2⟩
      by auto
  qed
qed

lemma sin-0-iff-normalized:
assumes sin φ = 0 –pi < φ φ ≤ pi
shows φ = 0 ∨ φ = pi

```

```

proof-
  obtain k::int where even k φ = real k * pi/2
    using sin-zero-iff-int[of φ] assms(1)
    by auto
  thus ?thesis
  proof (cases k > 2 ∨ k < 0)
    case True
    hence k ≥ 4 ∨ k ≤ -2
      using ⟨even k⟩
      by auto (smt even-difference odd-one-int)+
    hence φ ≥ 2*pi ∨ φ ≤ -pi
    proof
      assume 4 ≤ k
      hence 4 * pi/2 ≤ φ
        by (subst ⟨φ = real k * pi/2⟩) auto
      thus ?thesis
        by simp
    next
      assume k ≤ -2
      hence real k ≤ -2
        by simp
      hence -2*pi/2 ≥ φ
        by (subst ⟨φ = real k * pi/2⟩, metis mult-right-mono pi-half-ge-zero
          times-divide-eq-right)
      thus ?thesis
        by simp
    qed
    thus ?thesis
      using ⟨- pi < φ⟩ ⟨φ ≤ pi⟩
      by auto
  next
    case False
    hence k = 0 ∨ k = 1 ∨ k = 2
      by auto
    hence k = 0 ∨ k = 2
      using ⟨even k⟩
      by auto
    thus ?thesis
      using ⟨φ = real k * pi/2⟩
      by auto
    qed
  qed

lemma cos1-sin0:
  assumes cos φ = 1 sin φ = 0
  shows ∃ k::int. φ = 2*k*pi
proof-
  from ⟨sin φ = 0⟩
  obtain k::int where even k φ = real k * (pi / 2)

```

```

using sin-zero-iff-int[of  $\varphi$ ]
by auto
then obtain  $k'::int$  where  $k = 2*k'$ 
by (metis even-equiv-def)
hence  $\varphi = \text{real } k' * \pi$ 
using ⟨ $\varphi = \text{real } k * (\pi / 2)$ ⟩
by (auto simp add: field-simps)
hence even  $k'$ 
using ⟨ $\cos \varphi = 1$ ⟩ cos-kpi-odd[of  $k$ ]
by auto
thus ?thesis
using ⟨ $\varphi = \text{real } k' * \pi$ ⟩
unfolding even-def
by auto
qed

```

```

lemma sin-cos-eq:
fixes  $a b :: \text{real}$ 
assumes  $\cos a = \cos b$   $\sin a = \sin b$ 
shows  $\exists k::\text{int}. a - b = 2*k*\pi$ 
proof –
  from assms have  $\sin(a - b) = 0$   $\cos(a - b) = 1$ 
  using sin-diff[of  $a b$ ] cos-diff[of  $a b$ ]
  by auto
  thus ?thesis
  using cos1-sin0
  by auto
qed

lemma sin-monotone-2pi: assumes  $-(\pi / 2) \leq y$  and  $y < x$  and  $x \leq \pi / 2$ 
shows  $\sin y < \sin x$ 
proof –
  have  $0 \leq y + \pi / 2$  and  $y + \pi / 2 < x + \pi / 2$  and  $x + \pi / 2 \leq \pi$ 
  using pi-ge-two and assms by auto
  from cos-monotone-0-pi[OF this] show ?thesis unfolding minus-sin-cos-eq[symmetric]
  by auto
qed

lemma sin-inj:
assumes  $\alpha \neq \alpha' - \pi / 2 \leq \alpha \wedge \alpha \leq \pi / 2 - \pi / 2 \leq \alpha' \wedge \alpha' \leq \pi / 2$ 
shows  $\sin \alpha \neq \sin \alpha'$ 
using assms
using sin-monotone-2pi[of  $\alpha \alpha']$  sin-monotone-2pi[of  $\alpha' \alpha$ ]
by (cases  $\alpha < \alpha')$  auto

lemma arccos-le-pi2:
assumes  $a \geq 0$   $a \leq 1$ 

```

```

shows arccos a ≤ pi/2
using assms
by (smt antisym arccos-cos arccos-ubound cos-arccos cos-monotone-0-pi' cos-pi-half
pi-half-ge-zero)

definition atan2 where
atan2 y x =
(if x > 0 then arctan (y/x)
else if x < 0 then
  if y > 0 then arctan (y/x) + pi else arctan (y/x) - pi
else
  if y > 0 then pi/2 else if y < 0 then -pi/2 else 0)

lemma atan2-bounded: -pi ≤ atan2 y x ∧ atan2 y x < pi
using arctan-bounded[of y/x] zero-le-arctan-iff[of y/x] arctan-le-zero-iff[of y/x]
zero-less-arctan-iff[of y/x] arctan-less-zero-iff[of y/x]
using divide-neg-neg[of y x] divide-neg-pos[of y x] divide-pos-pos[of y x] divide-pos-neg[of
y x]
unfolding atan2-def
by (simp (no-asm-simp)) auto

lemma cos-periodic-nat[simp]: fixes n :: nat shows cos (x + n * (2 * pi)) = cos
x
proof (induct n arbitrary: x)
case (Suc n)
have split-pi-off: x + (Suc n) * (2 * pi) = (x + n * (2 * pi)) + 2 * pi
  unfolding Suc-eq-plus1 real-of-nat-add real-of-one distrib-right by auto
  show ?case unfolding split-pi-off using Suc by auto
qed auto

lemma cos-periodic-int[simp]: fixes i :: int shows cos (x + i * (2 * pi)) = cos x
proof (cases 0 ≤ i)
case True hence i-nat: real i = nat i by auto
show ?thesis unfolding i-nat by auto
next
case False hence i-nat: i = - real (nat (-i)) by auto
have cos x = cos (x + i * (2 * pi) - i * (2 * pi)) by auto
also have ... = cos (x + i * (2 * pi))
  unfolding i-nat mult-minus-left diff-minus-eq-add by (rule cos-periodic-nat)
finally show ?thesis by auto
qed

```

abbreviation canon-ang- P where
 $\text{canon-ang-}P \alpha \alpha' \equiv (-\pi < \alpha' \wedge \alpha' \leq \pi) \wedge (\exists k::int. \alpha - \alpha' = 2*k*\pi)$

```

definition canon-ang :: real  $\Rightarrow$  real ( $\lfloor \cdot \rfloor$ ) where
 $\lfloor \alpha \rfloor = (\text{THE } \alpha'. \text{canon-ang-}P \alpha \alpha')$ 

lemma canon-ang-ex:
  shows  $\exists \alpha'. \text{canon-ang-}P \alpha \alpha'$ 
proof-
  have ***:  $\forall \alpha: \text{real}. \exists \alpha'. 0 < \alpha' \wedge \alpha' \leq 1 \wedge (\exists k: \text{int}. \alpha' = \alpha - k)$ 
  proof
    fix  $\alpha: \text{real}$ 
    show  $\exists \alpha' > 0. \alpha' \leq 1 \wedge (\exists k: \text{int}. \alpha' = \alpha - \text{real } k)$ 
    proof (cases  $\alpha = \text{floor } \alpha$ )
      case True
      thus ?thesis
        by (rule-tac  $x=\alpha - \text{floor } \alpha + 1$  in exI, auto) (rule-tac  $x=\text{floor } \alpha - 1$  in exI, auto)
    next
      case False
      thus ?thesis
        using real-of-int-floor-ge-diff-one[of  $\alpha$ ]
        using real-of-int-floor-le[of  $\alpha$ ]
        by (rule-tac  $x=\alpha - \text{floor } \alpha$  in exI) (metis antisym diff-self floor-subtract
          le-cases le-iff-diff-le-0 less-int-code(1) not-leE zero-less-floor)
    qed
  qed

  have **:  $\forall \alpha: \text{real}. \exists \alpha'. 0 < \alpha' \wedge \alpha' \leq 2 \wedge (\exists k: \text{int}. \alpha - \alpha' = 2*k - 1)$ 
  proof
    fix  $\alpha: \text{real}$ 
    from **[rule-format, of  $(\alpha + 1)/2$ ]
    obtain  $\alpha'$  and  $k: \text{int}$  where  $0 < \alpha' \wedge \alpha' \leq 1 \wedge \alpha' = (\alpha + 1)/2 - k$ 
      by force
    hence  $0 < \alpha' \wedge \alpha' \leq 1 \wedge \alpha' = \alpha/2 - k + 1/2$ 
      by auto
    thus  $\exists \alpha' > 0. \alpha' \leq 2 \wedge (\exists k: \text{int}. \alpha - \alpha' = \text{real } (2 * k - 1))$ 
      by (rule-tac  $x=2*\alpha'$  in exI) auto
  qed
  have *:  $\forall \alpha: \text{real}. \exists \alpha'. -1 < \alpha' \wedge \alpha' \leq 1 \wedge (\exists k: \text{int}. \alpha - \alpha' = 2*k)$ 
  proof
    fix  $\alpha: \text{real}$ 
    from ** obtain  $\alpha'$  and  $k: \text{int}$  where
       $0 < \alpha' \wedge \alpha' \leq 2 \wedge \alpha - \alpha' = 2*k - 1$ 
      by force
    thus  $\exists \alpha' > -1. \alpha' \leq 1 \wedge (\exists k. \alpha - \alpha' = \text{real } (2 * (k: \text{int})))$ 
      by (rule-tac  $x=\alpha' - 1$  in exI) (auto simp add: field-simps)
  qed
  show ?thesis
    using *[rule-format, of  $\alpha / \pi$ ]
    apply auto

```

```

apply (rule-tac  $x=\alpha'*pi$  in exI)
by (auto simp add: field-simps) (metis mult.commute mult-minus1-right not-less
pi-gt-zero real-mult-le-cancel-iff2)
qed

lemma canon-ang-unique:
assumes canon-ang-P  $\alpha \alpha'$  canon-ang-P  $\alpha \alpha''$ 
shows  $\alpha' = \alpha''$ 
proof-
obtain  $k1::int$  where  $\alpha - \alpha' = 2*k1*pi$ 
using assms(1)
by auto
obtain  $k2::int$  where  $\alpha - \alpha'' = 2*k2*pi$ 
using assms(2)
by auto
hence  $-\alpha' + \alpha'' = 2*(k1 - k2)*pi$ 
using ⟨ $\alpha - \alpha' = 2*k1*pi$ ⟩
by (simp add:field-simps)
moreover
have  $-\alpha' + \alpha'' < 2 * pi$   $-\alpha' + \alpha'' > -2*pi$ 
using assms
by auto
ultimately
have  $-\alpha' + \alpha'' = 0$ 
by auto
thus ?thesis
by auto
qed

lemma canon-ang:
 $-pi < |\alpha| \ |\alpha| \leq pi \ \exists \ k::int. \alpha - |\alpha| = 2*k*pi$ 
proof-
obtain  $\alpha'$  where canon-ang-P  $\alpha \alpha'$ 
using canon-ang-ex[of  $\alpha$ ]
by auto
have canon-ang-P  $\alpha \ |\alpha|$ 
unfolding canon-ang-def
proof (rule theI[where  $a=\alpha'$ ])
show canon-ang-P  $\alpha \alpha'$ 
by fact
next
fix  $\alpha''$ 
assume canon-ang-P  $\alpha \alpha''$ 
thus  $\alpha'' = \alpha'$ 
using ⟨canon-ang-P  $\alpha \alpha'$ ⟩
using canon-ang-unique[of  $\alpha' \alpha \alpha''$ ]
by simp
qed

```

```

thus  $-pi < \lfloor \alpha \rfloor$   $\lfloor \alpha \rfloor \leq pi \exists k::int. \alpha - \lfloor \alpha \rfloor = 2*k*pi$ 
    by auto
qed

lemma canon-ang-id:
assumes  $-pi < \alpha \wedge \alpha \leq pi$ 
shows  $\lfloor \alpha \rfloor = \alpha$ 
using assms
using canon-ang-unique[of canon-ang  $\alpha$   $\alpha$   $\alpha$ ] canon-ang[of  $\alpha$ ]
by auto

lemma canon-ang-eq:
assumes  $\exists k::int. \alpha' - \lfloor \alpha' \rfloor = 2*k*pi$ 
shows  $\lfloor \alpha' \rfloor = \lfloor \alpha'' \rfloor$ 
proof-
obtain  $k'::int$  where  $*: -pi < \lfloor \alpha' \rfloor$   $\lfloor \alpha' \rfloor \leq pi$   $\alpha' - \lfloor \alpha' \rfloor = 2 * real k' * pi$ 
    using canon-ang[of  $\alpha'$ ]
    by auto

obtain  $k''::int$  where  $**: -pi < \lfloor \alpha'' \rfloor$   $\lfloor \alpha'' \rfloor \leq pi$   $\alpha'' - \lfloor \alpha'' \rfloor = 2 * real k'' * pi$ 
    using canon-ang[of  $\alpha''$ ]
    by auto

obtain  $k::int$  where  $***: \alpha' - \lfloor \alpha'' \rfloor = 2*k*pi$ 
    using assms
    by auto

have  $\exists m::int. \alpha' - \lfloor \alpha'' \rfloor = 2 * m * pi$ 
    using **(3) ***
    by (rule-tac  $x=k+k''$  in exI) (auto simp add: field-simps)

thus ?thesis
    using canon-ang-unique[of  $\lfloor \alpha' \rfloor$   $\alpha'$   $\lfloor \alpha'' \rfloor$ ] * **
    by auto
qed

lemma canon-ang-eqI:
assumes  $\exists k::int. \alpha' - \alpha = 2 * k * pi - pi < \alpha' \wedge \alpha' \leq pi$ 
shows  $\lfloor \alpha \rfloor = \alpha'$ 
using assms
using canon-ang-eq[of  $\alpha'$   $\alpha$ ]
using canon-ang-id[of  $\alpha'$ ]
by auto

lemma canon-ang-arg:
 $\lfloor arg z \rfloor = arg z$ 
using canon-ang-id[of arg z] arg-bounded
by simp

```

```

lemma canon-ang-uminus:
  assumes  $|\alpha| \neq pi$ 
  shows  $|- \alpha| = - |\alpha|$ 
  proof (rule canon-ang-eqI)
    show  $\exists x::int. - |\alpha| -- \alpha = 2 * real x * pi$ 
    using canon-ang(3)[of  $\alpha$ ]
    by (metis minus-diff-eq minus-diff-minus)
  next
    show  $- pi < - |\alpha| \wedge - |\alpha| \leq pi$ 
    using canon-ang(1)[of  $\alpha$ ] canon-ang(2)[of  $\alpha$ ] assms
    by auto
  qed

lemma canon-ang-uminus-pi:
  assumes  $|\alpha| = pi$ 
  shows  $|- \alpha| = |\alpha|$ 
  proof (rule canon-ang-eqI)
    obtain  $k::int$  where  $\alpha - |\alpha| = 2 * real k * pi$ 
    using canon-ang(3)[of  $\alpha$ ]
    by auto
    thus  $\exists x::int. |\alpha| -- \alpha = 2 * real x * pi$ 
    using assms
    by (rule-tac  $x=k+(1::int)$  in exI) (auto simp add: field-simps)
  next
    show  $- pi < |\alpha| \wedge |\alpha| \leq pi$ 
    using assms
    by auto
  qed

lemma canon-ang-diff:
   $|\alpha - \beta| = ||\alpha| - |\beta||$ 
  proof (rule canon-ang-eq)
    show  $\exists x::int. \alpha - \beta - (|\alpha| - |\beta|) = 2 * real x * pi$ 
  proof-
    obtain  $k1::int$  where  $\alpha - |\alpha| = 2*k1*pi$ 
    using canon-ang(3)
    by auto
    moreover
    obtain  $k2::int$  where  $\beta - |\beta| = 2*k2*pi$ 
    using canon-ang(3)
    by auto
    ultimately
    show ?thesis
    by (rule-tac  $x=k1 - k2$  in exI) (auto simp add: field-simps)
  qed
  qed

lemma canon-ang-sum:
   $|\alpha + \beta| = ||\alpha| + |\beta||$ 

```

```

proof (rule canon-ang-eq)
  show  $\exists x:\text{int}.$   $\alpha + \beta - (\lfloor \alpha \rfloor + \lfloor \beta \rfloor) = 2 * \text{real } x * pi$ 
  proof-
    obtain  $k1:\text{int}$  where  $\alpha - \lfloor \alpha \rfloor = 2*k1*pi$ 
      using canon-ang(3)
      by auto
    moreover
      obtain  $k2:\text{int}$  where  $\beta - \lfloor \beta \rfloor = 2*k2*pi$ 
        using canon-ang(3)
        by auto
    ultimately
      show ?thesis
        by (rule-tac x=k1 + k2 in exI) (auto simp add: field-simps)
  qed
qed

lemma canon-ang-plus-pi1:
  assumes  $0 < \alpha \leq 2*pi$ 
  shows  $\lfloor \alpha + pi \rfloor = \alpha - pi$ 
  proof (rule canon-ang-eqI)
    show  $\exists x:\text{int}.$   $\alpha - pi - (\alpha + pi) = 2 * \text{real } x * pi$ 
      by (rule-tac x=-1 in exI) auto
  next
    show  $-pi < \alpha - pi \wedge \alpha - pi \leq pi$ 
      using assms
      by auto
  qed

lemma canon-ang-plus-pi2:
  assumes  $-2*pi < \alpha \leq 0$ 
  shows  $\lfloor \alpha + pi \rfloor = \alpha + pi$ 
  proof (rule canon-ang-id)
    show  $-pi < \alpha + pi \wedge \alpha + pi \leq pi$ 
      using assms
      by auto
  qed

lemma canon-ang-minus-pi1:
  assumes  $0 < \alpha \leq 2*pi$ 
  shows  $\lfloor \alpha - pi \rfloor = \alpha - pi$ 
  proof (rule canon-ang-id)
    show  $-pi < \alpha - pi \wedge \alpha - pi \leq pi$ 
      using assms
      by auto
  qed

lemma canon-ang-minus-pi2:
  assumes  $-2*pi < \alpha \leq 0$ 
  shows  $\lfloor \alpha - pi \rfloor = \alpha + pi$ 

```

```

proof (rule canon-ang-eqI)
  show  $\exists x:\text{int. } \alpha + pi - (\alpha - pi) = 2 * \text{real } x * pi$ 
    by (rule-tac x=1 in exI) auto
next
  show  $-pi < \alpha + pi \wedge \alpha + pi \leq pi$ 
    using assms
    by auto
qed

lemma [simp]:  $\lfloor 0 \rfloor = 0$ 
  using canon-ang-eqI[of 0 0]
  by simp

lemma canon-ang-cos [simp]:  $\cos \lfloor \alpha \rfloor = \cos \alpha$ 
proof-
  obtain  $x:\text{int}$  where  $\alpha = \lfloor \alpha \rfloor + pi * (\text{real } x * 2)$ 
    using canon-ang(3)[of α]
    by (auto simp add: field-simps)
  thus ?thesis
    using cos-periodic-int[of ⌊α⌋ x]
    by (simp add: field-simps)
qed

lemma [simp]:  $cis \varphi * cis (-\varphi) = 1$ 
  by (metis cis-mult cis-zero right-minus)

lemma cis-eq:
  assumes  $cis a = cis b$ 
  shows  $\exists k:\text{int. } a - b = 2 * k * pi$ 
  using assms sin-cos-eq[of a b]
  using Re-cis[of a] Re-cis[of b] Im-cis[of a] Im-cis[of b]
  by (cases cis a, cases cis b) auto

lemma cis-inj:
  assumes  $cis \alpha = cis \alpha' - pi < \alpha \leq pi - pi < \alpha' \alpha' \leq pi$ 
  shows  $\alpha = \alpha'$ 
  using assms
  by (metis arg-unique sgn-cis)

lemma re-complex-zero-arg1:
  assumes  $\arg z = pi/2 \vee \arg z = -pi/2$ 
  shows  $\text{Re } z = 0$ 
  using assms
  using rcis-cmod-arg[of z] Re-rcis[of cmod z arg z]
  by (metis cos-minus cos-pi-half minus-divide-left mult-eq-0-iff)

```

```

lemma re-complex-zero-arg2:
  assumes Re z = 0 z ≠ 0
  shows arg z = pi/2 ∨ arg z = -pi/2
proof-
  have cos (arg z) = 0
  using assms
  by (metis Re-rcis no-zero-divisors norm-eq-zero rcis-cmod-arg)
thus ?thesis
  using arg-bounded[of z]
  using cos-0-iff-normalized[of arg z]
  by simp
qed

lemma im-complex-zero-arg1:
  assumes arg z = 0 ∨ arg z = pi
  shows Im z = 0
using assms
using rcis-cmod-arg[of z] Im-rcis[of cmod z arg z]
by auto

lemma im-complex-zero-arg2:
  assumes Im z = 0
  shows arg z = 0 ∨ arg z = pi
proof (cases z = 0)
  case True
  thus ?thesis
    by (auto simp add: arg-zero)
next
  case False
  hence sin (arg z) = 0
  using assms rcis-cmod-arg[of z] Im-rcis[of cmod z arg z]
  by auto
  thus ?thesis
    using arg-bounded[of z]
    using sin-0-iff-normalized
    by simp
qed

lemma arg-complex-of-real-positive:
  assumes k > 0
  shows arg (cor k) = 0
proof-
  have cos (arg (Complex k 0)) > 0
  using assms
  using rcis-cmod-arg[of Complex k 0] Re-rcis[of cmod (Complex k 0) arg (Complex k 0)]
  by auto
  thus ?thesis

```

```

using assms im-complex-zero-arg2[of cor k]
unfolding complex-of-real-def
by auto
qed

lemma arg-complex-of-real-negative:
assumes k < 0
shows arg (cor k) = pi
proof-
  have cos (arg (Complex k 0)) < 0
  using rcis-cmod-arg[of Complex k 0] Re-rcis[of cmod (Complex k 0) arg (Complex k 0)]
  by auto (metis assms less-asym' mult-eq-0-iff mult-pos-pos neqE zero-less-abs-iff)
  thus ?thesis
  using assms im-complex-zero-arg2[of cor k]
  unfolding complex-of-real-def
  by auto
qed

lemma
  [simp]: arg ii = pi/2
proof-
  have ii = cis (arg ii)
  using rcis-cmod-arg[of ii]
  by (simp add: rcis-def)
  hence cos (arg ii) = 0 sin (arg ii) = 1
  by (metis Re-cis complex-Re-i, metis Im-cis complex-Im-i)
  thus ?thesis
  using cos-0-iff-normalized[of arg ii] arg-bounded[of ii]
  by (auto simp add: field-simps)
qed

lemma
  [simp]: arg (-ii) = -pi/2
proof-
  have -ii = cis (arg (- ii))
  using rcis-cmod-arg[of -ii]
  by (simp add: rcis-def)
  hence cos (arg (-ii)) = 0 sin (arg (-ii)) = -1
  using Re-cis[of arg (-ii)] Im-cis[of arg (-ii)]
  by auto
  thus ?thesis
  using cos-0-iff-normalized[of arg (-ii)] arg-bounded[of -ii]
  by (metis one-neq-neg-numeral sin-pi-half)
qed

lemma arg-cis:
shows arg (cis φ) = |φ|
proof (rule canon-ang-eqI[symmetric])

```

```

show - pi < arg (cis φ) ∧ arg (cis φ) ≤ pi
  using arg-bounded
  by simp
next
  show ∃ k::int. arg (cis φ) - φ = 2*k*pi
  proof-
    have cis (arg (cis φ)) = cis φ
      using cis-arg[of cis φ]
      by auto
    thus ?thesis
      using cis-eq
      by auto
  qed
qed

lemma cos-arg:
  assumes z ≠ 0
  shows cos (arg z) = Re z / cmod z
  by (metis Complex.Re-sgn Re-cis assms cis-arg)

lemma sin-arg:
  assumes z ≠ 0
  shows sin (arg z) = Im z / cmod z
  by (metis Complex.Im-sgn Im-cis assms cis-arg)

lemma cis-arg-mult:
  assumes a * z ≠ 0
  shows cis (arg (a * z)) = cis (arg a + arg z)
  proof-
    have a * z = cor (cmod a) * cor (cmod z) * cis (arg a) * cis (arg z)
      using rcis-cmod-arg[of z, symmetric] rcis-cmod-arg[of a, symmetric]
      unfolding rcis-def
      by algebra
    hence a * z = cor (cmod (a * z)) * cis (arg a + arg z)
      using cis-mult[of arg a arg z]
      by auto
    hence cor (cmod (a * z)) * cis (arg a + arg z) = cor (cmod (a * z)) * cis (arg (a * z))
      using assms
      using rcis-cmod-arg[of a*z]
      unfolding rcis-def
      by auto
    thus ?thesis
      using mult-cancel-left[of cor (cmod (a * z)) cis (arg a + arg z) cis (arg (a * z))]
      using assms
      by auto
  qed

```

```

lemma arg-mult-2kpi:
  assumes a * z ≠ 0
  shows ∃ k:int. arg (a * z) = arg a + arg z + 2*k*pi
proof-
  have cis (arg (a*z)) = cis (arg a + arg z)
    by (rule cis-arg-mult[OF assms])
  thus ?thesis
    using cis-eq[of arg (a*z) arg a + arg z]
    by (auto simp add: field-simps)
qed

lemma arg-mult:
  assumes z1 * z2 ≠ 0
  shows arg(z1 * z2) = |arg z1 + arg z2|
proof-
  obtain k::int where arg(z1 * z2) = arg z1 + arg z2 + 2*k*pi
    using arg-mult-2kpi[of z1 z2]
    using assms
    by auto
  hence |arg(z1 * z2)| = |arg z1 + arg z2|
    using canon-ang-eq
    by(simp add:field-simps)
  thus ?thesis
    using canon-ang-arg[of z1*z2]
    by auto
qed

lemma arg-mult-real-positive:
  assumes k > 0
  shows arg (cor k * z) = arg z
proof (cases z = 0)
  case True
  thus ?thesis
    by (auto simp add: arg-zero)
next
  case False
  thus ?thesis
    using assms
    using arg-mult[of cor k z]
    by (auto simp add: arg-complex-of-real-positive canon-ang-arg)
qed

lemma arg-mult-real-negative:
  assumes k < 0
  shows arg (cor k * z) = arg (-z)
proof (cases z = 0)
  case True
  thus ?thesis

```

```

    by (auto simp add: arg-zero)
next
  case False
  thus ?thesis
    using assms
    using arg-mult[of cor k z]
    using arg-mult[of -1 z]
    using arg-complex-of-real-negative[of k] arg-complex-of-real-negative[of -1]
    by auto
qed

```

```

lemma arg-cnj1:
  assumes arg z = pi
  shows arg (cnj z) = pi
proof-
  have cos (arg (cnj z)) = cos (arg z)
    using rcis-cmod-arg[of z, symmetric] Re-rcis[of cmod z arg z]
    using rcis-cmod-arg[of cnj z, symmetric] Re-rcis[of cmod (cnj z) arg (cnj z)]
    by auto
  hence arg (cnj z) = arg z ∨ arg(cnj z) = -arg z
    using arg-bounded[of z] arg-bounded[of cnj z]
    by (metis arccos-cos arccos-cos2 less-eq-real-def linorder-le-cases minus-minus)
  thus ?thesis
    using assms
    using arg-bounded[of cnj z]
    by auto
qed

```

```

lemma arg-cnj2:
  assumes arg z ≠ pi
  shows arg (cnj z) = -arg z
proof(cases arg z = 0)
  case True
  thus ?thesis
    by (metis cnj-def complex-surj im-complex-zero-arg1 minus-zero)
next
  case False
  have cos (arg (cnj z)) = cos (arg z)
    using rcis-cmod-arg[of z] Re-rcis[of cmod z arg z]
    using rcis-cmod-arg[of cnj z] Re-rcis[of cmod (cnj z) arg (cnj z)]
    by auto
  hence arg (cnj z) = arg z ∨ arg(cnj z) = -arg z
    using arg-bounded[of z] arg-bounded[of cnj z]
    by (metis arccos-cos arccos-cos2 less-eq-real-def linorder-le-cases minus-minus)
moreover
have sin (arg (cnj z)) = -sin (arg z)
  using rcis-cmod-arg[of z] Im-rcis[of cmod z arg z]

```

```

using rcis-cmod-arg[of cnj z] Im-rcis[of cmod (cnj z) arg (cnj z)]
by auto (metis complex-Im-cnj complex-Im-zero complex-mod-cnj im-complex-zero-arg2
minus-mult-right norm-eq-zero real-mult-left-cancel sin-pi sin-zero)
hence arg (cnj z) ≠ arg z
using sin-0-iff-normalized[of arg (cnj z)] arg-bounded False assms
by auto
ultimately
show ?thesis
by auto
qed

lemma arg-div-real-positive:
assumes k ≠ 0 k > 0
shows arg (z / cor k) = arg z
proof(cases z = 0)
case True
thus ?thesis
by auto
next
case False
thus ?thesis
using assms
using arg-mult-real-positive[of 1/k z]
by auto
qed

lemma arg-inv1:
assumes z ≠ 0 arg z ≠ pi
shows arg (1 / z) = - arg z
proof-
have 1/z = cnj z / cor ((cmod z)2)
using ⟨z ≠ 0⟩ complex-mult-cnj-cmod[of z]
by (auto simp add:field-simps)
thus ?thesis
using arg-div-real-positive[of (cmod z)2 cnj z] ⟨z ≠ 0⟩
using arg-cnj2[of z] ⟨arg z ≠ pi⟩
by auto
qed

lemma arg-inv2:
assumes z ≠ 0 arg z = pi
shows arg (1 / z) = pi
proof-
have 1/z = cnj z / cor ((cmod z)2)
using ⟨z ≠ 0⟩ complex-mult-cnj-cmod[of z]
by (auto simp add:field-simps)
thus ?thesis
using arg-div-real-positive[of (cmod z)2 cnj z] ⟨z ≠ 0⟩

```

```

using arg-cnj1[of z]  $\langle \arg z = pi \rangle$ 
by auto
qed

lemma arg-inv-2kpi:
assumes  $z \neq 0$ 
shows  $\exists k:\text{int}. \arg(1/z) = -\arg z + 2*k*pi$ 
using arg-inv1[OF assms]
using arg-inv2[OF assms]
by (cases  $\arg z = pi$ ) (rule-tac  $x=1$  in exI, simp, rule-tac  $x=0$  in exI, simp)

lemma arg-inv:
assumes  $z \neq 0$ 
shows  $\arg(1/z) = \lfloor -\arg z \rfloor$ 
proof-
obtain  $k:\text{int}$  where  $\arg(1/z) = -\arg z + 2*k*pi$ 
using arg-inv-2kpi[of z]
using assms
by auto
hence  $\lfloor \arg(1/z) \rfloor = \lfloor -\arg z \rfloor$ 
using canon-ang-eq
by(simp add:field-simps)
thus ?thesis
using canon-ang-arg[of 1/z]
by auto
qed

lemma arg-div-2kpi:
assumes  $z1 \neq 0 z2 \neq 0$ 
shows  $\exists k:\text{int}. \arg(z1/z2) = \arg z1 - \arg z2 + 2*k*pi$ 
using assms
unfolding complex-divide-def[of z1 z2]
using inverse-eq-divide[of z2]
using arg-mult-2kpi[of z1 1/z2]
using arg-inv-2kpi[of z2]
by auto (metis comm-semiring-class.distrib distrib-left-numeral real-of-int-add)

lemma arg-div:
assumes  $z1 \neq 0 z2 \neq 0$ 
shows  $\arg(z1/z2) = \lfloor \arg z1 - \arg z2 \rfloor$ 
proof-
obtain  $k:\text{int}$  where  $\arg(z1/z2) = \arg z1 - \arg z2 + 2*k*pi$ 
using arg-div-2kpi[of z1 z2]
using assms
by auto
hence canon-ang( $\arg(z1/z2)$ ) = canon-ang( $\arg z1 - \arg z2$ )
using canon-ang-eq

```

```

by(simp add:field-simps)
thus ?thesis
  using canon-ang-arg[of z1/z2]
  by auto
qed

lemma arg-uminus:
  assumes z ≠ 0
  shows arg (-z) = |arg z + pi|
using assms
using arg-mult[of -1 z]
using arg-complex-of-real-negative[of -1]
by auto (metis comm-semiring-1-class.normalize-semiring-rules(24))

```

definition

$$\text{csqrt } z = \text{rcis} (\sqrt{(\text{cmod } z)}) (\arg z / 2)$$

lemma [simp]: $(\text{csqrt } x)^2 = x$
unfolding csqrt-def
by (subst DeMoivre2) (simp add: rcis-cmod-arg)

lemma ex-complex-sqrt: $\exists s::complex. s*s = z$
unfolding power2-eq-square[symmetric]
by (rule-tac x=csqrt z in exI) simp

lemma csqrt:
assumes $s * s = z$
shows $s = \text{csqrt } z \vee s = -\text{csqrt } z$
proof (cases $s = 0$)
case True
thus ?thesis
 using assms
 unfolding csqrt-def
 by simp
next
case False
then obtain k::int **where** cmod s * cmod s = cmod z 2 * arg s - arg z = 2*k*pi
 using assms
 using rcis-cmod-arg[of z] rcis-cmod-arg[of s]
 using arg-mult[of s s]
 using canon-ang(3)[of 2*arg s]
 by (auto simp add: norm-mult arg-mult)
have *: sqrt (cmod z) = cmod s

```

using ⟨cmod s * cmod s = cmod z⟩
by (smt norm-not-less-zero real-sqrt-abs2)

have **: arg z / 2 = arg s - k*pi
using ⟨2 * arg s - arg z = 2*k*pi⟩
by simp

have cis (arg s - k*pi) = cis (arg s) ∨ cis (arg s - k*pi) = -cis (arg s)
proof (cases even k)
  case True
  hence cis (arg s - k*pi) = cis (arg s)
    by (simp add: cis-def cos-diff sin-diff cos-kpi-even sin-kpi)
  thus ?thesis
    by simp
  next
    case False
    hence cis (arg s - k*pi) = -cis (arg s)
      by (simp add: cis-def cos-diff sin-diff cos-kpi-odd sin-kpi)
    thus ?thesis
      by simp
  qed
  thus ?thesis
proof
  assume ***: cis (arg s - real k * pi) = cis (arg s)
  hence s = csqrt z
    using rcis-cmod-arg[of s]
    unfolding csqrt-def rcis-def
    by (subst *, subst **, subst ***, simp)
  thus ?thesis
    by simp
  next
    assume ***: cis (arg s - real k * pi) = -cis (arg s)
    hence s = - csqrt z
      using rcis-cmod-arg[of s]
      unfolding csqrt-def rcis-def
      by (subst *, subst **, subst ***, simp)
    thus ?thesis
      by simp
  qed
qed

lemma [simp]: csqrt x = 0 ↔ x = 0
unfolding csqrt-def
by auto

lemma csqrt-mult: csqrt (a * b) = csqrt a * csqrt b ∨ csqrt (a * b) = - csqrt a
* csqrt b
proof (cases a = 0 ∨ b = 0)
  case True

```

```

thus ?thesis
  by auto
next
  case False
  obtain k::int where *:  $\lfloor \arg a + \arg b \rfloor = \arg a + \arg b - 2 * \text{real } k * \pi$ 
    using canon-ang(3)[of  $\arg a + \arg b$ ]
    by smt
  have cis ( $\lfloor \arg a + \arg b \rfloor / 2$ ) = cis ( $\arg a / 2 + \arg b / 2$ )  $\vee$  cis ( $\lfloor \arg a + \arg b \rfloor / 2$ ) = - cis ( $\arg a / 2 + \arg b / 2$ )
    using cos-kpi-even[of k] cos-kpi-odd[of k]
    by ((subst *)+, (subst diff-divide-distrib)+, (subst add-divide-distrib)+)
      (cases even k, auto simp add: cis-def cos-diff sin-diff sin-kpi)
  thus ?thesis
    using False
    unfolding csqrt-def
    by (simp add: rcis-mult real-sqrt-mult arg-mult)
      (auto simp add: rcis-def)
qed

lemma csqrt-real:
  assumes is-real x
  shows  $(\operatorname{Re} x \geq 0 \wedge \operatorname{csqrt} x = \operatorname{cor} (\sqrt{\operatorname{Re} x})) \vee (\operatorname{Re} x < 0 \wedge \operatorname{csqrt} x = i * \operatorname{cor} (\sqrt{-(\operatorname{Re} x)}))$ 
proof (cases x = 0)
  case True
  thus ?thesis
    by auto
next
  case False
  show ?thesis
  proof (cases  $\operatorname{Re} x > 0$ )
    case True
    hence arg x = 0
      using `is-real x`
      by (metis arg-complex-of-real-positive complex-of-real-Re)
    thus ?thesis
      using `Re x > 0`
      unfolding csqrt-def
      by simp (metis Re.simps complex-of-real-def rcis-cmod-arg rcis-zero-arg)
  next
    case False
    hence Re x < 0
      using `x ≠ 0` `is-real x`
      by (cases x, auto)
    hence arg x = pi
      using `is-real x`
      by (metis arg-complex-of-real-negative complex-of-real-Re)
    thus ?thesis
      using `Re x < 0`

```

```

unfolding csqrt-def
  by (simp add: rcis-def cis-def complex-of-real-def) (metis Complex-eq-0 False
    Re.simps assmss complex-minus-def complex-of-real-def cor-cmod-real le-less-linear
    norm-le-zero-iff)
  qed
qed

```

```

lemma is-real-rot-to-xaxis:
  assumes z ≠ 0
  shows is-real (cis (−arg z) * z)
proof (cases arg z = pi)
  case True
  thus ?thesis
    using im-complex-zero-arg1[of z]
    by auto
next
  case False
  hence |−arg z| = −arg z
  using canon-ang-eqI[of −arg z −arg z]
  using arg-bounded[of z]
  by (auto simp add: field-simps)
  hence arg (cis (−(arg z)) * z) = 0
  using arg-mult[of cis (−(arg z)) z] ⟨z ≠ 0⟩
  using arg-cis[of −arg z]
  by simp
  thus ?thesis
    using im-complex-zero-arg1[of cis (−arg z) * z]
    by auto
qed

```

```

lemma cmod-1-plus-mult-le:
  cmod (1 + z*w) ≤ sqrt((1 + (cmod z)^2) * (1 + (cmod w)^2))
proof–
  have Re ((1+z*w)*(1+cnj z*cnj w)) ≤ Re (1+z*cnj z)* Re (1+w*cnj w)
proof–
  have Re ((w − cnj z)*cnj(w − cnj z)) ≥ 0
  by (subst complex-mult-cnj-cmod) (simp add: power2-eq-square)
  hence Re (z*w + cnj z * cnj w) ≤ Re (w*cnj w) + Re(z*cnj z)
  by (simp only: complex-cnj complex-cnj-cnj field-simps complex-Re-diff complex-Re-add)
  thus ?thesis
    by (simp add: field-simps)
qed
hence (cmod (1 + z * w))^2 ≤ (1 + (cmod z)^2) * (1 + (cmod w)^2)
  by (subst cmod-square)+ simp

```

```

thus ?thesis
  by (metis abs-norm-cancel real-sqrt-abs real-sqrt-le-iff)
qed

lemma cmod-diff-ge: cmod (b - c) ≥ sqrt (1 + (cmod b)2) - sqrt (1 + (cmod c)2)
proof-
  have (cmod (b - c))2 + (1/2*Im(b*cnj c - c*cnj b))2 ≥ 0
    by simp
  hence (cmod (b - c))2 ≥ -(1/2*Im(b*cnj c - c*cnj b))2
    by simp
  hence (cmod (b - c))2 ≥ (1/2*Re(b*cnj c + c*cnj b))2 - Re(b*cnj b*c*cnj c)
    by (auto simp add: power2-eq-square field-simps)
  hence Re ((b - c)*(cnj b - cnj c)) ≥ (1/2*Re(b*cnj c + c*cnj b))2 - Re(b*cnj b*c*cnj c)
    by (subst (asm) cmod-square) (simp add: complex-cn)
  moreover
    have (1 + (cmod b)2) * (1 + (cmod c)2) = 1 + Re(b*cnj b) + Re(c*cnj c) + Re(b*cnj b*c*cnj c)
      by (subst cmod-square)+ (simp add: field-simps power2-eq-square)
  moreover
    have (1 + Re (scalprod b c))2 = 1 + 2*Re(scalprod b c) + ((Re (scalprod b c))2)
      by (subst power2-sum) simp
    hence (1 + Re (scalprod b c))2 = 1 + Re(b*cnj c + c*cnj b) + (1/2 * Re(b*cnj c + c*cnj b))
      by simp
  ultimately
    have (1 + (cmod b)2) * (1 + (cmod c)2) ≥ (1 + Re (scalprod b c))2
      by (simp add: field-simps)
  moreover
    have sqrt((1 + (cmod b)2) * (1 + (cmod c)2)) ≥ 0
      by (metis one-power2 real-sqrt-sum-squares-mult-ge-zero)
  ultimately
    have sqrt((1 + (cmod b)2) * (1 + (cmod c)2)) ≥ 1 + Re (scalprod b c)
      by (metis power2-le-imp-le real-sqrt-ge-0-iff real-sqrt-pow2-iff)
    hence Re ((b - c) * (cnj b - cnj c)) ≥ 1 + Re (c*cnj c) + 1 + Re (b*cnj b)
      - 2*sqrt((1 + (cmod b)2) * (1 + (cmod c)2))
      by (simp add: field-simps)
    hence *: (cmod (b - c))2 ≥ (sqrt (1 + (cmod b)2) - sqrt (1 + (cmod c)2))2
      apply (subst cmod-square)+
      apply (subst (asm) cmod-square)+
      apply (subst power2-diff)
      apply (subst real-sqrt-pow2, simp)
      apply (subst real-sqrt-pow2, simp)
      apply (simp add: real-sqrt-mult complex-cn)
      done
  thus ?thesis

```

```

proof (cases sqrt (1 + (cmod b)2) − sqrt (1 + (cmod c)2) > 0)
  case True
  thus ?thesis
    using square-cancel[OF *]
    by simp
next
  case False
  hence 0 ≥ sqrt (1 + (cmod b)2) − sqrt (1 + (cmod c)2)
    by (metis less_eq_real_def linorder_neqE_linordered_idom)
  moreover
  have cmod (b − c) ≥ 0
    by simp
  ultimately
  show ?thesis
    by (metis add_increasing monoid_add_class.add_right_neutral)
qed
qed

lemma cmod_diff_le: cmod (b − c) ≤ sqrt (1 + (cmod b)2) + sqrt (1 + (cmod c)2)
proof−
  have (cmod (b + c))2 + (1/2 * Im(b * cnj c − c * cnj b))2 ≥ 0
    by simp
  hence (cmod (b + c))2 ≥ −(1/2 * Im(b * cnj c − c * cnj b))2
    by simp
  hence (cmod (b + c))2 ≥ (1/2 * Re(b * cnj c + c * cnj b))2 − Re(b * cnj b * c * cnj c)
    by (auto simp add: power2_eq_square field_simps)
  hence Re ((b + c) * (cnj b + cnj c)) ≥ (1/2 * Re(b * cnj c + c * cnj b))2 − Re(b * cnj b * c * cnj c)
    by (subst (asm) cmod_square) (simp add: complex_cnj)
  moreover
  have (1 + (cmod b)2) * (1 + (cmod c)2) = 1 + Re(b * cnj b) + Re(c * cnj c) + Re(b * cnj b * c * cnj c)
    by (subst cmod_square) + (simp add: field_simps power2_eq_square)
  moreover
  have ++: 2 * Re(scalprod b c) = Re(b * cnj c + c * cnj b)
    by simp
  have (1 − Re (scalprod b c))2 = 1 − 2 * Re(scalprod b c) + ((Re (scalprod b c))2)
    by (subst power2_diff) simp
  hence (1 − Re (scalprod b c))2 = 1 − Re(b * cnj c + c * cnj b) + (1/2 * Re(b * cnj c + c * cnj b))2
    by (subst ++[symmetric]) simp
  ultimately
  have (1 + (cmod b)2) * (1 + (cmod c)2) ≥ (1 − Re (scalprod b c))2
    by (simp add: field_simps)
  moreover
  have sqrt((1 + (cmod b)2) * (1 + (cmod c)2)) ≥ 0

```

```

by (metis one-power2 real-sqrt-sum-squares-mult-ge-zero)
ultimately
have sqrt((1 + (cmod b)2) * (1 + (cmod c)2)) ≥ 1 - Re (scalprod b c)
  by (metis power2-le-imp-le real-sqrt-ge-0-iff real-sqrt-pow2-iff)
hence Re ((b - c) * (cnj b - cnj c)) ≤ 1 + Re (c*cnj c) + 1 + Re (b*cnj b)
+ 2*sqrt((1 + (cmod b)2) * (1 + (cmod c)2))
  by (simp add: field-simps)
hence *: (cmod (b - c))2 ≤ (sqrt (1 + (cmod b)2) + sqrt (1 + (cmod c)2))2
  apply (subst cmod-square)+
  apply (subst (asm) cmod-square)+
  apply (subst power2-sum)
  apply (subst real-sqrt-pow2, simp)
  apply (subst real-sqrt-pow2, simp)
  apply (simp add: real-sqrt-mult complex-cnj)
done
thus ?thesis
  using square-cancel[OF *]
  by simp
qed

```

```

definition cdist where
[simp]: cdist z1 z2 ≡ cmod (z2 - z1)

```

```

lemma [simp]:
fixes z1 z2 :: complex
assumes z1 ≠ 0 z2 ≠ 0
shows ∃ k. k ≠ 0 ∧ z2 = k * z1
using assms
by (rule-tac x=z2/z1 in exI) simp

```

```

lemma [simp]:
fixes z::complex
assumes z ≠ 0
shows ∃ k. k ≠ 0 ∧ k * z = 1
using assms
by (rule-tac x=1/z in exI) simp

```

```

lemma [simp]:
fixes z::complex
shows ∃ k. k ≠ 0 ∧ k * z = z
by (rule-tac x=1 in exI) simp

```

```

end

```

2 Systems of linear equations

```

theory LinearSystems
imports MoreComplex
begin

definition det2 where
[simp]: det2 a11 a12 a21 a22 ≡ a11*a22 - a12*a21

lemma regular-homogenous-system:
fixes a11::complex
assumes a11*a22 - a12*a21 ≠ 0 a11*x1 + a12*x2 = 0 a21*x1 + a22*x2 =
0
shows x1 = 0 ∧ x2 = 0
proof (cases a11 = 0)
case True
with assms(1) have a12 ≠ 0 a21 ≠ 0
by auto
thus ?thesis
using ⟨a11 = 0⟩ assms(2) assms(3)
by auto
next
case False
hence x1 = - a12*x2 / a11
using assms(2)
by (auto simp add: field-simps) (metis diff-divide-eq-iff diff-minus-eq-add divide-zero-left
eq-iff-diff-eq-0 minus-divide-left)
hence (a11*a22 - a12*a21)*x2 = 0
using assms(3) ⟨a11 ≠ 0⟩
by (auto simp add: field-simps)
thus ?thesis
using assms(1) assms(2) ⟨a11 ≠ 0⟩
by auto
qed

lemma regular-system:
fixes a11::complex
assumes a11*a22 - a12*a21 ≠ 0
shows ∃! x.
a11*(fst x) + a12*(snd x) = b1 ∧
a21*(fst x) + a22*(snd x) = b2
proof
let ?d = a11*a22 - a12*a21 and ?d1 = b1*a22 - b2*a12 and ?d2 = b2*a11
- b1*a21
let ?x = (?d1 / ?d, ?d2 / ?d)
have a11 * ?d1 + a12 * ?d2 = b1 * ?d a21 * ?d1 + a22 * ?d2 = b2 * ?d
by (auto simp add: field-simps)
thus a11 * fst ?x + a12 * snd ?x = b1 ∧ a21 * fst ?x + a22 * snd ?x = b2
using assms

```

```

by (metis (hide-lams, no-types) add-divide-distrib eq-divide-imp fst-eqD snd-eqD
times-divide-eq-right)

fix x'

$$\text{assume } a_{11} * \text{fst } x' + a_{12} * \text{snd } x' = b_1 \wedge a_{21} * \text{fst } x' + a_{22} * \text{snd } x' = b_2$$


$$\text{with } \langle a_{11} * \text{fst } ?x + a_{12} * \text{snd } ?x = b_1 \wedge a_{21} * \text{fst } ?x + a_{22} * \text{snd } ?x = b_2 \rangle$$


$$\text{have } a_{11} * (\text{fst } x' - \text{fst } ?x) + a_{12} * (\text{snd } x' - \text{snd } ?x) = 0 \wedge a_{21} * (\text{fst } x' -$$


$$\text{fst } ?x) + a_{22} * (\text{snd } x' - \text{snd } ?x) = 0$$

by (auto simp add: field-simps)
thus x' = ?x
using regular-homogenous-system[OF assms, of fst x' - fst ?x snd x' - snd ?x]
by (cases x') auto
qed

lemma singular-system:
fixes a11::complex
assumes a11*a22 - a12*a21 = 0 a11 ≠ 0 ∨ a12 ≠ 0
assumes *: a11*fst x0 + a12*snd x0 = b1 a21*fst x0 + a22*snd x0 = b2
assumes **: a11*fst x + a12*snd x = b1
shows a21*fst x + a22*snd x = b2
proof (cases a11 = 0)
case True
with assms have a21 = 0 a12 ≠ 0
by auto
let ?k = a22 / a12
have b2 = ?k * b1
using * ⟨a11 = 0⟩ ⟨a21 = 0⟩ ⟨a12 ≠ 0⟩
by auto
thus ?thesis
using ⟨a11 = 0⟩ ⟨a21 = 0⟩ ⟨a12 ≠ 0⟩ **
by auto
next
case False
let ?k = a21 / a11
from **
have ?k * a11 * fst x + ?k * a12 * snd x = ?k * b1
using ⟨a11 ≠ 0⟩
by (auto simp add: field-simps)
moreover
have a21 = ?k * a11 a22 = ?k * a12 b2 = ?k * b1
using assms(1) * ⟨a11 ≠ 0⟩
by (auto simp add: field-simps)
ultimately
show ?thesis
by auto
qed

lemma cnj-equation:
assumes a*z1 + b*z2 = c

```

```

shows  $\text{cnj } a * \text{cnj } z1 + \text{cnj } b * \text{cnj } z2 = \text{cnj } c$ 
using assms
by (auto simp add: complex-cnj-mult complex-cnj-add)

lemma regular-cnj-system:
assumes det2 a1 (cnj a1) a2 (cnj a2) ≠ 0 is-real b1 is-real b2
shows ∃! μ. a1 * cnj μ + cnj a1 * μ = b1 ∧
a2 * cnj μ + cnj a2 * μ = b2
proof-
have ∃! x. a1 * fst x + cnj a1 * snd x = b1 ∧
a2 * fst x + cnj a2 * snd x = b2
using regular-system assms(1)
by simp

then obtain x where
*: a1 * fst x + cnj a1 * snd x = b1
a2 * fst x + cnj a2 * snd x = b2
and **:
∀ x'. a1 * fst x' + cnj a1 * snd x' = b1 ∧
a2 * fst x' + cnj a2 * snd x' = b2 →
x' = x
unfolding Ex1-def
by blast
have cnj b1 = b1 cnj b2 = b2
using ⟨is-real b1⟩ ⟨is-real b2⟩
by (case-tac[!] b1, case-tac[!] b2) auto
hence a1 * cnj (snd x) + cnj a1 * cnj (fst x) = b1
a2 * cnj (snd x) + cnj a2 * cnj (fst x) = b2
using cnj-equation[OF *(1)] cnj-equation[OF *(2)] ⟨is-real b1⟩ ⟨is-real b2⟩
by (auto simp add: field-simps)
hence (cnj (snd x), cnj (fst x)) = x
using **
by auto
hence fst x = cnj (snd x)
by (cases x) auto
thus ?thesis
using **
unfolding Ex1-def
by (rule-tac x=snd x in exI, auto) (metis prod.inject)
qed

end

```

3 Quadratic equations

```

theory Quadratic
imports Complex MoreComplex
begin

```

```

lemma real-quadratic-equation:
  fixes  $\xi :: \text{real}$ 
  assumes  $\xi^2 + b * \xi + c = 0$   $b^2 - 4*c \geq 0$ 
  shows  $\xi = (-b + \sqrt{b^2 - 4*c}) / 2 \vee \xi = (-b - \sqrt{b^2 - 4*c}) / 2$ 
  using assms
  proof-
    from assms have  $(2 * (\xi + b/2))^2 = b^2 - 4*c$ 
    by (simp add: power2-eq-square field-simps)
    hence  $2 * (\xi + b/2) = \sqrt{b^2 - 4*c} \vee 2 * (\xi + b/2) = -\sqrt{b^2 - 4*c}$ 
    by (metis abs-minus-cancel power2-abs power2-eq-iff real-sqrt-abs)
    thus ?thesis
    by (auto simp add: field-simps)
  qed

lemma real-quadratic-equation':
  fixes  $\xi :: \text{real}$ 
  assumes  $b^2 - 4*c \geq 0$   $\xi = (-b + \sqrt{b^2 - 4*c}) / 2 \vee \xi = (-b - \sqrt{b^2 - 4*c}) / 2$ 
  shows  $\xi^2 + b * \xi + c = 0$ 
  using assms(2)
  proof
    assume  $*: \xi = (-b + \sqrt{b^2 - 4*c}) / 2$ 
    show ?thesis
    using assms(1)
    by ((subst *)+, subst power-divide, subst power2-sum, simp add: field-simps,
      simp add: power2-eq-square)
  next
    assume  $*: \xi = (-b - \sqrt{b^2 - 4*c}) / 2$ 
    show ?thesis
    using assms(1)
    by ((subst *)+, subst power-divide, subst power2-diff, simp add: field-simps,
      simp add: power2-eq-square)
  qed

lemma complex-quadratic-equation:
  fixes  $\xi :: \text{complex}$ 
  assumes  $\xi^2 + b * \xi + c = 0$ 
  shows  $\xi = (-b + \sqrt{b^2 - 4*c}) / 2 \vee \xi = (-b - \sqrt{b^2 - 4*c}) / 2$ 
  using assms
  proof-
    from assms have  $(2 * (\xi + b/2))^2 = b^2 - 4*c$ 
    by (simp add: power2-eq-square field-simps)
    (metis ab-semigroup-mult-class.mult-ac(1) comm-semiring-1-class.normalize-semiring-rules(34)
    comm-semiring-class.distrib mult-zero-left)
    hence  $2 * (\xi + b/2) = \sqrt{b^2 - 4*c} \vee 2 * (\xi + b/2) = -\sqrt{b^2 - 4*c}$ 
    using csqrt[of  $(2 * (\xi + b/2))$   $b^2 - 4*c$ ]
    by (simp add: power2-eq-square)
    thus ?thesis

```

```

using mult-cancel-right[of  $b + \xi * 2$  2 csqrt ( $b^2 - 4*c$ )]
using mult-cancel-right[of  $b + \xi * 2$  2 -csqrt ( $b^2 - 4*c$ )]
by (auto simp add: field-simps) (metis add-diff-cancel diff-minus-eq-add minus-diff-eq)
qed

lemma complex-quadratic-equation':
  fixes  $\xi :: complex$ 
  assumes  $\xi = (-b + csqrt(b^2 - 4*c)) / 2 \vee$ 
           $\xi = (-b - csqrt(b^2 - 4*c)) / 2$ 
  shows  $\xi^2 + b * \xi + c = 0$ 
  using assms
  proof
    assume *:  $\xi = (-b + csqrt(b^2 - 4*c)) / 2$ 
    show ?thesis
      by ((subst *)+) (subst power-divide, subst power2-sum, simp add: field-simps,
simp add: power2-eq-square)
    next
    assume *:  $\xi = (-b - csqrt(b^2 - 4*c)) / 2$ 
    show ?thesis
      by ((subst *)+, subst power-divide, subst power2-diff, simp add: field-simps,
simp add: power2-eq-square)
  qed

lemma complex-quadratic-equation-full:
  fixes  $\xi :: complex$ 
  assumes  $a * \xi^2 + b * \xi + c = 0$   $a \neq 0$ 
  shows  $\xi = (-b + csqrt(b^2 - 4*a*c)) / (2*a) \vee$ 
         $\xi = (-b - csqrt(b^2 - 4*a*c)) / (2*a)$ 
  proof-
    from assms have  $\xi^2 + (b/a) * \xi + (c/a) = 0$ 
      by (simp add: field-simps)
    hence  $\xi = (-(b/a) + csqrt((b/a)^2 - 4*(c/a))) / 2 \vee \xi = (-(b/a) - csqrt((b/a)^2$ 
 $- 4*(c/a))) / 2$ 
      using complex-quadratic-equation[of  $\xi b/a c/a$ ]
      by simp
    hence  $\exists k. \xi = (-(b/a) + (-1)^k * csqrt((b/a)^2 - 4*(c/a))) / 2$ 
      by safe (rule-tac x=2 in exI, simp, rule-tac x=1 in exI, simp)
    then obtain k1 where  $\xi = (-(b/a) + (-1)^{k1} * csqrt((b/a)^2 - 4*(c/a))) /$ 
 $2$ 
      by auto
    moreover
    have  $(b/a)^2 - 4 * (c/a) = (b^2 - 4 * a * c) * (1/a^2)$ 
      by (simp add: field-simps power2-eq-square)
    hence  $csqrt((b/a)^2 - 4 * (c/a)) = csqrt(b^2 - 4 * a * c) * csqrt(1/a^2) \vee$ 
 $csqrt((b/a)^2 - 4 * (c/a)) = -csqrt(b^2 - 4 * a * c) * csqrt(1/a^2)$ 
      using csqrt-mult[of  $b^2 - 4 * a * c 1/a^2$ ]
      by auto
    hence  $\exists k. csqrt((b/a)^2 - 4 * (c/a)) = (-1)^k * csqrt(b^2 - 4 * a * c)$ 
 $* csqrt(1/a^2)$ 
  
```

```

    by safe (rule-tac x=2 in exI, simp, rule-tac x=1 in exI, simp)
  then obtain k2 where csqrt ((b / a)2 - 4 * (c / a)) = (-1)k2 * csqrt (b2
- 4 * a * c) * csqrt (1 / a2)
    by auto
  moreover
  have csqrt (1 / a2) = 1/a ∨ csqrt (1 / a2) = -1/a
    using csqrt[of 1/a 1 / a2]
    by (auto simp add: power2-eq-square)
  hence ∃ k. csqrt (1 / a2) = (-1)k * 1/a
    by safe (rule-tac x=2 in exI, simp, rule-tac x=1 in exI, simp)
  then obtain k3 where csqrt (1 / a2) = (-1)k3 * 1/a
    by auto
  ultimately
  have ξ = -(b / a) + ((-1)k1 * (-1)k2 * (-1)k3) * csqrt (b2 - 4 *
a * c) * 1/a / 2
    by simp
  moreover
  have (-(1::complex))k1 * (-1)k2 * (-1)k3 = 1 ∨ (-(1::complex))k1 *
(-1)k2 * (-1)k3 = -1
    using neg-one-even-power[of k1 + k2 + k3]
    using neg-one-odd-power[of k1 + k2 + k3]
    by (simp add: comm-semiring-1-class.normalize-semiring-rules(26))
      (cases even (k1 + k2 + k3), auto)
  ultimately
  have ξ = -(b / a) + csqrt (b2 - 4 * a * c) * 1 / a / 2 ∨ ξ = -(b / a) -
csqrt (b2 - 4 * a * c) * 1 / a / 2
    by auto
  thus ?thesis
    using `a ≠ 0`
    by (simp add: field-simps)
qed

lemma complex-quadratic-two-solutions:
  fixes b c :: complex
  assumes b2 - 4*c ≠ 0
  shows ∃ k1 k2. k1 ≠ k2 ∧ k12 + b*k1 + c = 0 ∧ k22 + b*k2 + c = 0
proof-
  let ?ξ1 = (-b + csqrt(b2 - 4*c)) / 2
  let ?ξ2 = (-b - csqrt(b2 - 4*c)) / 2
  show ?thesis
    apply (rule-tac x=?ξ1 in exI)
    apply (rule-tac x=?ξ2 in exI)
    using assms complex-quadratic-equation'[of ?ξ1 b c] complex-quadratic-equation'[of
?ξ2 b c]
    by simp
qed

end

```

4 Vectors, Matrices

```
theory Matrices
imports MoreComplex LinearSystems Quadratic
begin
```

4.1 Vectors

Type of complex vector

```
type-synonym complex-vec = complex × complex
```

```
definition vec-zero :: complex-vec where
[simp]: vec-zero = (0, 0)
```

Vector scalar multiplication

```
fun mult-sv :: complex ⇒ complex-vec ⇒ complex-vec (infixl *sv 100) where
k *sv (x, y) = (k*x, k*y)
```

```
lemma fst-mult-sv [simp]: fst (k *sv v) = k * fst v
by (cases v) simp
```

```
lemma snd-mult-sv [simp]: snd (k *sv v) = k * snd v
by (cases v) simp
```

```
lemma mult-sv-mult-sv [simp]: k1 *sv (k2 *sv v) = (k1*k2) *sv v
by (cases v) simp
```

```
lemma one-mult-sv [simp]: 1 *sv v = v
by (cases v) simp
```

Multiplication of two vectors

```
fun mult-vv :: complex × complex ⇒ complex × complex ⇒ complex (infixl *vv
100) where
(x, y) *vv (a, b) = x*a + y*b
```

```
lemma mult-vv-commute: v1 *vv v2 = v2 *vv v1
by (cases v1, cases v2) auto
```

```
lemma mult-vv-scale-sv1:
(k *sv v1) *vv v2 = k * (v1 *vv v2)
by (cases v1, cases v2) (auto simp add: field-simps)
```

```
lemma mult-vv-scale-sv2:
v1 *vv (k *sv v2) = k * (v1 *vv v2)
by (cases v1, cases v2) (auto simp add: field-simps)
```

Conjugate vector

```
fun vec-map where
```

```

 $\text{vec-map } f \ (x, y) = (f x, f y)$ 

definition  $\text{vec-cnj}$  where  $\text{vec-cnj} = \text{vec-map cnj}$ 

lemma  $\text{vec-cnj-vec-cnj} [\text{simp}]: \text{vec-cnj} (\text{vec-cnj } v) = v$ 
by (cases v) (simp add: vec-cnj-def)

```

```

lemma  $\text{cnj-mult-vv}: \text{cnj} (v1 *_{vv} v2) = (\text{vec-cnj } v1) *_{vv} (\text{vec-cnj } v2)$ 
by (cases v1, cases v2) (simp add: vec-cnj-def complex-cnj)

```

```

lemma  $\text{vec-cnj-sv} [\text{simp}]: \text{vec-cnj} (k *_{sv} A) = \text{cnj } k *_{sv} \text{vec-cnj } A$ 
by (cases A) (auto simp add: vec-cnj-def complex-cnj)

```

```

lemma  $\text{scalsquare-vv-zero}:$ 
 $(\text{vec-cnj } v) *_{vv} v = 0 \longleftrightarrow v = \text{vec-zero}$ 
apply (cases v)
apply (auto simp add: vec-cnj-def field-simps complex-mult-cnj-cmod)
apply (smt norm-eq-zero of-real-add of-real-eq-0-iff of-real-power sum-power2-eq-zero-iff)+
done

```

4.2 Matrices

Type of complex matrices

```
type-synonym  $\text{complex-mat} = \text{complex} \times \text{complex} \times \text{complex} \times \text{complex}$ 
```

Matrix scalar multiplication

```
fun  $\text{mult-sm} :: \text{complex} \Rightarrow \text{complex-mat} \Rightarrow \text{complex-mat}$  (infixl  $*_{sm} 100$ ) where
 $k *_{sm} (a, b, c, d) = (k*a, k*b, k*c, k*d)$ 
```

```
lemma [simp]:  $k1 *_{sm} (k2 *_{sm} A) = (k1*k2) *_{sm} A$ 
by (cases A) auto
```

```
lemma [simp]:  $1 *_{sm} A = A$ 
by (cases A) auto
```

```
lemma  $\text{mult-sm-inv-l}:$ 
assumes  $k \neq 0$   $k *_{sm} A = B$ 
shows  $A = (1/k) *_{sm} B$ 
using assms
by auto
```

Matrix addition and subtraction

```
definition  $\text{mat-zero} :: \text{complex-mat}$  where [simp]:  $\text{mat-zero} = (0, 0, 0, 0)$ 
```

```
fun  $\text{mat-plus} :: \text{complex-mat} \Rightarrow \text{complex-mat} \Rightarrow \text{complex-mat}$  (infixl  $+_{mm} 100$ )
where
 $\text{mat-plus} (a1, b1, c1, d1) (a2, b2, c2, d2) = (a1+a2, b1+b2, c1+c2, d1+d2)$ 
```

```
fun mat-minus :: complex-mat  $\Rightarrow$  complex-mat  $\Rightarrow$  complex-mat (infixl  $-_{mm}$  100)
where
```

$$\text{mat-minus } (a_1, b_1, c_1, d_1) (a_2, b_2, c_2, d_2) = (a_1 - a_2, b_1 - b_2, c_1 - c_2, d_1 - d_2)$$

```
fun mat-uminus :: complex-mat  $\Rightarrow$  complex-mat where
```

$$\text{mat-uminus } (a, b, c, d) = (-a, -b, -c, -d)$$

```
lemma nonzero-mult-real:
```

```
assumes  $A \neq \text{mat-zero}$   $k \neq 0$ 
```

```
shows  $k *_{sm} A \neq \text{mat-zero}$ 
```

```
using assms
```

```
by (cases A) simp
```

Matrix multiplication

```
fun mult-mm :: complex-mat  $\Rightarrow$  complex-mat  $\Rightarrow$  complex-mat (infixl  $*_{mm}$  100)
where
```

$$(a_1, b_1, c_1, d_1) *_{mm} (a_2, b_2, c_2, d_2) =$$

$$(a_1*a_2 + b_1*c_2, a_1*b_2 + b_1*d_2, c_1*a_2 + d_1*c_2, c_1*b_2 + d_1*d_2)$$

```
lemma mult-mm-assoc:  $A *_{mm} (B *_{mm} C) = (A *_{mm} B) *_{mm} C$ 
by (cases A, cases B, cases C) (auto simp add: field-simps)
```

```
lemma mult-assoc-5:  $A *_{mm} (B *_{mm} C *_{mm} D) *_{mm} E = (A *_{mm} B) *_{mm} C$ 
 $*_{mm} (D *_{mm} E)$ 
```

```
by (simp only: mult-mm-assoc)
```

```
lemma mat-zero-r [simp]:  $A *_{mm} \text{mat-zero} = \text{mat-zero}$ 
by (cases A) simp
```

```
lemma mat-zero-l [simp]:  $\text{mat-zero} *_{mm} A = \text{mat-zero}$ 
by (cases A) simp
```

```
definition eye :: complex-mat where
[simp]: eye = (1, 0, 0, 1)
```

```
lemma mat-eye-l:
```

$$\text{eye} *_{mm} A = A$$

```
by (cases A) auto
```

```
lemma mat-eye-r:
```

$$A *_{mm} \text{eye} = A$$

```
by (cases A) auto
```

```
lemma mult-mm-sm [simp]:  $A *_{mm} (k *_{sm} B) = k *_{sm} (A *_{mm} B)$ 
```

```
by (cases A, cases B) (simp add: field-simps)
```

lemma *mult-sm-mm* [*simp*]: $(k *_{sm} A) *_{mm} B = k *_{sm} (A *_{mm} B)$
by (*cases A*, *cases B*) (*simp add: field-simps*)

lemma *mult-sm-eye-mm* [*simp*]: $k *_{sm} \text{eye} *_{mm} A = k *_{sm} A$
by (*cases A*) *simp*

Matrix determinant

fun *mat-det* **where** *mat-det* (*a, b, c, d*) = $a*d - b*c$

lemma *mat-det-mult* [*simp*]: *mat-det* (*A *_{mm} B*) = *mat-det A * mat-det B*
by (*cases A*, *cases B*) (*auto simp add: field-simps*)

lemma *mat-det-mult-sm* [*simp*]: *mat-det* (*k *_{sm} A*) = $(k*k) * \text{mat-det } A$
by (*cases A*) (*auto simp add: field-simps*)

Matrix inverse

fun *mat-inv* :: *complex-mat* \Rightarrow *complex-mat* **where**
mat-inv (*a, b, c, d*) = $(1/(a*d - b*c)) *_{sm} (d, -b, -c, a)$

lemma *mat-inv-r*:

assumes *mat-det A* $\neq 0$
shows *A *_{mm} (mat-inv A)* = *eye*
using *assms*
by (*cases A*, *auto simp add: field-simps*) *algebra*

lemma *mat-inv-l*:

assumes *mat-det A* $\neq 0$
shows *(mat-inv A) *_{mm} A* = *eye*
using *assms*
by (*cases A*, *auto simp add: field-simps*) *algebra*

lemma *mat-det-inv*:

assumes *mat-det A* $\neq 0$
shows *mat-det (mat-inv A)* = $1 / \text{mat-det } A$
proof –
have *mat-det eye* = *mat-det A * mat-det (mat-inv A)*
using *mat-inv-l[OF assms, symmetric]*
by *simp*
thus *?thesis*
using *assms*
by (*simp add: field-simps*)
qed

lemma *mult-mm-inv-l*:

assumes *mat-det A* $\neq 0$ *A *_{mm} B* = *C*
shows *B* = *mat-inv A *_{mm} C*
using *assms mat-eye-l[of B]*
by (*auto simp add: mult-mm-assoc mat-inv-l*)

```

lemma mult-mm-inv-r:
  assumes mat-det B ≠ 0 A *mm B = C
  shows A = C *mm mat-inv B
  using assms mat-eye-r[of A]
  by (auto simp add: mult-mm-assoc[symmetric] mat-inv-r)

lemma mult-mm-non-zero-l:
  assumes mat-det A ≠ 0 B ≠ mat-zero
  shows A *mm B ≠ mat-zero
  using assms mat-zero-r
  using mult-mm-inv-l[OF assms(1), of B mat-zero]
  by auto

lemma mat-inv-mult-mm:
  assumes mat-det A ≠ 0 mat-det B ≠ 0
  shows mat-inv (A *mm B) = mat-inv B *mm mat-inv A
  using assms
  proof-
    have (A *mm B) *mm (mat-inv B *mm mat-inv A) = eye
    using assms
    by (metis mat-inv-r mult-mm-assoc mult-mm-inv-r)
    thus ?thesis
      using mult-mm-inv-l[of A *mm B mat-inv B *mm mat-inv A eye] assms
      mat-eye-r
      by simp
  qed

lemma mult-mm-cancel-l:
  assumes mat-det M ≠ 0 M *mm A = M *mm B
  shows A = B
  using assms
  by (metis mult-mm-inv-l)

lemma mult-mm-cancel-r:
  assumes mat-det M ≠ 0 A *mm M = B *mm M
  shows A = B
  using assms
  by (metis mult-mm-inv-r)

lemma mult-mm-non-zero-r:
  assumes A ≠ mat-zero mat-det B ≠ 0
  shows A *mm B ≠ mat-zero
  using assms mat-zero-l
  using mult-mm-inv-r[OF assms(2), of A mat-zero]
  by auto

lemma mat-inv-mult-sm:
  assumes k ≠ 0

```

```

shows mat-inv ( $k *_{sm} A$ ) = ( $1 / k$ ) *sm mat-inv  $A$ 
proof-
  obtain a b c d where  $A = (a, b, c, d)$ 
    by (cases A) auto
  thus ?thesis
    using assms
    by auto (subst mult-assoc[of  $k a k*d$ ], subst mult-assoc[of  $k b k*c$ ], subst
right-diff-distrib[of  $k a*(k*d) b*(k*c)$ , symmetric], simp, simp add: field-simps)+
qed

lemma mat-inv-inv [simp]:
  assumes mat-det  $M \neq 0$ 
  shows mat-inv (mat-inv  $M$ ) =  $M$ 
proof-
  have mat-inv  $M *_{mm} M = eye$ 
  using mat-inv-l[OF assms]
  by simp
  thus ?thesis
    using assms mat-det-inv[of  $M$ ]
    using mult-mm-inv-l[of mat-inv  $M M eye$ ] mat-eye-r
    by (auto simp del: eye-def)
qed

```

Matrix transpose

```
fun mat-transpose where mat-transpose  $(a, b, c, d) = (a, c, b, d)$ 
```

```
lemma [simp]: mat-transpose (mat-transpose  $A$ ) =  $A$ 
by (cases A) auto
```

```
lemma [simp]: mat-transpose ( $k *_{sm} A$ ) =  $k *_{sm}$  (mat-transpose  $A$ )
by (cases A) simp
```

```
lemma [simp]: mat-transpose ( $A *_{mm} B$ ) = mat-transpose  $B *_{mm}$  mat-transpose
 $A$ 
by (cases A, cases B) auto
```

```
lemma mat-inv-transpose: mat-transpose (mat-inv  $M$ ) = mat-inv (mat-transpose
 $M$ )
by (cases M) auto
```

```
lemma mat-det-transpose:
  fixes  $M :: complex-mat$ 
  shows [simp]: mat-det (mat-transpose  $M$ ) = mat-det  $M$ 
by (cases M) auto
```

Diagonal matrices

```
fun mat-diagonal where
  mat-diagonal  $(A, B, C, D) = (B = 0 \wedge C = 0)$ 
```

Matrix conjugate

```

fun mat-map where
  mat-map f (a, b, c, d) = (f a, f b, f c, f d)

definition mat-cnj where mat-cnj = mat-map cnj

lemma [simp]: mat-cnj (mat-cnj A) = A
unfolding mat-cnj-def
by (cases A) auto

lemma mat-cnj-sm [simp]: mat-cnj (k *sm A) = cnj k *sm (mat-cnj A)
by (cases A) (simp add: mat-cnj-def complex-cnj)

lemma mat-det-cnj [simp]: mat-det (mat-cnj A) = cnj (mat-det A)
by (cases A) (simp add: mat-cnj-def complex-cnj)

lemma nonzero-mat-cnj: mat-cnj A = mat-zero  $\longleftrightarrow$  A = mat-zero
by (cases A) (auto simp add: mat-cnj-def)

lemma mat-inv-cnj: mat-cnj (mat-inv M) = mat-inv (mat-cnj M)
unfolding mat-cnj-def
by (cases M) (auto simp add: complex-cnj)

Matrix adjoint (conjugate)

definition mat-adj where mat-adj A = mat-cnj (mat-transpose A)

lemma mat-adj-mult-mm [simp]: mat-adj (A *mm B) = mat-adj B *mm mat-adj A
by (cases A, cases B) (auto simp add: mat-adj-def mat-cnj-def complex-cnj)

lemma mat-adj-mult-sm [simp]: mat-adj (k *sm A) = cnj k *sm mat-adj A
by (cases A) (auto simp add: mat-adj-def mat-cnj-def complex-cnj)

lemma mat-det-adj: mat-det (mat-adj A) = cnj (mat-det A)
by (cases A) (auto simp add: mat-adj-def mat-cnj-def complex-cnj)

lemma mat-adj-inv:
assumes mat-det M  $\neq$  0
shows mat-adj (mat-inv M) = mat-inv (mat-adj M)
by (cases M) (auto simp add: mat-adj-def mat-cnj-def complex-cnj)

lemma mat-transpose-mat-cnj: mat-transpose (mat-cnj A) = mat-adj A
by (cases A) (auto simp add: mat-adj-def mat-cnj-def)

lemma [simp]: mat-adj (mat-adj A) = A
unfolding mat-adj-def
by (subst mat-transpose-mat-cnj) (simp add: mat-adj-def)

Matrix trace

fun mat-trace where

```

mat-trace (a, b, c, d) = $a + d$

Multiplication of matrix and a vector

```
fun mult-mv :: complex-mat ⇒ complex-vec ⇒ complex-vec (infixl *mv 100) where
  (a, b, c, d) *mv (x, y) = (x*a + y*b, x*c + y*d)
```

```
fun mult-vm :: complex-vec ⇒ complex-mat ⇒ complex-vec (infixl *vm 100) where
  (x, y) *vm (a, b, c, d) = (x*a + y*c, x*b + y*d)
```

```
lemma eye-mv-l [simp]: eye *mv v = v
by (cases v) simp
```

```
lemma mult-mv-mv [simp]: B *mv (A *mv v) = (B *mm A) *mv v
by (cases v, cases A, cases B) (auto simp add: field-simps)
```

```
lemma mult-vm-vm [simp]: (v *vm A) *vm B = v *vm (A *mm B)
by (cases v, cases A, cases B) (auto simp add: field-simps)
```

```
lemma mult-mv-inv:
  assumes x = A *mv y mat-det A ≠ 0
  shows y = (mat-inv A) *mv x
using assms
by (cases y) (simp add: mat-inv-l)
```

```
lemma mult-vm-inv:
  assumes x = y *vm A mat-det A ≠ 0
  shows y = x *vm (mat-inv A)
using assms
by (cases y) (simp add: mat-inv-r)
```

```
lemma mult-mv-cancel-l:
  assumes mat-det A ≠ 0 A *mv v = A *mv v'
  shows v = v'
using assms
using mult-mv-inv
by blast
```

```
lemma mult-vm-cancel-r:
  assumes mat-det A ≠ 0 v *vm A = v' *vm A
  shows v = v'
using assms
using mult-vm-inv
by blast
```

```
lemma vec-zero-l [simp]:
  A *mv vec-zero = vec-zero
by (cases A) simp
```

```
lemma vec-zero-r [simp]:
```

```

vec-zero *vm A = vec-zero
by (cases A) simp

lemma mult-mv-nonzero:
assumes v ≠ vec-zero mat-det A ≠ 0
shows A *mv v ≠ vec-zero
apply (rule ccontr)
using assms mult-mv-inv[of vec-zero A v] mat-inv-l vec-zero-l
by auto

lemma mult-vm-nonzero:
assumes v ≠ vec-zero mat-det A ≠ 0
shows v *vm A ≠ vec-zero
apply (rule ccontr)
using assms mult-vm-inv[of vec-zero v A] mat-inv-r vec-zero-r
by auto

lemma mult-sv-mv: k *sv (A *mv v) = (A *mv (k *sv v))
by (cases A, cases v) (simp add: field-simps)

lemma mult-mv-mult-vm: A *mv x = x *vm (mat-transpose A)
by (cases A, cases x) auto

lemma mult-mv-vv: A *mv v1 *vv v2 = v1 *vv (mat-transpose A *mv v2)
by (cases v1, cases v2, cases A) (auto simp add: field-simps)

lemma mult-vv-mv: x *vv (A *mv y) = (x *vm A) *vv y
by (cases x, cases y, cases A) (auto simp add: field-simps)

lemma vec-cnj-mult-mv:
shows vec-cnj (A *mv x) = (mat-cnj A) *mv (vec-cnj x)
using assms
by (cases A, cases x) (auto simp add: vec-cnj-def mat-cnj-def complex-cnj)

lemma vec-cnj-mult-vm: vec-cnj (v *vm A) = vec-cnj v *vm mat-cnj A
unfolding vec-cnj-def mat-cnj-def
by (cases A, cases v, auto simp add: complex-cnj)

```

4.3 Eigenvalues and eigenvectors

```

definition eigenpair where
[simp]: eigenpair k v H ⟷ v ≠ vec-zero ∧ H *mv v = k *sv v

definition eigenval where
[simp]: eigenval k H ⟷ (∃ v. v ≠ vec-zero ∧ H *mv v = k *sv v)

lemma eigen-equation:
shows eigenval k H ⟷ k2 − mat-trace H * k + mat-det H = 0 (is ?lhs ⟷

```

```

?rhs)
proof-
  obtain A B C D where HH: H = (A, B, C, D)
    by (cases H) auto
  show ?thesis
  proof
    assume ?lhs
    then obtain v where v ≠ vec-zero H *mv v = k *sv v
      unfolding eigenval-def
      by blast
    obtain v1 v2 where vv: v = (v1, v2)
      by (cases v) auto
    from ‹H *mv v = k *sv v› have (H −mm (k *sm eye)) *mv v = vec-zero
      using HH vv
      by (auto simp add: field-simps)
    hence mat-det (H −mm (k *sm eye)) = 0
      using ‹v ≠ vec-zero› vv HH
      using regular-homogenous-system[of A − k D − k B C v1 v2]
      by (auto simp add: field-simps)
    thus ?rhs
      using HH
      by (auto simp add: power2-eq-square field-simps)
  next
    assume ?rhs
    hence *: mat-det (H −mm (k *sm eye)) = 0
      using HH
      by (auto simp add: field-simps power2-eq-square)
    show ?lhs
    proof (cases H −mm (k *sm eye) = mat-zero)
      case True
      thus ?thesis
        using HH
        by (auto) (rule-tac x=1 in exI, simp)
    next
      case False
      hence (A − k ≠ 0 ∨ B ≠ 0) ∨ (D − k ≠ 0 ∨ C ≠ 0)
        using HH
        by auto
      thus ?thesis
      proof
        assume A − k ≠ 0 ∨ B ≠ 0
        hence C * B + (D − k) * (k − A) = 0
          using * singular-system[of A − k D − k B C (0, 0) 0 0 (B, k − A)] HH
          by (auto simp add: field-simps)
        hence (B, k − A) ≠ vec-zero (H −mm (k *sm eye)) *mv (B, k − A) =
          vec-zero
          using HH ‹A − k ≠ 0 ∨ B ≠ 0›
          by (auto simp add: field-simps)
        then obtain v where v ≠ vec-zero ∧ (H −mm (k *sm eye)) *mv v =

```

```

 $\text{vec-zero}$ 
  by blast
  thus ?thesis
    using HH
    unfolding eigenval-def
    by (rule-tac x=v in exI) (case-tac v, simp add: field-simps)
next
  assume D - k ≠ 0 ∨ C ≠ 0
  hence C * B + (D - k) * (k - A) = 0
    using * singular-system[of D-k A-k C B (0, 0) 0 0 (C, k-D)] HH
    by (auto simp add: field-simps)
    hence (k-D, C) ≠ vec-zero (H -_mm (k *_sm eye)) *_mv (k-D, C) =
 $\text{vec-zero}$ 
  using HH {D - k ≠ 0 ∨ C ≠ 0}
  by (auto simp add: field-simps)
  then obtain v where v ≠ vec-zero ∧ (H -_mm (k *_sm eye)) *_mv v =
 $\text{vec-zero}$ 
  by blast
  thus ?thesis
    using HH
    unfolding eigenval-def
    by (rule-tac x=v in exI) (case-tac v, simp add: field-simps)
qed
qed
qed
qed

```

4.4 Bilinear and Quadratic forms; Congruence

Bilinear forms

definition bilinear-form **where**

[simp]: bilinear-form v1 v2 H = (vec-cnj v1) *_vm H *_vv v2

lemma bilinear-form-scale-m:

shows bilinear-form v1 v2 (k *_sm H) = k * bilinear-form v1 v2 H
by (cases v1, cases v2, cases H) (simp add: vec-cnj-def complex-cnj field-simps)

lemma bilinear-form-scale-v1:

shows bilinear-form (k *_sv v1) v2 H = cnj k * bilinear-form v1 v2 H
by (cases v1, cases v2, cases H) (simp add: vec-cnj-def complex-cnj field-simps)

lemma bilinear-form-scale-v2:

shows bilinear-form v1 (k *_sv v2) H = k * bilinear-form v1 v2 H
by (cases v1, cases v2, cases H) (simp add: vec-cnj-def complex-cnj field-simps)

Quadratic forms

definition quad-form **where**

[simp]: quad-form v H = (vec-cnj v) *_vm H *_vv v

```

lemma quad-form v H = bilinear-form v v H
by simp

lemma quad-form-scale-v:
  shows quad-form (k *sv v) H = cor ((cmod k)2) * quad-form v H
  using bilinear-form-scale-v1 bilinear-form-scale-v2
  by (simp add: complex-mult-cnj-cmod field-simps)

lemma quad-form-scale-m:
  shows quad-form v (k *sm H) = k * quad-form v H
  using bilinear-form-scale-m
  by simp

lemma cnj-quad-form [simp]: cnj (quad-form z H) = quad-form z (mat-adj H)
by (cases H, cases z) (auto simp add: mat-adj-def mat-cnj-def vec-cnj-def complex-cnj
field-simps)

```

Matrix congruence

```

abbreviation congruence where
  congruence M H ≡ mat-adj M *mm H *mm M

lemma bilinear-form-congruence:
  assumes mat-det M ≠ 0
  shows bilinear-form v1 v2 H = bilinear-form (M *mv v1) (M *mv v2) (congruence
(mat-inv M) H)
proof-
  have mat-det (mat-adj M) ≠ 0
  using assms
  by (simp add: mat-det-adj)
  show ?thesis
    unfolding bilinear-form-def
    apply (subst mult-mv-mult-vm)
    apply (subst vec-cnj-mult-vm)
    apply (subst mat-adj-def[symmetric])
    apply (subst mult-vm-vm)
    apply (subst mult-vv-mv)
    apply (subst mult-vm-vm)
    apply (subst mat-adj-inv[OF ⟨mat-det M ≠ 0⟩])
    apply (subst mult-assoc-5)
    apply (subst mat-inv-r[OF ⟨mat-det (mat-adj M) ≠ 0⟩])
    apply (subst mat-inv-l[OF ⟨mat-det M ≠ 0⟩])
    apply (subst mat-eye-l, subst mat-eye-r)
    by simp
  qed

lemma quad-form-congruence:
  assumes mat-det M ≠ 0
  shows quad-form (M *mv z) (congruence (mat-inv M) H) = quad-form z H
  using bilinear-form-congruence[OF assms]

```

by *simp*

lemma *congruence-nonzero*:

assumes $H \neq \text{mat-zero}$ mat-det $M \neq 0$

shows congruence $M H \neq \text{mat-zero}$

using *assms*

by (subst mult-mm-non-zero-r, subst mult-mm-non-zero-l) (auto simp add: mat-det-adj)

lemma *congruence-congruence*:

shows congruence $M1 (\text{congruence } M2 A) = \text{congruence } (M2 *_{mm} M1) A$

apply (subst mult-mm-assoc)

apply (subst mult-mm-assoc)

apply (subst mat-adj-mult-mm)

apply (subst mult-mm-assoc)

by *simp*

lemma [simp]: congruence eye $A = A$

by (cases A) (simp add: mat-adj-def mat-cnj-def)

lemma *congruence-congruence-inv*:

assumes mat-det $M \neq 0$

shows congruence $M (\text{congruence } (\text{mat-inv } M) A) = A$

using *assms congruence-congruence*[of M mat-inv $M A$]

using mat-inv-l[of M] mat-eye-l

by (simp del: eye-def)

lemma *congruence-inv*:

assumes mat-det $M \neq 0$ congruence $M A = B$

shows congruence (mat-inv M) $B = A$

using *assms*

using <mat-det $M \neq 0$ mult-mm-inv-l[of mat-adj $M A *_{mm} M B$]

using mult-mm-inv-r[of $M A$ mat-inv (mat-adj M) *_{mm} B]

by (simp add: mat-det-adj mult-mm-assoc mat-adj-inv)

lemma *congruence-scale-m*:

shows congruence $A (k *_{sm} B) = k *_{sm} (\text{congruence } A B)$

by (cases A , cases B) (auto simp add: mat-adj-def mat-cnj-def field-simps)

lemma *inj-congruence*:

assumes mat-det $M \neq 0$ congruence $M H = \text{congruence } M H'$

shows $H = H'$

proof –

have $H *_{mm} M = H' *_{mm} M$

using *assms*

using mult-mm-cancel-l[of mat-adj $M H *_{mm} M H' *_{mm} M$]

by (simp add: mat-det-adj mult-mm-assoc)

thus ?thesis

using *assms*

using mult-mm-cancel-r[of $M H H'$]

```

by simp
qed

definition similarity where similarity I M = mat-inv I *mm M *mm I

lemma mat-det-similarity:
assumes mat-det I ≠ 0
shows mat-det (similarity I M) = mat-det M
using assms
unfolding similarity-def
by (simp add: mat-det-inv)

lemma mat-trace-similarity:
assumes mat-det I ≠ 0
shows mat-trace (similarity I M) = mat-trace M
proof-
obtain a b c d where II: I = (a, b, c, d)
  by (cases I) auto
obtain A B C D where MM: M = (A, B, C, D)
  by (cases M) auto
have A * (a * d) / (a * d - b * c) + D * (a * d) / (a * d - b * c) =
  A + D + A * (b * c) / (a * d - b * c) + D * (b * c) / (a * d - b * c)
  using assms II
  by (simp add: field-simps)
thus ?thesis
  using II MM
  by (simp add: field-simps similarity-def)
qed

end

```

5 Unitary matrices

```

theory UnitaryMatrices
imports Matrices
begin

```

```

definition unitary where
unitary M ↔ mat-adj M *mm M = eye

definition unitary-gen where
unitary-gen M ↔ (∃ k::complex. k ≠ 0 ∧ mat-adj M *mm M = k *sm eye)

lemma unary-gen-scale [simp]:

```

```

assumes unitary-gen M k ≠ 0
shows unitary-gen (k *sm M)
using assms
unfolding unitary-gen-def
by auto

lemma unitary-unitary-gen [simp]: unitary M ⇒ unitary-gen M
unfolding unitary-gen-def unitary-def
by auto

lemma unitary-gen-real:
assumes unitary-gen M
shows (∃ k::real. k > 0 ∧ mat-adj M *mm M = cor k *sm eye)
proof-
obtain k where *: mat-adj M *mm M = k *sm eye k ≠ 0
using assms
by (auto simp add: unitary-gen-def)
obtain a b c d where M = (a, b, c, d)
by (cases M) auto
hence k = cor ((cmod a)2) + cor ((cmod c)2)
using *
by (subst complex-mult-cnj-cmod[symmetric])+
(auto simp add: mat-adj-def mat-cnj-def)
hence is-real k Re k > 0
using ‹k ≠ 0›
by (auto simp add: power2-eq-square) (metis comm-semiring-1-class.normalize-semiring-rules(6)
mult-eq-0-iff of-real-eq-0-iff sum-squares-gt-zero-iff)
thus ?thesis
using *
by (rule-tac x=Re k in exI) (simp add: complex-of-real-Re)
qed

lemma unitary-gen-regular:
assumes unitary-gen M
shows mat-det M ≠ 0
proof-
from assms obtain k where
k ≠ 0 mat-adj M *mm M = k *sm eye
unfolding unitary-gen-def
by auto
hence mat-det (mat-adj M *mm M) ≠ 0
by simp
thus ?thesis
by (simp add: mat-det-adj)
qed

```

lemmas unitary-regular = unitary-gen-regular[*OF* unitary-unitary-gen]

```

lemma unitary-gen M  $\longleftrightarrow$  ( $\exists k::complex. k \neq 0 \wedge mat-adj M *_{mm} (1, 0, 0, 1) *_{mm}$ 
 $M = k *_{sm} (1, 0, 0, 1)$ )
unfolding unitary-gen-def
using mat-eye-r
by (auto simp add: mult-assoc)

lemma unitary-comp:
assumes unitary M1 unitary M2
shows unitary (M1 *mm M2)
using assms
unfolding unitary-def
by (simp del: eye-def) (metis mat-eye-r mult-mm-assoc)

lemma unitary-gen-comp:
assumes unitary-gen M1 unitary-gen M2
shows unitary-gen (M1 *mm M2)
proof-
obtain k1 k2 where *:  $k1 * k2 \neq 0$  mat-adj M1 *mm M1 = k1 *sm eye mat-adj
M2 *mm M2 = k2 *sm eye
using assms
unfolding unitary-gen-def
by auto
have mat-adj M2 *mm mat-adj M1 *mm (M1 *mm M2) = mat-adj M2 *mm
(mat-adj M1 *mm M1) *mm M2
by (auto simp add: mult-mm-assoc)
also have ... = mat-adj M2 *mm ((k1 *sm eye) *mm M2)
using *
by (auto simp add: mult-mm-assoc)
also have ... = mat-adj M2 *mm (k1 *sm M2)
using mult-sm-eye-mm[of k1 M2]
by (simp del: eye-def)
also have ... = k1 *sm (k2 *sm eye)
using *
by auto
finally
show ?thesis
using *
unfolding unitary-gen-def
by (rule-tac x=k1*k2 in exI, simp del: eye-def)
qed

lemma unitary-adj-eq-inv:
unitary M  $\longleftrightarrow$  mat-det M  $\neq 0 \wedge mat-adj M = mat-inv M$ 
using unitary-regular[of M] mult-mm-inv-r[of M mat-adj M eye] mat-eye-l[of
mat-inv M] mat-inv-l[of M]
unfolding unitary-def
by – (rule, simp-all)

```

```

lemma unitary-inv:
  assumes unitary M
  shows unitary (mat-inv M)
using assms
unfolding unitary-adj-eq-inv
using mat-adj-inv[of M] mat-det-inv[of M]
by simp

lemma unitary-gen-unitary:
  shows unitary-gen M  $\longleftrightarrow$  ( $\exists k M'. k > 0 \wedge \text{unitary } M' \wedge M = (\text{cor } k *_{sm} \text{eye}) *_{mm} M'$ ) (is ?lhs = ?rhs)
proof
  assume ?lhs
  then obtain k where  $*: k > 0 \text{ mat-adj } M *_{mm} M = \text{cor } k *_{sm} \text{eye}$ 
  using unitary-gen-real[of M]
  by auto

  let ?k' = cor (sqrt k)
  have ?k' * cnj ?k' = cor k
    using ⟨k > 0⟩
    by simp
  moreover
  have Re ?k' > 0 is-real ?k' ?k' ≠ 0
    using ⟨k > 0⟩
    by auto
  ultimately
  show ?rhs
    using * mat-eye-l
    unfolding unitary-gen-def unitary-def
    by (rule-tac x=Re ?k' in exI) (rule-tac x=(1/?k')*sm M in exI, simp add: complex-cnj mult-sm-mm[symmetric])
next
  assume ?rhs
  then obtain k M' where  $k > 0 \text{ unitary } M' M = (\text{cor } k *_{sm} \text{eye}) *_{mm} M'$ 
  by blast
  hence  $M = \text{cor } k *_{sm} M'$ 
    using mult-sm-mm[of cor k eye M] mat-eye-l
    by simp
  thus ?lhs
    using ⟨unitary M'⟩ ⟨k > 0⟩
    by (simp add: unitary-gen-def unitary-def)
qed

lemma unitary-gen-inv:
  assumes unitary-gen M
  shows unitary-gen (mat-inv M)
proof-
  obtain k M' where  $0 < k \text{ unitary } M' M = \text{cor } k *_{sm} \text{eye} *_{mm} M'$ 

```

```

using unitary-gen-unitary[of M] assms
by blast
hence mat-inv M = cor (1/k) *sm mat-inv M'
  by (metis mat-inv-mult-sm mult-sm-eye-mm norm-not-less-zero of-real-1
of-real-divide of-real-eq-0-iff sgn-1-neg sgn-greater sgn-if sgn-pos sgn-sgn)
thus ?thesis
  using ‹k > 0› ⟨unitary M'⟩
  by (subst unitary-gen-unitary[of mat-inv M]) (rule-tac x=1/k in exI, rule-tac
x=mat-inv M' in exI, metis divide-pos-pos mult-sm-eye-mm unitary-inv zero-less-one)

```

qed

lemma unitary-special:

```

assumes unitary M mat-det M = 1
shows ∃ a b. M = (a, b, -cnj b, cnj a)

```

proof –

```

have mat-adj M = mat-inv M
  using assms mult-mm-inv-r[of M mat-adj M eye] mat-eye-r mat-eye-l
  by (simp add: unitary-def)
thus ?thesis
  using ⟨mat-det M = 1⟩
  by (cases M) (auto simp add: mat-adj-def mat-cnj-def)

```

qed

lemma unitary-gen-special:

```

assumes unitary-gen M mat-det M = 1
shows ∃ a b. M = (a, b, -cnj b, cnj a)

```

proof –

```

from assms
obtain k where *: k ≠ 0 mat-adj M *mm M = k *sm eye
  unfolding unitary-gen-def
  by auto
hence mat-det (mat-adj M *mm M) = k*k
  by simp
hence k*k = 1
  using assms(2)
  by (simp add: mat-det-adj)
hence k = 1 ∨ k = -1
  using square-eq-1-iff[of k]
  by simp

```

moreover

have mat-adj M = k *_{sm} mat-inv M

```

using *
using assms mult-mm-inv-r[of M mat-adj M k *sm eye] mat-eye-r mat-eye-l
by simp (metis mult-sm-eye-mm *(2))

```

moreover

obtain a b c d where M = (a, b, c, d)

by (cases M) auto

ultimately

```

have  $M = (a, b, -cnj b, cnj a) \vee M = (a, b, cnj b, -cnj a)$ 
  using assms(2)
  by (auto simp add: mat-adj-def mat-cnj-def)
moreover
have  $Re(-(\operatorname{cor}(\operatorname{cmod} a))^2 - (\operatorname{cor}(\operatorname{cmod} b))^2) < 1$ 
  by (auto simp add: power2-eq-square) (smt add-increasing2 add-nonneg-nonneg
is-num-normalize(8) less-le minus-add-distrib neg-le-0-iff-le norm-ge-zero norm-mult
not-one-le-zero real-0-le-add-iff zero-le-one)
hence  $-(\operatorname{cor}(\operatorname{cmod} a))^2 - (\operatorname{cor}(\operatorname{cmod} b))^2 \neq 1$ 
  by force
hence  $M \neq (a, b, cnj b, -cnj a)$ 
  using ⟨mat-det  $M = 1$ ⟩ complex-mult-cnj-cmod[of  $a$ ] complex-mult-cnj-cmod[of
 $b$ ]
  by auto
ultimately
show ?thesis
  by auto
qed

lemma unitary-gen-iff:
shows unitary-gen  $M \longleftrightarrow (\exists a b k . k \neq 0 \wedge \text{mat-det}(a, b, -cnj b, cnj a) \neq 0 \wedge (M = k *_{sm} (a, b, -cnj b, cnj a)))$  (is ?lhs = ?rhs)
proof
assume ?lhs
obtain d where *:  $d * d = \text{mat-det } M$ 
  using ex-complex-sqrt
  by auto
hence  $d \neq 0$ 
  using unitary-gen-regular[OF ⟨unitary-gen  $M$ ⟩]
  by auto
from ⟨unitary-gen  $M$ ⟩
obtain k where  $k \neq 0 \text{ mat-adj } M *_{mm} M = k *_{sm} \text{eye}$ 
  unfolding unitary-gen-def
  by auto
hence  $\text{mat-adj}((1/d) *_{sm} M) *_{mm} ((1/d) *_{sm} M) = (k / (d * cnj d)) *_{sm} \text{eye}$ 
  by (simp add: complex-cnj)
obtain a b where  $(a, b, -cnj b, cnj a) = (1 / d) *_{sm} M$ 
  using unitary-gen-special[of  $(1 / d) *_{sm} M$ ] ⟨unitary-gen  $M$ ⟩ * unitary-gen-regular[of
 $M$ ]  $d \neq 0$ 
  by force
moreover
hence  $\text{mat-det}(a, b, -cnj b, cnj a) \neq 0$ 
  using unitary-gen-regular[OF ⟨unitary-gen  $M$ ⟩]  $d \neq 0$ 
  by auto
ultimately
show ?rhs
apply (rule-tac x=a in exI, rule-tac x=b in exI, rule-tac x=d in exI)
using mult-sm-inv-l[of  $1/d M$ ]
by (auto simp add: field-simps)

```

```

next
assume ?rhs
then obtain a b k where k ≠ 0 ∧ mat-det (a, b, - cnj b, cnj a) ≠ 0 ∧ M =
k *sm (a, b, - cnj b, cnj a)
by auto
thus ?lhs
unfolding unitary-gen-def
apply (auto simp add: mat-adj-def mat-cnj-def complex-cnj)
using mult-eq-0-iff[of cnj k * k cnj a * a + cnj b * b]
by (auto simp add: field-simps)
qed

lemma unitary-iff:
shows unitary M  $\longleftrightarrow$ 
(∃ a b k . (cmod a)2 + (cmod b)2 ≠ 0 ∧ (cmod k)2 = 1 / ((cmod a)2 + (cmod
b)2) ∧ M = k *sm (a, b, -cnj b, cnj a)) (is ?lhs = ?rhs)
proof
assume ?lhs
obtain k a b where *: M = k *sm (a, b, -cnj b, cnj a) k ≠ 0 mat-det (a, b,
-cnj b, cnj a) ≠ 0
using unitary-gen-iff unitary-unitary-gen[OF ⟨unitary M⟩]
by auto

have md: mat-det (a, b, -cnj b, cnj a) = cor ((cmod a)2 + (cmod b)2)
by (auto simp add: complex-mult-cnj-cmod)

have k * cnj k * mat-det (a, b, -cnj b, cnj a) = 1
using ⟨unitary M⟩ *
unfolding unitary-def
by (auto simp add: mat-adj-def mat-cnj-def complex-cnj field-simps)
hence (cmod k)2 * ((cmod a)2 + (cmod b)2) = 1
by (subst (asm) complex-mult-cnj-cmod, subst (asm) md, subst (asm) cor-mult[symmetric])
(metis of-real-1 of-real-eq-iff)
thus ?rhs
using * mat-eye-l
apply (rule-tac x=a in exI, rule-tac x=b in exI, rule-tac x=k in exI)
apply (auto simp add: complex-mult-cnj-cmod)
by (metis `((cmod k)2 * ((cmod a)2 + (cmod b)2) = 1) mult-eq-0-iff nonzero-eq-divide-eq
zero-neq-one)
next
assume ?rhs
then obtain a b k where *: (cmod a)2 + (cmod b)2 ≠ 0 (cmod k)2 = 1 /
((cmod a)2 + (cmod b)2) M = k *sm (a, b, -cnj b, cnj a)
by auto
have (k * cnj k) * (a * cnj a) + (k * cnj k) * (b * cnj b) = 1
apply (subst complex-mult-cnj-cmod)+
using *(1-2)
apply (auto simp add: field-simps)
apply (metis cor-add cor-mult of-real-1 of-real-power)+
```

```

done
thus ?lhs
  using *
  unfolding unitary-def
  by (simp add: mat-adj-def mat-cnj-def complex-cnj field-simps)
qed

```

```

definition unitary11 where
  unitary11 M  $\longleftrightarrow$  mat-adj M *mm (1, 0, 0, -1) *mm M = (1, 0, 0, -1)

```

```

definition unitary11-gen where
  unitary11-gen M  $\longleftrightarrow$  ( $\exists$  k. k  $\neq$  0  $\wedge$  mat-adj M *mm (1, 0, 0, -1) *mm M = k *sm (1, 0, 0, -1))

```

```

lemma unitary11-gen-real:
  unitary11-gen M  $\longleftrightarrow$  ( $\exists$  k. k  $\neq$  0  $\wedge$  mat-adj M *mm (1, 0, 0, -1) *mm M = cor k *sm (1, 0, 0, -1))
  unfolding unitary11-gen-def
  proof auto
    fix k
    assume k  $\neq$  0 congruence M (1, 0, 0, -1) = (k, 0, 0, -k)
    hence mat-det (congruence M (1, 0, 0, -1)) = -k*k
      by simp
    moreover
      have is-real (mat-det (congruence M (1, 0, 0, -1))) Re (mat-det (congruence M (1, 0, 0, -1)))  $\leq$  0
        by (auto simp add: mat-det-adj) (smt real-minus-mult-self-le)
    ultimately
      have is-real (k*k) Re (-k*k)  $\leq$  0
        by auto
      hence is-real k
        using (k  $\neq$  0)
        by auto (smt not-real-square-gt-zero)
      thus  $\exists$  ka. ka  $\neq$  0  $\wedge$  k = cor ka
        using (k  $\neq$  0)
        by (rule-tac x=Re k in exI) (cases k, auto simp add: complex-of-real-Re)
qed

```

```

lemma unitary11-unitary11-gen [simp]: unitary11 M  $\implies$  unitary11-gen M
  unfolding unitary11-gen-def unitary11-def
  by (rule-tac x=1 in exI, auto)

```

```

lemma unitary11-gen-regular:
  assumes unitary11-gen M

```

shows $\text{mat-det } M \neq 0$
proof–
from assms **obtain** k **where**
 $k \neq 0 \text{ mat-adj } M *_{mm} (1, 0, 0, -1) *_{mm} M = \text{cor } k *_{sm} (1, 0, 0, -1)$
unfolding $\text{unitary11-gen-real}$
by auto
hence $\text{mat-det} (\text{mat-adj } M *_{mm} (1, 0, 0, -1) *_{mm} M) \neq 0$
by simp
thus ?thesis
by ($\text{simp add: mat-det-adj}$)
qed

lemmas $\text{unitary11-regular} = \text{unitary11-gen-regular}[\text{OF } \text{unitary11-unitary11-gen}]$

lemma $\text{unitary11-gen-mult-sm}:$
assumes $k \neq 0 \text{ unitary11-gen } M$
shows $\text{unitary11-gen} (k *_{sm} M)$
proof–
have $k * \text{cnj } k = \text{cor} (\text{Re} (k * \text{cnj } k))$
by ($\text{subst complex-of-real-Re}$) auto
thus ?thesis
using assms
unfolding $\text{unitary11-gen-real}$
by $\text{auto} (\text{rule-tac } x=\text{Re} (k * \text{cnj } k) * ka \text{ in } \text{exI}, \text{auto})$
qed

lemma $\text{unitary11-gen-div-sm}:$
assumes $k \neq 0 \text{ unitary11-gen } (k *_{sm} M)$
shows $\text{unitary11-gen } M$
using $\text{assms unitary11-gen-mult-sm}[\text{of } 1/k k *_{sm} M]$
by simp

lemma $\text{unitary11-special}:$
assumes $\text{unitary11 } M \text{ mat-det } M = 1$
shows $\exists a b. M = (a, b, \text{cnj } b, \text{cnj } a)$
proof–
have $\text{mat-adj } M *_{mm} (1, 0, 0, -1) = (1, 0, 0, -1) *_{mm} \text{mat-inv } M$
using $\text{assms mult-mm-inv-r}$
by ($\text{simp add: unitary11-def}$)
thus ?thesis
using $\text{assms}(2)$
by ($\text{cases } M$) ($\text{simp add: mat-adj-def mat-cnj-def}$)
qed

lemma $\text{unitary11-gen-special}:$
assumes $\text{unitary11-gen } M \text{ mat-det } M = 1$
shows $\exists a b. M = (a, b, \text{cnj } b, \text{cnj } a) \vee M = (a, b, -\text{cnj } b, -\text{cnj } a)$
proof–
from assms

```

obtain k where *:  $k \neq 0$  mat-adj  $M *_{mm} (1, 0, 0, -1) *_{mm} M = cor k *_{sm}$ 
 $(1, 0, 0, -1)$ 
  unfolding unitary11-gen-real
  by auto
hence mat-det (mat-adj  $M *_{mm} (1, 0, 0, -1) *_{mm} M$ ) =  $- cor k * cor k$ 
  by simp
hence mat-det (mat-adj  $M *_{mm} M$ ) =  $cor k * cor k$ 
  by simp
hence  $cor k * cor k = 1$ 
  using assms(2)
  by (simp add: mat-det-adj)
hence  $cor k = 1 \vee cor k = -1$ 
  using square-eq-1-iff[of cor k]
  by simp
moreover
have mat-adj  $M *_{mm} (1, 0, 0, -1) = (cor k *_{sm} (1, 0, 0, -1)) *_{mm} mat-inv$ 
 $M$ 
  using *
  using assms mult-mm-inv-r mat-eye-r mat-eye-l
  by auto
moreover
obtain a b c d where  $M = (a, b, c, d)$ 
  by (cases M) auto
ultimately
have  $M = (a, b, cnj b, cnj a) \vee M = (a, b, -cnj b, -cnj a)$ 
  using assms(2)
  by (auto simp add: mat-adj-def mat-cnj-def)
thus ?thesis
  by auto
qed

lemma unitary11-gen-iff':
  shows unitary11-gen  $M \longleftrightarrow$ 
     $(\exists a b k . k \neq 0 \wedge mat-det (a, b, cnj b, cnj a) \neq 0 \wedge$ 
       $(M = k *_{sm} (a, b, cnj b, cnj a) \vee M = k *_{sm} (-1, 0, 0, 1) *_{mm} (a,$ 
       $b, cnj b, cnj a)))$  (is ?lhs = ?rhs)
proof
  assume ?lhs
  obtain d where *:  $d * d = mat-det M$ 
    using ex-complex-sqrt
    by auto
  hence  $d \neq 0$ 
    using unitary11-gen-regular[OF `unitary11-gen M`]
    by auto
  from `unitary11-gen M`
  obtain k where  $k \neq 0$  mat-adj  $M *_{mm} (1, 0, 0, -1) *_{mm} M = cor k *_{sm} (1,$ 
 $0, 0, -1)$ 
    unfolding unitary11-gen-real
    by auto

```

```

hence mat-adj ((1/d)*smM)*mm (1, 0, 0, -1) *mm ((1/d)*smM) = (cor k / (d*cnj d)) *sm (1, 0, 0, -1)
  by (simp add: complex-cnj)
moreover
  have is-real (cor k / (d * cnj d))
    by (metis complex-In-mult-cnj-zero div-reals is-real-complex-of-real)
  hence cor (Re (cor k / (d * cnj d))) = cor k / (d * cnj d)
    by (simp add: complex-of-real-Re)
ultimately
  have unitary11-gen ((1/d)*smM)
    unfolding unitary11-gen-real
    using ⟨d ≠ 0⟩ ⟨k ≠ 0⟩
    by (rule-tac x=Re (cor k / (d * cnj d)) in exI, auto)
moreover
  have mat-det ((1 / d) *sm M) = 1
    using * unitary11-gen-regular[of M] ⟨unitary11-gen M⟩
    by auto
ultimately
  obtain a b where (a, b, cnj b, cnj a) = (1 / d) *sm M ∨ (a, b, -cnj b, -cnj a) = (1 / d) *sm M
    using unitary11-gen-special[of (1 / d) *sm M]
    by force
  thus ?rhs
proof
  assume (a, b, cnj b, cnj a) = (1 / d) *sm M
  moreover
  hence mat-det (a, b, cnj b, cnj a) ≠ 0
    using unitary11-gen-regular[OF ⟨unitary11-gen M⟩] ⟨d ≠ 0⟩
    by auto
  ultimately
  show ?rhs
    using ⟨d ≠ 0⟩
    by (rule-tac x=a in exI, rule-tac x=b in exI, rule-tac x=d in exI, simp)
next
  assume*: (a, b, -cnj b, -cnj a) = (1 / d) *sm M
  hence (1 / d) *sm M = (a, b, -cnj b, -cnj a)
    by simp
  hence M = (a * d, b * d, - (d * cnj b), - (d * cnj a))
    using ⟨d ≠ 0⟩
    using mult-sm-inv-l[of 1/d M (a, b, -cnj b, -cnj a), symmetric]
    by (simp add: field-simps)
  moreover
  have mat-det (a, b, -cnj b, -cnj a) ≠ 0
    using* unitary11-gen-regular[OF ⟨unitary11-gen M⟩] ⟨d ≠ 0⟩
    by auto
  ultimately
  show ?thesis
    using ⟨d ≠ 0⟩
    by (rule-tac x=a in exI, rule-tac x=b in exI, rule-tac x=-d in exI) (simp)

```

```

add: field-simps)
qed
next
assume ?rhs
then obtain a b k where k ≠ 0 mat-det (a, b, cnj b, cnj a) ≠ 0
M = k *sm (a, b, cnj b, cnj a) ∨ M = k *sm (-1, 0, 0, 1) *mm (a, b, cnj b,
cnj a)
by auto
moreover
let ?x = cnj k * cnj a * (k * a) + - (cnj k * b * (k * cnj b))
have ?x = (k*cnj k)*(a*cnj a - b*cnj b)
by (auto simp add: field-simps)
hence is-real ?x
by simp
hence cor (Re ?x) = ?x
by (rule complex-of-real-Re)
moreover
have ?x ≠ 0
using mult-eq-0-iff[of cnj k * k (cnj a * a + - cnj b * b)]
using ⟨mat-det (a, b, cnj b, cnj a) ≠ 0⟩ ⟨k ≠ 0⟩
by (auto simp add: field-simps)
hence Re ?x ≠ 0
using ⟨is-real ?x⟩
by (cases ?x) simp
ultimately
show ?lhs
unfolding unitary11-gen-real
by (auto simp add: mat-adj-def mat-cnj-def complex-cnj)
qed

```

```

lemma unitary11-gen-cis-blaschke:
assumes k ≠ 0 M = k *sm (a, b, cnj b, cnj a) a ≠ 0 mat-det (a, b, cnj b, cnj
a) ≠ 0
shows ∃ k' φ a'. k' ≠ 0 ∧ a' * cnj a' ≠ 1 ∧ M = k' *sm (cis φ, 0, 0, 1) *mm
(1, -a', -cnj a', 1)
proof-
have a = cnj a * cis (2 * arg a)
using rcis-cmod-arg[of a] rcis-cnj[of a]
using cis-rcis-eq rcis-mult
by simp
thus ?thesis
using assms
by (rule-tac x=k*cnj a in exI, rule-tac x=2*arg a in exI, rule-tac x=- b / a
in exI) (auto simp add: field-simps complex-cnj)
qed

```

lemma unitary11-gen-cis-blaschke':

assumes $k \neq 0 M = k *_{sm} (-1, 0, 0, 1) *_{mm} (a, b, cnj b, cnj a)$ $a \neq 0 mat-det$
 $(a, b, cnj b, cnj a) \neq 0$
shows $\exists k' \varphi a'. k' \neq 0 \wedge a' * cnj a' \neq 1 \wedge M = k' *_{sm} (cis \varphi, 0, 0, 1) *_{mm}$
 $(1, -a', -cnj a', 1)$
proof–
obtain $k' \varphi a'$ **where** $*: k' \neq 0 k *_{sm} (a, b, cnj b, cnj a) = k' *_{sm} (cis \varphi, 0,$
 $0, 1) *_{mm} (1, -a', -cnj a', 1) a' * cnj a' \neq 1$
using *unitary11-gen-cis-blaschke*[OF $\langle k \neq 0 \rangle - \langle a \neq 0 \rangle$] [$mat-det (a, b, cnj b,$
 $cnj a) \neq 0$]
by *blast*
have $(cis \varphi, 0, 0, 1) *_{mm} (-1, 0, 0, 1) = (cis (\varphi + pi), 0, 0, 1)$
by (*simp add: cis-def*)
thus *?thesis*
using $* \langle M = k *_{sm} (-1, 0, 0, 1) *_{mm} (a, b, cnj b, cnj a) \rangle$
by (*rule-tac x=k' in exI, rule-tac x=varphi+pi in exI, rule-tac x=a' in exI, simp*)
 $(metis minus-mult-right equation-minus-iff minus-mult-left minus-mult-right)$
qed

lemma *unitary11-gen-cis-blaschke-rev*:
assumes $k' \neq 0 M = k' *_{sm} (cis \varphi, 0, 0, 1) *_{mm} (1, -a', -cnj a', 1) a' * cnj$
 $a' \neq 1$
shows $\exists k a b. k \neq 0 \wedge mat-det (a, b, cnj b, cnj a) \neq 0 \wedge M = k *_{sm} (a, b,$
 $cnj b, cnj a)$
using *assms*
by (*rule-tac x=k'*cis(varphi/2) in exI, rule-tac x=cis(varphi/2) in exI, rule-tac x=-a'*cis(varphi/2)*
in exI) (*simp add: complex-cnj cis-mult, simp add: cis-def*)

lemma *unitary11-gen-cis-inversion*:
assumes $k \neq 0 M = k *_{sm} (0, b, cnj b, 0) b \neq 0$
shows $\exists k' \varphi. k' \neq 0 \wedge M = k' *_{sm} (cis \varphi, 0, 0, 1) *_{mm} (0, 1, 1, 0)$
using *assms*
using *rcis-cmod-arg*[of b , *symmetric*] *rcis-cnj*[of b] *cis-rcis-eq*
by (*simp (rule-tac x=2*arg b in exI, simp add: rcis-mult)*)

lemma *unitary11-gen-cis-inversion'*:
assumes $k \neq 0 M = k *_{sm} (-1, 0, 0, 1) *_{mm} (0, b, cnj b, 0) b \neq 0$
shows $\exists k' \varphi. k' \neq 0 \wedge M = k' *_{sm} (cis \varphi, 0, 0, 1) *_{mm} (0, 1, 1, 0)$
proof–
obtain $k' \varphi$ **where** $*: k' \neq 0 k *_{sm} (0, b, cnj b, 0) = k' *_{sm} (cis \varphi, 0, 0, 1)$
 $*_{mm} (0, 1, 1, 0)$
using *unitary11-gen-cis-inversion*[OF $\langle k \neq 0 \rangle - \langle b \neq 0 \rangle$]
by *metis*
have $(cis \varphi, 0, 0, 1) *_{mm} (-1, 0, 0, 1) = (cis (\varphi + pi), 0, 0, 1)$
by (*simp add: cis-def*)
thus *?thesis*
using $* \langle M = k *_{sm} (-1, 0, 0, 1) *_{mm} (0, b, cnj b, 0) \rangle$
by (*rule-tac x=k' in exI, rule-tac x=varphi+pi in exI, simp*)
 $(metis minus-mult-right)$
qed

```

lemma unitary11-gen-cis-inversion-rev:
  assumes  $k' \neq 0$   $M = k' *_{sm} (\text{cis } \varphi, 0, 0, 1) *_{mm} (0, 1, 1, 0)$ 
  shows  $\exists k a b. k \neq 0 \wedge \text{mat-det}(a, b, \text{cnj } b, \text{cnj } a) \neq 0 \wedge M = k *_{sm} (a, b, \text{cnj } b, \text{cnj } a)$ 
  using assms
  by (rule-tac  $x=k'*\text{cis}(\varphi/2)$  in exI, rule-tac  $x=0$  in exI, rule-tac  $x=\text{cis}(\varphi/2)$  in exI) (simp add: cis-mult, simp add: cis-def)

lemma unitary11-gen-iff:
  shows unitary11-gen  $M \longleftrightarrow (\exists k a b. k \neq 0 \wedge \text{mat-det}(a, b, \text{cnj } b, \text{cnj } a) \neq 0 \wedge M = k *_{sm} (a, b, \text{cnj } b, \text{cnj } a))$  (is ?lhs = ?rhs)
  proof
    assume ?lhs
    then obtain a b k where  $*: k \neq 0 \text{ mat-det}(a, b, \text{cnj } b, \text{cnj } a) \neq 0 M = k *_{sm} (a, b, \text{cnj } b, \text{cnj } a) \vee M = k *_{sm} (-1, 0, 0, 1) *_{mm} (a, b, \text{cnj } b, \text{cnj } a)$ 
    using unitary11-gen-iff'
    by auto
    show ?rhs
    proof (cases  $M = k *_{sm} (a, b, \text{cnj } b, \text{cnj } a)$ )
      case True
      thus ?thesis
        using *
        by auto
    next
      case False
      hence **:  $M = k *_{sm} (-1, 0, 0, 1) *_{mm} (a, b, \text{cnj } b, \text{cnj } a)$ 
      using *
      by simp
      show ?thesis
      proof (cases  $a = 0$ )
        case True
        hence  $b \neq 0$ 
        using *
        by auto
        show ?thesis
        using unitary11-gen-cis-inversion-rev[of - M]
        using ** ⟨ $a = 0$ ⟩
        using unitary11-gen-cis-inversion'[OF ⟨ $k \neq 0$ ⟩ - ⟨ $b \neq 0$ ⟩, of M]
        by auto
    next
      case False
      show ?thesis
      using unitary11-gen-cis-blaschke-rev[of - M]
      using **
      using unitary11-gen-cis-blaschke'[OF ⟨ $k \neq 0$ ⟩ - ⟨ $a \neq 0$ ⟩, of M b] ⟨mat-det
       $(a, b, \text{cnj } b, \text{cnj } a) \neq 0$ ⟩

```

```

    by blast
qed
qed
next
assume ?rhs
thus ?lhs
  using unitary11-gen-iff'
  by auto
qed

lemma unitary11-iff:
  shows unitary11 M  $\longleftrightarrow$ 
     $(\exists a b k . (cmod a)^2 > (cmod b)^2 \wedge (cmod k)^2 = 1 / ((cmod a)^2 - (cmod b)^2)$ 
 $\wedge M = k *_{sm} (a, b, cnj b, cnj a)) (\text{is } ?lhs = ?rhs)$ 
proof
  assume ?lhs
  obtain k a b where *:
     $M = k *_{sm} (a, b, cnj b, cnj a) \text{mat-det } (a, b, cnj b, cnj a) \neq 0 k \neq 0$ 
    using unitary11-gen-iff unitary11-unitary11-gen[OF `unitary11 M`]
    by auto

  have md: mat-det (a, b, cnj b, cnj a) = cor ((cmod a)^2 - (cmod b)^2)
    by (auto simp add: complex-mult-cnj-cmod)
  hence **:  $(cmod a)^2 \neq (cmod b)^2$ 
    using `mat-det (a, b, cnj b, cnj a) \neq 0`
    by auto (metis of-real-power)

  have k * cnj k * mat-det (a, b, cnj b, cnj a) = 1
    using `M = k *_{sm} (a, b, cnj b, cnj a)`
    using `unitary11 M`
    unfolding unitary11-def
    by (auto simp add: mat-adj-def mat-cnj-def complex-cnj) (simp add: field-simps)
  hence  $(cmod k)^2 * ((cmod a)^2 - (cmod b)^2) = 1$ 
    by (subst (asm) complex-mult-cnj-cmod, subst (asm) md, subst (asm) cor-mult[symmetric])
      (metis of-real-1 of-real-eq-iff)
  thus ?rhs
    using `M = k *_{sm} (a, b, cnj b, cnj a)` ** mat-eye-l
    apply (rule-tac x=a in exI, rule-tac x=b in exI, rule-tac x=k in exI)
    apply (auto simp add: complex-mult-cnj-cmod)
    apply (metis less-iff-diff-less-0 linorder-neqE-linordered-idom mult-pow2-lt0
      mult-zero-left not-one-less-zero zero-eq-power2 zero-neq-one)
    apply (metis `(cmod k)^2 * ((cmod a)^2 - (cmod b)^2) = 1` mult-eq-0-iff nonzero-eq-divide-eq
      zero-neq-one)
    done
next
assume ?rhs
then obtain a b k where  $(cmod b)^2 < (cmod a)^2 \wedge (cmod k)^2 = 1 / ((cmod a)^2 - (cmod b)^2)$ 
   $\wedge M = k *_{sm} (a, b, cnj b, cnj a)$ 
  by auto

```

```

moreover
have  $\text{cnj } k * \text{cnj } a * (k * a) + - (\text{cnj } k * b * (k * \text{cnj } b)) = (\text{cor } ((\text{cmod } k)^2 * ((\text{cmod } a)^2 - (\text{cmod } b)^2)))$ 
proof-
have  $\text{cnj } k * \text{cnj } a * (k * a) = \text{cor } ((\text{cmod } k)^2 * (\text{cmod } a)^2)$ 
  using complex-mult-cnj-cmod[of a] complex-mult-cnj-cmod[of k]
  by (auto simp add: field-simps)
moreover
have  $\text{cnj } k * b * (k * \text{cnj } b) = \text{cor } ((\text{cmod } k)^2 * (\text{cmod } b)^2)$ 
  using complex-mult-cnj-cmod[of b, symmetric] complex-mult-cnj-cmod[of k]
  by (auto simp add: field-simps)
ultimately
show ?thesis
  by (auto simp add: field-simps)
qed
ultimately
show ?lhs
  unfolding unitary11-def
  by (auto simp add: mat-adj-def mat-cnj-def complex-cnj field-simps)
qed

```

```

lemma unitary11-inv:
assumes  $k \neq 0 M = k *_{sm} (a, b, \text{cnj } b, \text{cnj } a) \text{ mat-det } (a, b, \text{cnj } b, \text{cnj } a) \neq 0$ 
shows  $\exists k' a' b'. k' \neq 0 \wedge \text{mat-inv } M = k' *_{sm} (a', b', \text{cnj } b', \text{cnj } a') \wedge \text{mat-det } (a', b', \text{cnj } b', \text{cnj } a') \neq 0$ 
using assms
by (subst assms, subst mat-inv-mult-sm[OF assms(1)])
  (rule-tac  $x=1/(k * \text{mat-det } (a, b, \text{cnj } b, \text{cnj } a))$  in exI, rule-tac  $x=\text{cnj } a$  in exI,
  rule-tac  $x=-b$  in exI, simp add: complex-cnj field-simps)

```

```

lemma unitary11-comp:
assumes  $k1 \neq 0 M1 = k1 *_{sm} (a1, b1, \text{cnj } b1, \text{cnj } a1) \text{ mat-det } (a1, b1, \text{cnj } b1, \text{cnj } a1) \neq 0$ 
assumes  $k2 \neq 0 M2 = k2 *_{sm} (a2, b2, \text{cnj } b2, \text{cnj } a2) \text{ mat-det } (a2, b2, \text{cnj } b2, \text{cnj } a2) \neq 0$ 
shows  $\exists k a b. k \neq 0 \wedge M1 *_{mm} M2 = k *_{sm} (a, b, \text{cnj } b, \text{cnj } a) \wedge \text{mat-det } (a, b, \text{cnj } b, \text{cnj } a) \neq 0$ 
using assms
apply (rule-tac  $x=k1*k2$  in exI)
apply (rule-tac  $x=a1*a2 + b1*cnj b2$  in exI)
apply (rule-tac  $x=a1*b2 + b1*cnj a2$  in exI)
apply (auto simp add: field-simps complex-cnj)
apply algebra
done

```

```

lemma unitary11-gen-mat-inv:
assumes unitary11-gen M mat-det M  $\neq 0$ 

```

```

shows unitary11-gen (mat-inv M)
proof-
  obtain k a b where k ≠ 0 ∧ mat-det (a, b, cnj b, cnj a) ≠ 0 ∧ M = k *sm (a,
  b, cnj b, cnj a)
    using assms unitary11-gen-iff[of M]
    by auto
  then obtain k' a' b' where k' ≠ 0 ∧ mat-inv M = k' *sm (a', b', cnj b', cnj
  a') ∧ mat-det (a', b', cnj b', cnj a') ≠ 0
    using unitary11-inv [of k M a b]
    by auto
  thus ?thesis
    using unitary11-gen-iff[of mat-inv M]
    by auto
qed

```

```

lemma unitary11-gen-comp:
  assumes unitary11-gen M1 mat-det M1 ≠ 0 unitary11-gen M2 mat-det M2 ≠ 0
  shows unitary11-gen (M1 *mm M2)
proof-
  from assms obtain k1 k2 a1 a2 b1 b2 where
    k1 ≠ 0 ∧ mat-det (a1, b1, cnj b1, cnj a1) ≠ 0 ∧ M1 = k1 *sm (a1, b1, cnj
    b1, cnj a1)
    k2 ≠ 0 ∧ mat-det (a2, b2, cnj b2, cnj a2) ≠ 0 ∧ M2 = k2 *sm (a2, b2, cnj
    b2, cnj a2)
    using unitary11-gen-iff[of M1] unitary11-gen-iff[of M2]
    by blast
  then obtain k a b where k ≠ 0 ∧ M1 *mm M2 = k *sm (a, b, cnj b, cnj a)
  ∧ mat-det (a, b, cnj b, cnj a) ≠ 0
    using unitary11-comp[of k1 M1 a1 b1 k2 M2 a2 b2]
    by blast
  thus ?thesis
    using unitary11-gen-iff[of M1 *mm M2]
    by blast
qed

```

```

lemma unitary11-sgn-det-orientation:
  assumes k ≠ 0 mat-det (a, b, cnj b, cnj a) ≠ 0 M = k *sm (a, b, cnj b, cnj a)
  shows ∃ k'. sgn k' = sgn (Re (mat-det (a, b, cnj b, cnj a))) ∧ congruence M
  (1, 0, 0, -1) = cor k' *sm (1, 0, 0, -1)
proof-
  let ?x = cnj k * cnj a * (k * a) - (cnj k * b * (k * cnj b))
  have *: ?x = k * cnj k * (a * cnj a - b * cnj b)
    by (auto simp add: field-simps)
  hence is-real ?x
    by auto
  hence cor (Re ?x) = ?x
    by (rule complex-of-real-Re)

```

```

moreover
have  $\text{sgn}(\text{Re } ?x) = \text{sgn}(\text{Re}(a * \text{cnj } a - b * \text{cnj } b))$ 
proof-
  have  $*: \text{Re } ?x = (\text{cmod } k)^2 * \text{Re}(a * \text{cnj } a - b * \text{cnj } b)$ 
  by (subst *, subst complex-mult-cnj-cmod, subst Re-mult-real) (metis Im-complex-of-real,
metis Re-complex-of-real)
  show ?thesis
    using ⟨ $k \neq 0$ ⟩
    by (subst *) (simp add: sgn-mult)
  qed
ultimately
show ?thesis
  using assms(3)
  by (rule-tac x=Re ?x in exI) (auto simp add: mat-adj-def mat-cnj-def complex-cnj)
qed

lemma unitary11-sgn-det:
assumes  $k \neq 0 \text{ mat-det } (a, b, \text{cnj } b, \text{cnj } a) \neq 0 M = k *_{sm} (a, b, \text{cnj } b, \text{cnj } a)$ 
 $M = (A, B, C, D)$ 
shows  $\text{sgn}(\text{Re}(\text{mat-det}(a, b, \text{cnj } b, \text{cnj } a))) = (\text{if } b = 0 \text{ then } 1 \text{ else } \text{sgn}(\text{Re}((A*D)/(B*C)) - 1))$ 
proof (cases b = 0)
  case True
  thus ?thesis
    using assms
    by (simp only: mat-det.simps, subst complex-mult-cnj-cmod, subst complex-Re-diff,
subst Re-complex-of-real, simp)
  next
    case False
    from assms have  $*: A = k * a B = k * b C = k * \text{cnj } b D = k * \text{cnj } a$ 
      by auto
    hence  $*: (A*D)/(B*C) = (a * \text{cnj } a) / (b * \text{cnj } b)$ 
    using ⟨ $k \neq 0$ ⟩
    by simp
    show ?thesis
    using ⟨ $b \neq 0$ ⟩
    apply (subst *, subst Re-divide-real, simp, simp)
    apply (simp only: mat-det.simps)
    apply (subst complex-mult-cnj-cmod)+
    apply ((subst Re-complex-of-real)+, subst complex-Re-diff, (subst Re-complex-of-real)+, simp add: field-simps sgn-if)
    done
qed

lemma unitary11-orientation:
assumes unitary11-gen M M = (A, B, C, D)
shows  $\exists k'. \text{sgn } k' = \text{sgn}(\text{if } B = 0 \text{ then } 1 \text{ else } \text{sgn}(\text{Re}((A*D)/(B*C)) - 1))$ 
 $\wedge \text{congruence } M (1, 0, 0, -1) = \text{cor } k' *_{sm} (1, 0, 0, -1)$ 
proof-

```

```

from ⟨unitary11-gen M⟩
obtain k a b where *:  $k \neq 0$  mat-det (a, b, cnj b, cnj a) ≠ 0 M = k *sm (a, b, cnj b, cnj a)
    using unitary11-gen-iff[of M]
    by auto
moreover
have b = 0  $\longleftrightarrow$  B = 0
    using ⟨M = (A, B, C, D)⟩ *
    by auto
ultimately
show ?thesis
    using unitary11-sgn-det-orientation[OF *] unitary11-sgn-det[OF *] ⟨M = (A, B, C, D)⟩]
    by auto
qed

lemma unitary11-sgn-det-orientation':
assumes congruence M (1, 0, 0, -1) = cor k' *sm (1, 0, 0, -1) k' ≠ 0
shows  $\exists a b k. k \neq 0 \wedge M = k *_{sm} (a, b, cnj b, cnj a) \wedge sgn k' = sgn (Re (mat-det (a, b, cnj b, cnj a)))$ 
proof-
obtain a b k where
    k ≠ 0 mat-det (a, b, cnj b, cnj a) ≠ 0 M = k *sm (a, b, cnj b, cnj a)
    using assms
    using unitary11-gen-iff[of M]
    unfolding unitary11-gen-def
    by auto
moreover
have sgn k' = sgn (Re (mat-det (a, b, cnj b, cnj a)))
proof-
    let ?x = cnj k * cnj a * (k * a) - (cnj k * b * (k * cnj b))
    have *: ?x = k * cnj k * (a * cnj a - b * cnj b)
        by (auto simp add: field-simps)
    hence is-real ?x
        by auto
    hence cor (Re ?x) = ?x
        by (rule complex-of-real-Re)

    have **: sgn (Re ?x) = sgn (Re (a * cnj a - b * cnj b))
proof-
    have *: Re ?x = (cmod k)2 * Re (a * cnj a - b * cnj b)
        by (subst *, subst complex-mult-cnj-cmod, subst Re-mult-real) (metis
        Im-complex-of-real, metis Re-complex-of-real)
    show ?thesis
        using ⟨k ≠ 0⟩
        by (subst *) (simp add: sgn-mult)
qed
moreover
have ?x = cor k'

```

```

using ⟨M = k *sm (a, b, cnj b, cnj a)⟩ assms
by (simp add: mat-adj-def mat-cnj-def complex-cnj complex-diff-def)
hence sgn (Re ?x) = sgn k'
  using ⟨cor (Re ?x) = ?x⟩
  unfolding complex-of-real-def
  by simp
ultimately
show ?thesis
  by simp
qed
ultimately
show ?thesis
  by (rule-tac x=a in exI, rule-tac x=b in exI, rule-tac x=k in exI) simp
qed

end

```

6 Hermitean matrices

```

theory HermiteanMatrices
imports UnitaryMatrices
begin

Hermitean matrices

definition hermitean :: complex-mat ⇒ bool where
hermitean A ⟷ mat-adj A = A

lemma hermitean A ⟷ mat-transpose A = mat-cnj A
unfolding hermitean-def
by (cases A) (auto simp add: mat-adj-def mat-cnj-def)

lemma hermitean-mat-cnj: hermitean H ⟷ hermitean (mat-cnj H)
by (cases H) (auto simp add: hermitean-def mat-adj-def mat-cnj-def)

lemma hermitean-mult-real:
assumes hermitean H
shows hermitean ((cor k) *sm H)
using assms
unfolding hermitean-def
by simp

lemma hermitean-congruence:
assumes hermitean H
shows hermitean (congruence M H)
using assms
unfolding hermitean-def
by (auto simp add: mult-mm-assoc)

lemma hermitean-elems:

```

```

assumes hermitean (A, B, C, D)
shows is-real A is-real D B = cnj C cnj B = C
using assms eq-cnj-iff-real[of A] eq-cnj-iff-real[of D]
by (auto simp add: hermitean-def mat-adj-def mat-cnj-def)

lemma mat-det-hermitean-real:
assumes hermitean A
shows is-real (mat-det A)
using assms
unfolding hermitean-def
by (cases A, auto simp add: mat-adj-def mat-cnj-def) (metis add-0-iff eq-cnj-iff-real
mult-eq-0-iff)

lemma Re-det-sgn-congruence:
assumes hermitean H mat-det M ≠ 0
shows sgn (Re (mat-det (congruence M H))) = sgn (Re (mat-det H))
proof-
have *: mat-det (mat-adj M *mm H *mm M) =
(cmod ((mat-det M)2)) * mat-det H
using complex-mult-cnj-cmod[of mat-det M]
by (auto simp add: mat-det-adj field-simps)
have *: Re (mat-det (mat-adj M *mm H *mm M)) =
(cmod (mat-det M))2 * Re (mat-det H)
by (subst *, subst Re-mult-real, rule is-real-complex-of-real) (subst Re-complex-of-real,
simp)
show ?thesis
using assms
by (subst *) (auto simp add: sgn-mult)
qed

lemma det-sgn-congruence:
assumes hermitean H mat-det M ≠ 0
shows sgn (mat-det (congruence M H)) = sgn (mat-det H)
proof-
have *: mat-det (mat-adj M *mm H *mm M) =
(cmod ((mat-det M)2)) * mat-det H
using complex-mult-cnj-cmod[of mat-det M]
by (auto simp add: mat-det-adj field-simps)
thus ?thesis
using assms
by (subst *, auto simp add: sgn-mult power2-eq-square) (smt mult-eq-0-iff
norm-divide norm-mult norm-sgn of-real-1 of-real-divide of-real-mult sgn-eq times-divide-times-eq)
qed

lemma bilinear-form-hermitean-commute:
assumes hermitean H
shows bilinear-form v1 v2 H = cnj (bilinear-form v2 v1 H)
proof-
have v2 *vm mat-cnj H *vv vec-cnj v1 = vec-cnj v1 *vv (mat-adj H *mv v2)

```

```

by (subst mult-vv-commute, subst mult-mv-mult-vm, simp add: mat-adj-def
mat-transpose-mat-cnj)
also
have ... = bilinear-form v1 v2 H
  using assms
  by (simp add: mult-vv-mv hermitean-def)
finally
show ?thesis
  by (simp add: cnj-mult-vv vec-cnj-mult-vm)
qed

lemma quad-form-hermitean-real:
  assumes hermitean H
  shows is-real (quad-form z H)
  using assms
  by (subst eq-cnj-iff-real[symmetric]) (simp del: quad-form-def add: hermitean-def)

```

Eigenvalues, eigenvectors and diagonalization of Hermitean matrices

```

lemma hermitean-eigenval-real:
  assumes hermitean H eigenval k H
  shows is-real k
proof-
  from assms obtain v where v ≠ vec-zero H *mv v = k *sv v
    unfolding eigenval-def
    by blast
  have k * (v *vv vec-cnj v) = (k *sv v) *vv (vec-cnj v)
    by (simp add: mult-vv-scale-sv1)
  also have ... = (H *mv v) *vv (vec-cnj v)
    using ⟨H *mv v = k *sv v⟩
    by simp
  also have ... = v *vv (mat-transpose H *mv (vec-cnj v))
    by (simp add: mult-mv-vv)
  also have ... = v *vv (vec-cnj (mat-cnj (mat-transpose H) *mv v))
    by (simp add: vec-cnj-mult-mv)
  also have ... = v *vv (vec-cnj (H *mv v))
    using ⟨hermitean H⟩
    by (simp add: hermitean-def mat-adj-def)
  also have ... = v *vv (vec-cnj (k *sv v))
    using ⟨H *mv v = k *sv v⟩
    by simp
  finally have k * (v *vv vec-cnj v) = cnj k * (v *vv vec-cnj v)
    by (simp add: mult-vv-scale-sv2)
  hence k = cnj k
    using ⟨v ≠ vec-zero⟩
    using scalsquare-vv-zero[of v]
    by (simp add: mult-vv-commute)
  thus ?thesis
    by (metis eq-cnj-iff-real)
qed

```

```

lemma hermitean-distinct-eigenvals:
  assumes hermitean H
  shows ( $\exists k_1 k_2. k_1 \neq k_2 \wedge \text{eigenval } k_1 H \wedge \text{eigenval } k_2 H$ )  $\vee$  mat-diagonal H
  proof-
    obtain A B C D where HH:  $H = (A, B, C, D)$ 
      by (cases H) auto
    show ?thesis
    proof (cases B = 0)
      case True
      thus ?thesis
        using ⟨hermitean H⟩ hermitean-elems[of A B C D] HH
        by auto
    next
      case False
      have (mat-trace H)2  $\neq 4 * \text{mat-det } H$ 
      proof (rule ccontr)
        have C = cnj B is-real A is-real D
          using hermitean-elems HH ⟨hermitean H⟩
          by auto
        assume  $\neg$  ?thesis
        hence  $(A + D)^2 = 4*(A*D - B*C)$ 
          using HH
          by auto
        hence  $(A - D)^2 = -4*B*cnj B$ 
          using ⟨C = cnj B⟩
          by (auto simp add: power2-eq-square field-simps) algebra
        hence  $(A - D)^2 / \text{cor}((\text{cmod } B)^2) = -4$ 
          using ⟨B ≠ 0⟩ complex-mult-cnj-cmod[of B]
          by (auto simp add: field-simps)
        hence  $(\text{Re } A - \text{Re } D)^2 / (\text{cmod } B)^2 = -4$ 
          using ⟨is-real A⟩ ⟨is-real D⟩ ⟨B ≠ 0⟩
          using Re-divide-real[of cor ((cmod B)2) (A - D)2]
          by (auto simp add: power2-eq-square)
        thus False
        by (metis abs-neg-numeral abs-power2 neg-numeral-neq-numeral power-divide)
    qed
    show ?thesis
      apply (rule disjI1)
      apply (subst eigen-equation)+
      using complex-quadratic-two-solutions[of -mat-trace H mat-det H] ⟨(mat-trace H)2  $\neq 4 * \text{mat-det } H$ ⟩
      apply auto
      apply (rule-tac x=k1 in exI, rule-tac x=k2 in exI)
      apply (simp add: complex-diff-def)
      done
    qed
  qed

```

```

lemma hermitean-ortho-eigenvecs:
  assumes hermitean H
  assumes eigenpair k1 v1 H eigenpair k2 v2 H k1 ≠ k2
  shows vec-cnj v2 *vv v1 = 0 vec-cnj v1 *vv v2 = 0
proof-
  from assms
  have v1 ≠ vec-zero H *mv v1 = k1 *sv v1
    using eigenpair-def
    by auto
  have real-k: is-real k1 is-real k2
    using assms
    using hermitean-eigenval-real[of H k1]
    using hermitean-eigenval-real[of H k2]
    unfoldng eigenpair-def eigenval-def
    by blast+
  have vec-cnj (H *mv v2) = vec-cnj (k2 *sv v2)
    using ⟨H *mv v2 = k2 *sv v2⟩
    by auto
  hence vec-cnj v2 *vm H = k2 *sv vec-cnj v2
    using ⟨hermitean H⟩ real-k eq-cnj-iff-real[of k1] eq-cnj-iff-real[of k2]
    unfoldng hermitean-def
    by (cases H, cases v2) (auto simp add: mat-adj-def mat-cnj-def vec-cnj-def
complex-cnj)
  have k2 * (vec-cnj v2 *vv v1) = k1 * (vec-cnj v2 *vv v1)
    using ⟨H *mv v1 = k1 *sv v1⟩
    using ⟨vec-cnj v2 *vm H = k2 *sv vec-cnj v2⟩
    by (cases v1, cases v2, cases H) (auto simp add: vec-cnj-def field-simps, algebra)
  thus vec-cnj v2 *vv v1 = 0
    using ⟨k1 ≠ k2⟩
    by simp
  hence cnj (vec-cnj v2 *vv v1) = 0
    by simp
  thus vec-cnj v1 *vv v2 = 0
    by (simp add: cnj-mult-vv mult-vv-commute)
qed

```

```

lemma hermitean-diagonizable:
  assumes hermitean H
  shows ∃ k1 k2 M. mat-det M ≠ 0 ∧ unitary M ∧ congruence M H = (k1, 0,
0, k2) ∧
  is-real k1 ∧ is-real k2 ∧ sgn (Re k1 * Re k2) = sgn (Re (mat-det
H))
proof-
  from assms
  have (∃ k1 k2. k1 ≠ k2 ∧ eigenval k1 H ∧ eigenval k2 H) ∨ mat-diagonal H
    using hermitean-distinct-eigenvals[of H]
    by simp

```

```

thus ?thesis
proof
  assume  $\exists k_1 k_2. k_1 \neq k_2 \wedge \text{eigenval } k_1 H \wedge \text{eigenval } k_2 H$ 
  then obtain  $k_1 k_2$  where  $k_1 \neq k_2 \text{ eigenval } k_1 H \text{ eigenval } k_2 H$ 
    using hermitean-distinct-eigenvals
    by blast
  then obtain  $v_1 v_2$  where  $\text{eigenpair } k_1 v_1 H \text{ eigenpair } k_2 v_2 H$ 
     $v_1 \neq \text{vec-zero} v_2 \neq \text{vec-zero}$ 
    unfolding eigenval-def eigenpair-def
    by blast
  hence  $*: \text{vec-cnj } v_2 *_{vv} v_1 = 0 \text{ vec-cnj } v_1 *_{vv} v_2 = 0$ 
    using  $\langle k_1 \neq k_2 \rangle \text{ hermitean-ortho-eigenvecs } \langle \text{hermitean } H \rangle$ 
    by auto
  obtain  $v_{11} v_{12} v_{21} v_{22}$  where  $vv: v_1 = (v_{11}, v_{12}) v_2 = (v_{21}, v_{22})$ 
    by (cases v1, cases v2) auto
  let  $?nv1' = \text{vec-cnj } v_1 *_{vv} v_1$  and  $?nv2' = \text{vec-cnj } v_2 *_{vv} v_2$ 
  let  $?nv1 = \text{cor } (\text{sqrt } (\text{Re } ?nv1'))$ 
  let  $?nv2 = \text{cor } (\text{sqrt } (\text{Re } ?nv2'))$ 
  have  $?nv1' \neq 0 ?nv2' \neq 0$ 
    using  $\langle v_1 \neq \text{vec-zero} \rangle \langle v_2 \neq \text{vec-zero} \rangle vv$ 
    by (simp add: scalsquare-vv-zero)+
  moreover
  have is-real ?nv1' is-real ?nv2'
    using vv
    by (auto simp add: vec-cnj-def)
  ultimately
  have  $?nv1 \neq 0 ?nv2 \neq 0$ 
    by - (cases ?nv1', cases ?nv2', auto)+
  have  $\text{Re } (?nv1') \geq 0 \text{ Re } (?nv2') \geq 0$ 
    using vv
    by (auto simp add: vec-cnj-def)
  obtain  $nv1 nv2$  where  $nv1 = ?nv1 nv1 \neq 0 nv2 = ?nv2 nv2 \neq 0$ 
    using  $\langle ?nv1 \neq 0 \rangle \langle ?nv2 \neq 0 \rangle$ 
    by auto
  let  $?M = (1/nv1 * v_{11}, 1/nv2 * v_{21}, 1/nv1 * v_{12}, 1/nv2 * v_{22})$ 

  have is-real  $k_1$  is-real  $k_2$ 
    using  $\langle \text{eigenval } k_1 H \rangle \langle \text{eigenval } k_2 H \rangle \langle \text{hermitean } H \rangle$ 
    by (auto simp add: hermitean-eigenval-real)
  moreover
  have mat-det ?M  $\neq 0$ 
  proof (rule ccontr)
    assume  $\neg ?thesis$ 
    hence  $v_{11} * v_{22} = v_{12} * v_{21}$ 
      using  $\langle nv1 \neq 0 \rangle \langle nv2 \neq 0 \rangle$ 
      by (auto simp add: field-simps)
    hence  $\exists k. k \neq 0 \wedge v_2 = k *_{sv} v_1$ 
      using vv  $\langle v_1 \neq \text{vec-zero} \rangle \langle v_2 \neq \text{vec-zero} \rangle$ 
      apply auto

```

```

apply (rule-tac x=v21/v11 in exI, force simp add: field-simps)
apply (rule-tac x=v21/v11 in exI, force simp add: field-simps)
apply (rule-tac x=v22/v12 in exI, force simp add: field-simps)
apply (rule-tac x=v22/v12 in exI, force simp add: field-simps)
done
thus False
  using <vec-cnj v1 *vv v2 = 0> vv <?nv1' ≠ 0>
  by (auto simp add: vec-cnj-def field-simps) (metis comm-semiring-1-class.normalize-semiring-rules(34)
mult-eq-0-iff)
qed
moreover
have unitary ?M
proof-
  have **: cnj nv1 * nv1 = ?nv1' cnj nv2 * nv2 = ?nv2'
    using <nv1 = ?nv1> <nv1 ≠ 0> <nv2 = ?nv2> <nv2 ≠ 0> <is-real ?nv1'>
<is-real ?nv2'>
    using <Re (?nv1') ≥ 0> <Re (?nv2') ≥ 0>
    by (auto simp add: complex-of-real-Re)
  have ***: cnj nv1 * nv2 ≠ 0 cnj nv2 * nv1 ≠ 0
    using vv <nv1 = ?nv1> <nv1 ≠ 0> <nv2 = ?nv2> <nv2 ≠ 0> <is-real ?nv1'>
<is-real ?nv2'>
    by auto
show ?thesis
  unfolding unitary-def
  using vv ** <?nv1' ≠ 0> <?nv2' ≠ 0> * ***
  apply (auto simp add: mat-adj-def mat-cnj-def vec-cnj-def complex-cnj)
  apply (metis add-divide-distrib divide-self-if)
  apply (metis add-divide-distrib divide-zero-left)
  apply (metis add-divide-distrib divide-zero-left)
  apply (metis add-divide-distrib divide-self-if)
done
qed
moreover
have congruence ?M H = (k1, 0, 0, k2)
proof-
  have mat-inv ?M *mm H *mm ?M = (k1, 0, 0, k2)
  proof-
    have *: H *mm ?M = ?M *mm (k1, 0, 0, k2)
      using <eigenpair k1 v1 H> <eigenpair k2 v2 H> vv <?nv1 ≠ 0> <?nv2 ≠ 0>
      unfolding eigenpair-def
      apply (cases H)
      apply (auto simp add: vec-cnj-def)
      apply (metis add-divide-distrib mult.commute)+
    done
  show ?thesis
    using mult-mm-inv-l[of ?M (k1, 0, 0, k2) H *mm ?M, OF <mat-det ?M
≠ 0> *[symmetric], symmetric]
    by (simp add: mult-mm-assoc)

```

```

qed
moreover
have mat-inv ?M = mat-adj ?M
  using ⟨mat-det ?M ≠ 0⟩ ⟨unitary ?M⟩ mult-mm-inv-r[of ?M mat-adj ?M
eye]
  by (simp add: unitary-def)
ultimately
show ?thesis
  by simp
qed
moreover
have sgn (Re k1 * Re k2) = sgn (Re (mat-det H))
  using ⟨congruence ?M H = (k1, 0, 0, k2)⟩ ⟨is-real k1⟩ ⟨is-real k2⟩
  using Re-det-sgn-congruence[of H ?M] ⟨mat-det ?M ≠ 0⟩ ⟨hermitean H⟩
  by simp
ultimately
show ?thesis
  by (rule-tac x=k1 in exI, rule-tac x=k2 in exI, rule-tac x=?M in exI) simp
next
assume mat-diagonal H
then obtain A D where H = (A, 0, 0, D)
  by (cases H) auto
moreover
hence is-real A is-real D
  using ⟨hermitean H⟩ hermitean-elems[of A 0 0 D]
  by auto
ultimately
show ?thesis
  by (rule-tac x=A in exI, rule-tac x=D in exI, rule-tac x=eye in exI) (simp
add: unitary-def mat-adj-def mat-cnj-def)
qed
qed

end

```

7 Elementary complex geometry

```

theory ElementaryComplexGeometry
imports MoreComplex LinearSystems
begin

```

```

definition colinear :: complex ⇒ complex ⇒ complex ⇒ bool where
  colinear z1 z2 z3 ⟷ z1 = z2 ∨ Im ((z3 - z1)/(z2 - z1)) = 0

lemma colinear-ex-real:
  colinear z1 z2 z3 ⟷ (∃ k::real. z1 = z2 ∨ z3 - z1 = complex-of-real k * (z2
  - z1))

```

```

unfolding colinear-def
by (auto split: split-if-asm) (metis Im.simps complex.exhaust complex-of-real-def
eq-iff-diff-eq-0 nonzero-divide-eq-eq)

lemma colinear-sym1:
  colinear z1 z2 z3  $\longleftrightarrow$  colinear z1 z3 z2
unfolding colinear-def
using div-reals[of 1  $(z3 - z1)/(z2 - z1)$ ] div-reals[of 1  $(z2 - z1)/(z3 - z1)$ ]
by auto

lemma colinear-sym2':
  assumes colinear z1 z2 z3
  shows colinear z2 z1 z3
proof-
  obtain k where z1 = z2  $\vee$  z3 - z1 = complex-of-real k * (z2 - z1)
  using assms
  unfolding colinear-ex-real
  by auto
  thus ?thesis
  proof
    assume z3 - z1 = complex-of-real k * (z2 - z1)
    thus ?thesis
      unfolding colinear-ex-real
      by (rule-tac x=1-k in exI) (auto simp add: field-simps)
    qed (simp add: colinear-def)
  qed

lemma colinear-sym2:
  colinear z1 z2 z3  $\longleftrightarrow$  colinear z2 z1 z3
  using colinear-sym2'[of z1 z2 z3] colinear-sym2'[of z2 z1 z3]
  by auto

lemma colinear-trans1:
  assumes colinear z0 z2 z1 colinear z0 z3 z1 z0  $\neq$  z1
  shows colinear z0 z2 z3
  using assms
  unfolding colinear-ex-real
  by (cases z0 = z2, auto) (rule-tac x=k/ka in exI, case-tac ka = 0, auto simp
add: field-simps)

lemma colinear-det:
  assumes  $\neg$  colinear z2 z3 z1
  shows det2 (z1 - z2) (cnj (z1 - z2)) (z2 - z3) (cnj (z2 - z3))  $\neq$  0
proof-
  from assms have  $((z1 - z2) / (z3 - z2)) - cnj ((z1 - z2) / (z3 - z2)) \neq 0$ 
  z3  $\neq$  z2
  unfolding colinear-def
  using im-complex[of  $(z1 - z2) / (z3 - z2)$ ]
  by auto

```

```

thus ?thesis
  by (auto simp add: field-simps complex-cnj-divide complex-cnj-add complex-cnj-diff)
qed

```

```

definition line :: complex  $\Rightarrow$  complex  $\Rightarrow$  complex set where
  line z1 z2 = {z. colinear z1 z2 z}

```

```

lemma line-points-colinear:
  assumes z1  $\in$  line z z' z2  $\in$  line z z' z3  $\in$  line z z' z  $\neq$  z'
  shows colinear z1 z2 z3
  using assms
  unfolding line-def
  by auto (smt colinear-sym1 colinear-sym2 colinear-trans1)

```

```

lemma line-param:
  shows z1 + complex-of-real k * (z2 - z1)  $\in$  line z1 z2
  unfolding line-def
  by (auto simp add: colinear-def)

```

```

definition circle :: complex  $\Rightarrow$  real  $\Rightarrow$  complex set where
  circle  $\mu$  r = {z. cmod (z -  $\mu$ ) = r}

```

```

lemma line-equation:
  assumes z1  $\neq$  z2  $\mu$  = rot90 (z2 - z1)
  shows line z1 z2 = {z. cnj  $\mu$ *z +  $\mu$ *cnj z - (cnj  $\mu$  * z1 +  $\mu$  * cnj z1) = 0}
  proof-
  {
    fix z
    have z  $\in$  line z1 z2  $\longleftrightarrow$  Im ((z - z1)/(z2 - z1)) = 0
      using assms
      by (simp add: line-def colinear-def)
    also have ...  $\longleftrightarrow$  (z - z1)/(z2 - z1) = cnj ((z - z1)/(z2 - z1))
      using complex-diff-cnj[of (z - z1)/(z2 - z1)]
      by auto
    also have ...  $\longleftrightarrow$  (z - z1)*(cnj z2 - cnj z1) = (cnj z - cnj z1)*(z2 - z1)
      using assms(1)
      by auto
    by auto (metis (lifting) complex-cnj-cancel-iff complex-cnj-diff complex-cnj-divide
      frac-eq-eq right-minus-eq)+
    also have ...  $\longleftrightarrow$  cnj(z2 - z1)*z - (z2 - z1)*cnj z - (cnj(z2 - z1)*z1 -
      (z2 - z1)*cnj z1) = 0
      by (simp add: complex-cnj-diff field-simps)
    also have ...  $\longleftrightarrow$  cnj  $\mu$  * z +  $\mu$  * cnj z - (cnj  $\mu$  * z1 +  $\mu$  * cnj z1) = 0
      using assms cnj-mix-minus
      by simp
    finally have z  $\in$  line z1 z2  $\longleftrightarrow$  cnj  $\mu$  * z +  $\mu$  * cnj z - (cnj  $\mu$  * z1 +  $\mu$  *

```

```

 $cnj z1) = 0$ 
.
}

thus ?thesis
  by auto
qed

lemma circle-equation:
assumes  $r \geq 0$ 
shows  $\text{circle } \mu r = \{z. z * cnj z - z * cnj \mu - cnj z * \mu + \mu * cnj \mu - \text{complex-of-real}(r * r) = 0\}$ 
proof (safe)
fix  $z$ 

assume  $z \in \text{circle } \mu r$ 
hence  $(z - \mu) * cnj(z - \mu) = \text{complex-of-real}(r * r)$ 
  unfolding circle-def
  using complex-mult-cnj-cmod[of  $z - \mu$ ]
  by (auto simp add: power2-eq-square)
thus  $z * cnj z - z * cnj \mu - cnj z * \mu + \mu * cnj \mu - \text{complex-of-real}(r * r) = 0$ 
  by (auto simp add: field-simps complex-cnj-diff)
next
fix  $z$ 
assume  $z * cnj z - z * cnj \mu - cnj z * \mu + \mu * cnj \mu - \text{complex-of-real}(r * r) = 0$ 
hence  $(z - \mu) * cnj(z - \mu) = \text{complex-of-real}(r * r)$ 
  by (auto simp add: field-simps complex-cnj-diff)
thus  $z \in \text{circle } \mu r$ 
  using assms
  using complex-mult-cnj-cmod[of  $z - \mu$ ]
  using power2-eq-imp-eq[of cmod( $z - \mu$ )  $r$ ]
  unfolding circle-def power2-eq-square[symmetric] complex-of-real-def
  by auto
qed

```

```

definition circline where
circline A BC D = {z. cor A * z * cnj z + cnj BC * z + BC * cnj z + cor D = 0}

lemma circline-circle:
assumes  $A \neq 0$   $A * D \leq (\text{cmod } BC)^2$ 
cl = circline A BC D
 $\mu = -BC / \text{complex-of-real } A$   $r^2 = ((\text{cmod } BC)^2 - A * D) / A^2$   $r = \sqrt{r^2}$ 
shows cl = circle  $\mu r$ 
proof-
have *: cl = {z. z * cnj z + cnj(BC / \text{complex-of-real } A) * z + (BC / \text{complex-of-real } A) * cnj z + \text{complex-of-real}(D / A) = 0}

```

```

using ⟨cl = circline A BC D⟩ ⟨A ≠ 0⟩
by (auto simp add: circline-def complex-cnj-divide field-simps)

have r2 ≥ 0
proof-
  have (cmod BC)2 − A * D ≥ 0
    using ⟨A * D ≤ (cmod BC)2⟩
    by auto
  thus ?thesis
    using ⟨A ≠ 0⟩ ⟨r2 = ((cmod BC)2 − A*D) / A2⟩
    by (metis zero-le-divide-iff zero-le-power2)
qed
hence **: r * r = r2 r ≥ 0
using ⟨r = sqrt r2⟩
by (auto simp add: real-sqrt-mult[symmetric])

have ***: − μ * − cnj μ − complex-of-real r2 = complex-of-real (D / A)
  using ⟨μ = − BC / complex-of-real A⟩ ⟨r2 = ((cmod BC)2 − A*D) / A2⟩
  by (auto simp add: complex-cnj-divide complex-cnj-minus complex-mult-cnj-cmod
power2-eq-square complex-of-real-def complex-divide-def div-reals field-simps intro!
complex-eqI)
  thus ?thesis
    using ⟨r2 = ((cmod BC)2 − A*D) / A2⟩ ⟨μ = − BC / complex-of-real A⟩
    by (subst *, subst circle-equation[of r μ, OF ⟨r ≥ 0⟩], subst **) (auto simp
add: complex-cnj-minus complex-cnj-divide field-simps power2-eq-square)
qed

lemma circline-ex-circle:
assumes A ≠ 0 A * D ≤ (cmod BC)2
cl = circline A BC D
shows ∃ μ r. cl = circle μ r
using circline-circle[OF assms]
by auto

lemma circle-circline:
assumes cl = circle μ r r ≥ 0
shows cl = circline 1 (−μ) ((cmod μ)2 − r2)
proof-
  have complex-of-real ((cmod μ)2 − r2) = μ * cnj μ − complex-of-real (r2)
    by (auto simp add: complex-mult-cnj-cmod)
  thus cl = circline 1 (− μ) ((cmod μ)2 − r2)
    using assms
    using circle-equation[of r μ]
    unfolding circline-def power2-eq-square
    by (simp add: complex-cnj-minus field-simps)
qed

lemma circle-ex-circline:
assumes cl = circle μ r r ≥ 0

```

```

shows  $\exists A BC D. A \neq 0 \wedge A*D \leq (\text{cmod } BC)^2 \wedge cl = \text{circline } A BC D$ 
using circle-circline[OF assms]
using  $\langle r \geq 0 \rangle$ 
by (rule-tac  $x=1$  in exI, rule-tac  $x=-\mu$  in exI, rule-tac  $x=\text{Re } (\mu * \text{cnj } \mu) - (r * r)$  in exI) (simp add: complex-mult-cnj-cmod power2-eq-square)

```

lemma circline-line:

assumes

$$A = 0 \quad BC \neq 0$$

$$cl = \text{circline } A BC D$$

$$z1 = -\text{cor } D * BC / (2 * BC * \text{cnj } BC)$$

$$z2 = z1 + ii * \text{sgn} (\text{if arg } BC > 0 \text{ then } -BC \text{ else } BC)$$

shows

$$cl = \text{line } z1 z2$$

proof-

$$\text{have } cl = \{z. \text{cnj } BC*z + BC*\text{cnj } z + \text{complex-of-real } D = 0\}$$

using *assms*

by (simp add: circline-def)

$$\text{have } \{z. \text{cnj } BC*z + BC*\text{cnj } z + \text{complex-of-real } D = 0\} =$$

$$\{z. \text{cnj } BC*z + BC*\text{cnj } z - (\text{cnj } BC*z1 + BC*\text{cnj } z1) = 0\}$$

using $\langle BC \neq 0 \rangle$ *assms*

by (auto simp add: complex-cnj-minus complex-cnj-divide complex-cnj-mult)

moreover

$$\text{have } z1 \neq z2$$

using $\langle BC \neq 0 \rangle$ *assms*

by (auto simp add: sgn-eq)

moreover

$$\text{have } \exists k. k \neq 0 \wedge BC = \text{cor } k * \text{rot90} (z2 - z1)$$

using *assms*

apply auto

apply (rule-tac $x=(\text{cmod } BC)$ in exI, simp, metis Complex.Re-sgn Im-sgn cmod-cis mult.commute complex-surj eq-divide-eq mult-zero-left sgn-eq)

apply (rule-tac $x=-(\text{cmod } BC)$ in exI, simp, metis Complex.Re-sgn Im-sgn cis-arg cmod-cis mult.commute complex-minus-def minus-minus minus-mult-left)

done

$$\text{then obtain } k \text{ where } \text{cor } k \neq 0 \quad BC = \text{cor } k * \text{rot90} (z2 - z1)$$

by auto

moreover

$$\text{have } *: \bigwedge z. \text{cnj-mix } (BC / \text{cor } k) z - \text{cnj-mix } (BC / \text{cor } k) z1 = (1 / \text{cor } k)$$

$$* (\text{cnj-mix } BC z - \text{cnj-mix } BC z1)$$

using $\langle \text{cor } k \neq 0 \rangle$

by (simp add: complex-cnj field-simps)

$$\text{hence } \{z. \text{cnj-mix } BC z - \text{cnj-mix } BC z1 = 0\} = \{z. \text{cnj-mix } (BC / \text{cor } k) z - \text{cnj-mix } (BC / \text{cor } k) z1 = 0\}$$

using $\langle \text{cor } k \neq 0 \rangle$

by auto

ultimately

$$\text{have } cl = \text{line } z1 z2$$

```

using line-equation[of z1 z2 BC/cor k] ⟨cl = {z. cnj BC*z + BC*cnj z +
complex-of-real D = 0}⟩
by auto
thus ?thesis
using ⟨z1 ≠ z2⟩
by blast
qed

```

```

lemma circline-ex-line:
assumes
  A = 0 BC ≠ 0
  cl = circline A BC D
shows ∃ z1 z2. z1 ≠ z2 ∧ cl = line z1 z2
proof-
  let ?z1 = - cor D * BC / (2 * BC * cnj BC)
  let ?z2 = ?z1 + i * sgn (if 0 < arg BC then - BC else BC)
  have ?z1 ≠ ?z2
    using ⟨BC ≠ 0⟩
    by (simp add: sgn-eq)
  thus ?thesis
    using circline-line[OF assms, of ?z1 ?z2] ⟨BC ≠ 0⟩
    by (rule-tac x=?z1 in exI, rule-tac x=?z2 in exI, simp)
qed

```

```

lemma line-ex-circline:
assumes cl = line z1 z2 z1 ≠ z2
shows ∃ BC D. BC ≠ 0 ∧ cl = circline 0 BC D
proof-
  let ?BC = rot90 (z2 - z1)
  let ?D = Re (- 2 * scalprod z1 ?BC)
  show ?thesis
  proof (rule-tac x=?BC in exI, rule-tac x=?D in exI, rule conjI)
    show ?BC ≠ 0
    using ⟨z1 ≠ z2⟩
    by (metis complex-minus-def eq-iff-diff-eq-0 i-mult-Complex minus-diff-eq
mult-zero-right)
  next
    have *: complex-of-real (Re (- 2 * scalprod z1 (rot90 (z2 - z1)))) = -
(cnj-mix z1 (rot90 (z2 - z1)))
    by (cases z1, cases z2, auto simp add: complex-of-real-def field-simps)
    show cl = circline 0 ?BC ?D
    apply (subst assms, subst line-equation[of z1 z2 ?BC])
    unfolding circline-def
    by (fact, simp, subst *, simp add: field-simps)
  qed
qed
end

```

```

theory Angles
imports MoreComplex
begin

```

```

definition ang-vec () where
  [simp]:  $z1 z2 \equiv |\arg z2 - \arg z1|$ 

```

```

definition ang-vec-c (c) where
  [simp]:  $c z1 z2 \equiv \text{abs}(z1 z2)$ 

```

```

definition acute-ang where
  [simp]:  $\text{acute-ang } \alpha = (\text{if } \alpha > \pi / 2 \text{ then } \pi - \alpha \text{ else } \alpha)$ 

```

```

definition ang-vec-a (a) where
  [simp]:  $a z1 z2 \equiv \text{acute-ang}(c z1 z2)$ 

```

```

lemma ang-vec-sym:
  assumes  $z1 z2 \neq pi$ 
  shows  $z1 z2 = - z2 z1$ 
  using assms
  unfolding ang-vec-def
  using canon-ang-uminus[of arg z2 - arg z1]
  by simp

```

```

lemma ang-vec-sym-pi:
  assumes  $z1 z2 = pi$ 
  shows  $z1 z2 = z2 z1$ 
  using assms
  unfolding ang-vec-def
  using canon-ang-uminus-pi[of arg z2 - arg z1]
  by simp

```

```

lemma ang-vec-c-sym:
  shows  $c z1 z2 = c z2 z1$ 
  unfolding ang-vec-c-def
  using ang-vec-sym-pi[of z1 z2] ang-vec-sym[of z1 z2]
  by (cases z1 z2 = pi) auto

```

```

lemma ang-vec-a-sym:

```

```

a z1 z2 = a z2 z1
unfolding ang-vec-a-def
using ang-vec-c-sym
by auto

```

```

lemma ang-vec-c-bounded: 0 ≤ c z1 z2 ∧ c z1 z2 ≤ pi
using canon-ang(1)[of arg z2 - arg z1] canon-ang(2)[of arg z2 - arg z1]
by auto

```

```

lemma ortho-c-scalprod0:
assumes z1 ≠ 0 z2 ≠ 0
shows c z1 z2 = pi/2 ↔ scalprod z1 z2 = 0
proof
assume c z1 z2 = pi / 2
have |arg z2 - arg z1| = arg (z2 / z1)
using arg-div[of z2 z1] assms
by auto
hence arg (z2 / z1) = pi/2 ∨ arg (z2 / z1) = -pi/2
using <c z1 z2 = pi/2>
unfolding ang-vec-c-def
unfolding ang-vec-def
by auto
hence Re (z2 / z1) = 0
using re-complex-zero-arg1[of z2/z1]
by auto
hence z2 / z1 + cnj (z2 / z1) = 0
using re-complex[of z2/z1]
by (auto simp add: complex-of-real-def[symmetric])
thus scalprod z1 z2 = 0
using assms complex-cnj-divide[of z2 z1]
using add-frac-eq[of z1 cnj z1 z2 cnj z2]
using divide-eq-0-iff[of z2 * cnj z1 + cnj z2 * z1 z1 * cnj z1]
by (auto simp add:field-simps)

```

```

next
assume scalprod z1 z2 = 0
hence z2 * cnj z1 + cnj z2 * z1 = 0
by (simp add:field-simps)
hence z2 / z1 + cnj (z2 / z1) = 0
using assms complex-cnj-divide[of z2 z1]
using add-frac-eq[of z1 cnj z1 z2 cnj z2]
using divide-eq-0-iff[of z2 * cnj z1 + cnj z2 * z1 z1 * cnj z1]
by auto
hence Re (z2 / z1) = 0
using re-complex[of z2/z1]
by auto
have z2 / z1 ≠ 0
using assms
by auto

```

```

hence  $\arg(z2 / z1) = pi/2 \vee \arg(z2 / z1) = -pi/2$ 
  using  $\langle Re(z2 / z1) = 0 \rangle$  re-complex-zero-arg2[of z2/z1]
  by auto
have  $\lfloor \arg z2 - \arg z1 \rfloor = \arg(z2 / z1)$ 
  using arg-div[of z2 z1] assms
  by auto
thus  $c z1 z2 = pi / 2$ 
  using  $\langle \arg(z2 / z1) = pi/2 \vee \arg(z2 / z1) = -pi/2 \rangle$ 
  unfolding ang-vec-c-def
  unfolding ang-vec-def
  by (metis abs-minus-cancel abs-of-nonneg minus-divide-left pi-half-ge-zero)
qed

```

```

lemma ortho-a-scalprod0:
  assumes  $z1 \neq 0$   $z2 \neq 0$ 
  shows  $a z1 z2 = pi/2 \longleftrightarrow \text{scalprod } z1 z2 = 0$ 
  unfolding ang-vec-a-def
  using assms ortho-c-scalprod0[of z1 z2]
  by auto

```

```

lemma canon-ang-plus-pi1:
  assumes  $z1 z2 > 0$ 
  shows  $\lfloor z1 z2 + pi \rfloor = z1 z2 - pi$ 
proof (rule canon-ang-eqI)
  show  $\exists x::int. z1 z2 - pi - (z1 z2 + pi) = 2 * \text{real } x * pi$ 
    by (rule-tac x=-1 in exI) auto
next
  show  $-pi < z1 z2 - pi \wedge z1 z2 - pi \leq pi$ 
    using assms
    unfolding ang-vec-def
    using canon-ang(1)[of arg z2 - arg z1] canon-ang(2)[of arg z2 - arg z1]
    by auto
qed

```

```

lemma canon-ang-plus-pi2:
  assumes  $z1 z2 \leq 0$ 
  shows  $\lfloor z1 z2 + pi \rfloor = z1 z2 + pi$ 
proof (rule canon-ang-id)
  show  $-pi < z1 z2 + pi \wedge z1 z2 + pi \leq pi$ 
    using assms
    unfolding ang-vec-def
    using canon-ang(1)[of arg z2 - arg z1] canon-ang(2)[of arg z2 - arg z1]
    by auto
qed

```

```

lemma ang-vec-opposite1:

```

```

assumes z1 ≠ 0
shows (‐z1) z2 = | z1 z2 – pi |
unfolding ang-vec-def
apply (subst arg-uminus[OF assms])
apply (subst canon-ang-arg[of z2, symmetric])
apply (subst canon-ang-diff[of arg z2 arg z1 + pi, symmetric])
apply (subst canon-ang-id[of pi, symmetric]) back
apply simp
apply (subst canon-ang-diff[of arg z2 – arg z1 pi, symmetric])
apply (simp add: field-simps)
done

lemma ang-vec-opposite2:
assumes z2 ≠ 0
shows z1 (‐z2) = | z1 z2 + pi |
unfolding ang-vec-def
using arg-mult[of ‐1 z2] assms
using arg-complex-of-real-negative[of ‐1]
using canon-ang-diff[of arg ‐1 + arg z2 arg z1]
using canon-ang-sum[of arg z2 – arg z1 pi]
using canon-ang-id[of pi] canon-ang-arg[of z1]
by auto (metis (hide-lams, no-types) ab-diff-minus ab-semigroup-add-class.add-ac(1)
minus-add minus-add-distrib minus-minus)

lemma ang-vec-opposite-opposite:
assumes z1 ≠ 0 z2 ≠ 0
shows (‐z1) (‐z2) = z1 z2
apply (subst ang-vec-opposite1[OF assms(1)])
apply (subst ang-vec-opposite2[OF assms(2)])
apply (subst canon-ang-id[of pi, symmetric]) back
apply simp
apply (subst canon-ang-diff[symmetric])
apply (simp del: ang-vec-def)
by (metis ang-vec-def canon-ang(1) canon-ang(2) canon-ang-id)

lemma ang-vec-a-opposite2:
a z1 z2 = a z1 (‐z2)
proof(cases z2 = 0)
case True
thus ?thesis
by (metis minus-zero)
next
case False
thus ?thesis
proof(cases z1 z2 < –pi / 2)
case True
hence z1 z2 < 0
by auto (metis less-trans minus-pi-half-less-zero)
have a z1 z2 = pi + z1 z2

```

```

using True ⊢ z1 z2 < 0
unfolding ang-vec-a-def ang-vec-c-def ang-vec-a-def abs-real-def
by auto
moreover
have a z1 (-z2) = pi + z1 z2
  unfolding ang-vec-a-def ang-vec-c-def abs-real-def
  using canon-ang(1)[of arg z2 - arg z1] canon-ang(2)[of arg z2 - arg z1]
  using canon-ang-plus-pi2[of z1 z2] True ⊢ z1 z2 < 0 ⊢ z2 ≠ 0
  using ang-vec-opposite2[of z2 z1]
  by auto
ultimately
show ?thesis
  by auto
next
case False
show ?thesis
proof (cases z1 z2 ≤ 0)
  case True
    have a z1 z2 = - z1 z2
      using ⊢ z1 z2 < - pi / 2 ⊢ True
      unfolding ang-vec-a-def ang-vec-c-def ang-vec-a-def abs-real-def
      by auto
    moreover
    have a z1 (-z2) = - z1 z2
      using ⊢ z1 z2 < - pi / 2 ⊢ True
      unfolding ang-vec-a-def ang-vec-c-def abs-real-def
      using canon-ang-plus-pi2[of z1 z2]
      using canon-ang(1)[of arg z2 - arg z1] canon-ang(2)[of arg z2 - arg z1]
      using ⊢ z2 ≠ 0 ang-vec-opposite2[of z2 z1]
      by auto
    ultimately
    show ?thesis
      by simp
  next
  case False
  show ?thesis
  proof (cases z1 z2 < pi / 2)
    case True
      have a z1 z2 = z1 z2
        using ⊢ z1 z2 ≤ 0 ⊢ True
        unfolding ang-vec-a-def ang-vec-c-def ang-vec-a-def abs-real-def
        by auto
    moreover
    have a z1 (-z2) = z1 z2
      using ⊢ z1 z2 ≤ 0 ⊢ True
      unfolding ang-vec-a-def ang-vec-c-def abs-real-def
      using canon-ang-plus-pi1[of z1 z2]
      using canon-ang(1)[of arg z2 - arg z1] canon-ang(2)[of arg z2 - arg z1]
      using ⊢ z2 ≠ 0 ang-vec-opposite2[of z2 z1]

```

```

    by auto
ultimately
show ?thesis
    by simp
next
case False
have z1 z2 > 0
    using False
    by (metis less-linear less-trans pi-half-gt-zero)
have a z1 z2 = pi - z1 z2
    using False ⟨ z1 z2 > 0 ⟩
    unfolding ang-vec-a-def ang-vec-c-def ang-vec-a-def abs-real-def
    by auto
moreover
have a z1 (-z2) = pi - z1 z2
    unfolding ang-vec-a-def ang-vec-c-def abs-real-def
    using False ⟨ z1 z2 > 0 ⟩
    using canon-ang-plus-pi1[of z1 z2]
    using canon-ang(1)[of arg z2 - arg z1] canon-ang(2)[of arg z2 - arg z1]
    using ⟨z2 ≠ 0⟩ ang-vec-opposite2[of z2 z1]
    by auto
ultimately
show ?thesis
    by auto
qed
qed
qed
qed

lemma ang-vec-a-opposite1:
a z1 z2 = a (-z1) z2
using ang-vec-a-sym[of -z1 z2] ang-vec-a-opposite2[of z2 z1] ang-vec-a-sym[of z2 z1]
by auto

lemma ang-vec-a-scale1:
assumes k ≠ 0
shows a (complex-of-real k * z1) z2 = a z1 z2
proof (cases k > 0)
case True
thus ?thesis
    unfolding ang-vec-a-def ang-vec-c-def ang-vec-def
    using arg-mult-real-positive[of k z1]
    by auto
next
case False
hence k < 0
using assms
by auto

```

```

thus ?thesis
  using arg-mult-real-negative[of k z1]
  using ang-vec-a-opposite1[of z1 z2]
  unfolding ang-vec-a-def ang-vec-c-def ang-vec-def
  by simp
qed

lemma ang-vec-a-scale2:
  assumes k ≠ 0
  shows a z1 (complex-of-real k * z2) = a z1 z2
  using ang-vec-a-sym[of z1 complex-of-real k * z2]
  using ang-vec-a-scale1[OF assms, of z2 z1]
  using ang-vec-a-sym[of z1 z2]
  by auto

lemma ang-vec-a-scale:
  assumes k1 ≠ 0 k2 ≠ 0
  shows a (complex-of-real k1 * z1) (complex-of-real k2 * z2) = a z1 z2
  using ang-vec-a-scale1[OF assms(1)] ang-vec-a-scale2[OF assms(2)]
  by auto

lemma ang-a-cnj-cnj:
  shows a z1 z2 = a (cnj z1) (cnj z2)
  unfolding ang-vec-a-def ang-vec-c-def ang-vec-def
  proof(cases arg z1 ≠ pi ∧ arg z2 ≠ pi)
    case True
    thus acute-ang ||arg z2 - arg z1|| = acute-ang ||arg (cnj z2) - arg (cnj z1)||
      using arg-cnj2[of z1] arg-cnj2[of z2]
      apply (auto simp del:acute-ang-def)
      proof(cases |arg z2 - arg z1| = pi)
        case True
        thus acute-ang ||arg z2 - arg z1|| = acute-ang ||- arg z2 + arg z1||
          using canon-ang-uminus-pi[of arg z2 - arg z1]
          by (auto simp add:field-simps del:acute-ang-def)
      qed
    next
      case False
      thus acute-ang ||arg z2 - arg z1|| = acute-ang ||- arg z2 + arg z1||
        using canon-ang-uminus[of arg z2 - arg z1]
        by (auto simp add:field-simps del:acute-ang-def)
    qed
  next
    case False
    thus acute-ang ||arg z2 - arg z1|| = acute-ang ||arg (cnj z2) - arg (cnj z1)||
      proof(cases arg z1 = pi)
        case False
        hence arg z2 = pi
          using ⟨¬ (arg z1 ≠ pi ∧ arg z2 ≠ pi)⟩
          by auto
        thus ?thesis
      qed
    qed
  qed

```

```

using False
using arg-cnj2[of z1] arg-cnj1[of z2]
apply (auto simp del:acute-ang-def)
proof(cases arg z1 > 0)
  case True
  hence -arg z1 ≤ 0
    by auto
  thus acute-ang ||pi - arg z1|| = acute-ang ||pi + arg z1||
    using True MoreComplex.canon-ang-plus-pi1[of arg z1]
    using arg-bounded[of z1] MoreComplex.canon-ang-plus-pi2[of -arg z1]
    by (auto simp add:field-simps del:acute-ang-def)
next
  case False
  hence -arg z1 ≥ 0
    by simp
  thus acute-ang ||pi - arg z1|| = acute-ang ||pi + arg z1||
  proof(cases arg z1 = 0)
    case True
    thus ?thesis
      by (auto simp del:acute-ang-def)
  next
    case False
    hence -arg z1 > 0
      using ⟨-arg z1 ≥ 0⟩
      by auto
    thus ?thesis
      using False MoreComplex.canon-ang-plus-pi1[of -arg z1]
      using arg-bounded[of z1] MoreComplex.canon-ang-plus-pi2[of arg z1]
      by (auto simp add:field-simps del:acute-ang-def)
  qed
qed
next
  case True
  thus ?thesis
    using arg-cnj1[of z1]
    apply (auto simp del:acute-ang-def)
  proof(cases arg z2 = pi)
    case True
    thus acute-ang ||arg z2 - pi|| = acute-ang ||arg (cnj z2) - pi||
      using arg-cnj1[of z2]
      by auto
  next
    case False
    thus acute-ang ||arg z2 - pi|| = acute-ang ||arg (cnj z2) - pi||
    using arg-cnj2[of z2]
      apply (auto simp del:acute-ang-def)
  proof(cases arg z2 > 0)
    case True
    hence -arg z2 ≤ 0

```

```

    by auto
thus acute-ang ||arg z2 - pi|| = acute-ang ||- arg z2 - pi||
  using True canon-ang-minus-pi1[of arg z2]
  using arg-bounded[of z2] canon-ang-minus-pi2[of -arg z2]
  by (auto simp add:field-simps del:acute-ang-def)
next
  case False
  hence -arg z2 ≥ 0
    by simp
  thus acute-ang ||arg z2 - pi|| = acute-ang ||- arg z2 - pi||
  proof(cases arg z2 = 0)
    case True
    thus ?thesis
      by (auto simp del:acute-ang-def)
  next
    case False
    hence -arg z2 > 0
      using (-arg z2 ≥ 0)
      by auto
    thus ?thesis
      using False canon-ang-minus-pi1[of -arg z2]
      using arg-bounded[of z2] canon-ang-minus-pi2[of arg z2]
      by (auto simp add:field-simps del:acute-ang-def)
qed
qed
qed
qed
qed

```

abbreviation *sgn-bool* **where**
 $\text{sgn-bool } p \equiv \text{if } p \text{ then } 1 \text{ else } -1$

definition *circ-tang-vec* :: *complex* \Rightarrow *complex* \Rightarrow *bool* \Rightarrow *complex* **where**
 $\text{circ-tang-vec } \mu E p = \text{sgn-bool } p * ii * (E - \mu)$

lemma *circ-tang-vec-ortho*:
 $\text{scalprod } (E - \mu) (\text{circ-tang-vec } \mu E p) = 0$
unfolding *circ-tang-vec-def* *Let-def*
by (auto simp add: complex-cnj-mult)

lemma *circ-tang-vec-opposite-orient*:
 $\text{circ-tang-vec } \mu E p = - \text{circ-tang-vec } \mu E (\neg p)$
unfolding *circ-tang-vec-def*
by auto

definition *ang-circ* **where**
 $\text{ang-circ } E \mu1 \mu2 p1 p2 = (\text{circ-tang-vec } \mu1 E p1) (\text{circ-tang-vec } \mu2 E p2)$

```

definition ang-circ-c where
  ang-circ-c E μ1 μ2 p1 p2 = c (circ-tang-vec μ1 E p1) (circ-tang-vec μ2 E p2)

definition ang-circ-a where
  ang-circ-a E μ1 μ2 p1 p2 = a (circ-tang-vec μ1 E p1) (circ-tang-vec μ2 E p2)

lemma ang-circ-simp:
  assumes E ≠ μ1 E ≠ μ2
  shows ang-circ E μ1 μ2 p1 p2 = canon-ang (arg (E - μ2) - arg (E - μ1) +
  sgn-bool p1 * pi / 2 - sgn-bool p2 * pi / 2)
  unfolding ang-circ-def ang-vec-def circ-tang-vec-def
  apply (rule canon-ang-eq)
  using assms
  using arg-mult-2kpi[of sgn-bool p2*ii E - μ2]
  using arg-mult-2kpi[of sgn-bool p1*ii E - μ1]
  apply auto
  apply (rule-tac x=x-xa in exI, auto simp add: field-simps)
  apply (rule-tac x=-1+x-xa in exI, auto simp add: field-simps)
  apply (rule-tac x=1+x-xa in exI, auto simp add: field-simps)
  apply (rule-tac x=x-xa in exI, auto simp add: field-simps)
  done

lemma ang-circ-c-simp:
  assumes E ≠ μ1 E ≠ μ2
  shows ang-circ-c E μ1 μ2 p1 p2 = abs (canon-ang (arg(E - μ2) - arg(E -
  μ1) + (sgn-bool p1) * pi/2 - (sgn-bool p2) * pi/2))
  unfolding ang-circ-c-def ang-vec-c-def
  using ang-circ-simp[OF assms]
  unfolding ang-circ-def
  by auto

lemma ang-circ-a-simp:
  assumes E ≠ μ1 E ≠ μ2
  shows ang-circ-a E μ1 μ2 p1 p2 = acute-ang (abs (canon-ang (arg(E - μ2) -
  arg(E - μ1) + (sgn-bool p1) * pi/2 - (sgn-bool p2) * pi/2)))
  unfolding ang-circ-a-def ang-vec-a-def
  using ang-circ-c-simp[OF assms]
  unfolding ang-circ-c-def
  by auto

lemma ang-circ-a-pTrue:
  assumes E ≠ μ1 E ≠ μ2
  shows ang-circ-a E μ1 μ2 p1 p2 = ang-circ-a E μ1 μ2 True True
  proof (cases p1)
    case True
    show ?thesis
  proof (cases p2)
    case True
    show ?thesis

```

```

using ⟨p1⟩ ⟨p2⟩
by simp
next
case False
show ?thesis
  using ⟨p1⟩ ⊥⟨p2⟩
  unfolding ang-circ-a-def
  using circ-tang-vec-opposite-orient[of μ2 E p2]
  using ang-vec-a-opposite2
  by simp
qed
next
case True
show ?thesis
proof (cases p2)
  case True
  show ?thesis
    using ⊥⟨p1⟩ ⟨p2⟩
    unfolding ang-circ-a-def
    using circ-tang-vec-opposite-orient[of μ1 E p1]
    using ang-vec-a-opposite1
    by simp
  next
  case False
  show ?thesis
    using ⊥⟨p1⟩ ⊥⟨p2⟩
    unfolding ang-circ-a-def
    using circ-tang-vec-opposite-orient[of μ1 E p1] circ-tang-vec-opposite-orient[of
μ2 E p2]
    using ang-vec-a-opposite1 ang-vec-a-opposite2
    by simp
  qed
qed

lemma ang-circ-a-simp1:
assumes E ≠ μ1 E ≠ μ2
shows ang-circ-a E μ1 μ2 p1 p2 = a (E - μ1) (E - μ2)
unfolding ang-vec-a-def ang-vec-c-def ang-vec-def
by (subst ang-circ-a-pTrue[OF assms, of p1 p2], subst ang-circ-a-simp[OF assms,
of True True]) (metis add-diff-cancel)

abbreviation ang-circ-a' where
ang-circ-a' E μ1 μ2 ≡ ang-circ-a E μ1 μ2 True True

lemma ang-circ-a'-simp:
assumes z ≠ μ1 z ≠ μ2
shows ang-circ-a' z μ1 μ2 = a (z - μ1) (z - μ2)
by (rule ang-circ-a-simp1[OF assms])

```

```

lemma cos-cmod-scalprod:
  shows cmod b * cmod c * (cos ( b c)) = Re (scalprod b c)
  proof (cases b = 0 ∨ c = 0)
    case True
    thus ?thesis
      by auto
  next
    case False
    thus ?thesis
      by (simp add: cos-diff cos-arg sin-arg field-simps)
  qed

lemma law-of-cosines:
  shows (cdist B C)2 = (cdist A C)2 + (cdist A B)2 - 2*(cdist A C)*(cdist A B)*(cos ((C-A) (B-A)))
  proof-
    let ?a = C-B and ?b = C-A and ?c = B-A
    have ?a = ?b - ?c
      by simp
    hence (cmod ?a)2 = (cmod (?b - ?c))2
      by metis
    also have ... = Re (scalprod (?b - ?c) (?b - ?c))
      by (simp add: cmod-square)
    also have ... = (cmod ?b)2 + (cmod ?c)2 - 2*Re (scalprod ?b ?c)
      by (simp add: cmod-square field-simps)
    finally
      show ?thesis
        using cos-cmod-scalprod[of ?b ?c]
        by simp
  qed

declare ang-vec-c-def[simp del]

lemma cos-c: cos (c z1 z2) = cos ( z1 z2)
unfolding ang-vec-c-def
by (smt cos-minus)

lemma cos-a-c: cos (a z1 z2) = abs (cos (c z1 z2))
unfolding ang-vec-a-def
using ang-vec-c-bounded[of z1 z2] cos-lt-zero[of c z1 z2] cos-gt-zero-pi[of c z1 z2]
by (cases c z1 z2 = pi/2) (auto, smt cos-minus cos-periodic-pi3)

end

```

8 Homogeneous coordinates in extended complex plane

```

theory HomogeneousCoordinates
imports MoreComplex Matrices
begin

typedef homo-coords = {v. v ≠ vec-zero}
by (rule-tac x=(1, 0) in exI, simp)

lemma obtain-homo-coords:
fixes x::homo-coords
obtains A B where
Rep-homo-coords x = (A, B) A ≠ 0 ∨ B ≠ 0
by (cases x) (auto simp add: Abs-homo-coords-inverse)

definition homo-coords-eq :: homo-coords ⇒ homo-coords ⇒ bool (infix ≈ 50)
where
[simp]: z1 ≈ z2 ↔
(let z1 = Rep-homo-coords z1;
z2 = Rep-homo-coords z2
in (∃ k. k ≠ (0::complex) ∧ z2 = k *sv z1))

lemma homo-coords-eq-reflp:
reflp homo-coords-eq
by (auto simp add: reflp-def, rule-tac x=1 in exI, simp)

lemma homo-coords-eq-symp:
symp homo-coords-eq
by (auto simp add: symp-def, rule-tac x=1/k in exI, simp)

lemma homo-coords-eq-transp:
transp homo-coords-eq
by (auto simp add: transp-def, rule-tac x=ka*k in exI, simp)

lemma homo-coords-eq-equivp:
equivp homo-coords-eq
by (auto intro: equivpI homo-coords-eq-reflp homo-coords-eq-symp homo-coords-eq-transp)

lemma homo-coords-eq-refl [simp]:
z ≈ z
using homo-coords-eq-reflp
by (auto simp add: reflp-def refl-on-def)

lemma homo-coords-eq-trans:
assumes z1 ≈ z2 z2 ≈ z3
shows z1 ≈ z3
using assms homo-coords-eq-transp
unfolding transp-def

```

by *blast*

lemma *homo-coords-eq-sym*:

assumes $z1 \approx z2$

shows $z2 \approx z1$

using *assms homo-coords-eq-symp*

unfolding *symp-def*

by *blast*

lemma *homo-coords-eq-mix*:

assumes *Rep-homo-coords* $z1 = (z1', z1'')$ *Rep-homo-coords* $z2 = (z2', z2'')$

shows $z1 \approx z2 \longleftrightarrow z2'*z1'' = z1'*z2''$

using *assms*

proof (cases $z1'' \neq 0 \vee z2'' \neq 0$)

case *False*

thus *?thesis*

using *assms* using *Rep-homo-coords*[of $z1$] *Rep-homo-coords*[of $z2$]

by *auto*

next

case *True*

thus *?thesis*

using *assms*

apply *auto*

apply (rule-tac $x=z2''/z1''$ in *exI*)

using *Rep-homo-coords*[of $z2$]

apply (auto simp add: *field-simps*)

apply (rule-tac $x=z2''/z1''$ in *exI*)

using *Rep-homo-coords*[of $z1$]

apply (auto simp add: *field-simps*)

done

qed

lemma [*simp*]: *Rep-homo-coords* (*Abs-homo-coords* (*Rep-homo-coords* x)) = *Rep-homo-coords* x

using *Rep-homo-coords*[of x]

by (*simp add: Abs-homo-coords-inverse*)

Quotient of homogeneous coordinates

quotient-type

complex-homo = *homo-coords* / *homo-coords-eq*

by (rule *homo-coords-eq-equivp*)

Infinite point

definition *inf-homo-rep* **where** [*simp*]: *inf-homo-rep* = *Abs-homo-coords* (1, 0)

lift-definition *inf-homo* :: *complex-homo* (∞_h) **is** *inf-homo-rep*

done

lemma [*simp*]: *Rep-homo-coords* (*Abs-homo-coords* (1, 0)) = (1, 0)
by (*simp add: Abs-homo-coords-inverse*)

```

lemma [simp]: Rep-homo-coords inf-homo-rep = (1, 0)
by simp

lemma inf-snd-0: z ≈ inf-homo-rep  $\longleftrightarrow$  (let (z1, z2) = Rep-homo-coords z in z1
≠ 0  $\wedge$  z2 = 0)
using Rep-homo-coords[of z]
by auto

lemma not-inf-snd-not0:
assumes  $\neg$  z ≈ inf-homo-rep
shows let (z1, z2) = Rep-homo-coords z in z2 ≠ 0
using assms Rep-homo-coords[of z] inf-snd-0[of z]
by auto

Zero

definition zero-homo-rep where [simp]: zero-homo-rep = Abs-homo-coords (0, 1)
lift-definition zero-homo :: complex-homo (0h) is zero-homo-rep
done

lemma [simp]: Rep-homo-coords (Abs-homo-coords (0, 1)) = (0, 1)
by (simp add: Abs-homo-coords-inverse)

lemma [simp]: Rep-homo-coords zero-homo-rep = (0, 1)
by simp

lemma zero-fst-0: z ≈ zero-homo-rep  $\longleftrightarrow$  (let (z1, z2) = Rep-homo-coords z in
z1 = 0  $\wedge$  z2 ≠ 0)
using Rep-homo-coords[of z]
by auto

One

definition one-homo-rep where [simp]: one-homo-rep = Abs-homo-coords (1, 1)
lift-definition one-homo :: complex-homo (1h) is one-homo-rep
done

lemma [simp]: Rep-homo-coords (Abs-homo-coords (1, 1)) = (1, 1)
by (simp add: Abs-homo-coords-inverse)

lemma [simp]: Rep-homo-coords one-homo-rep = (1, 1)
by simp

lemma [simp]: 1h ≠ ∞h 0h ≠ ∞h 0h ≠ 1h 1h ≠ 0h ∞h ≠ 0h ∞h ≠ 1h
by (transfer, auto)+
```

definition ii-homo-rep **where** ii-homo-rep = Abs-homo-coords (ii, 1)

lift-definition *ii-homo* :: *complex-homo* (*ii_h*) **is** *ii-homo-rep*
done

lemma [*simp*]: *Rep-homo-coords* (*Abs-homo-coords* (*ii*, 1)) = (*ii*, 1)
by (*simp add: Abs-homo-coords-inverse*)

lemma [*simp*]: *Rep-homo-coords ii-homo-rep* = (*ii*, 1)
by (*simp add: ii-homo-rep-def*)

lemma *ex-3-different-points*:
fixes *z::complex-homo*
shows $\exists z1 z2. z \neq z1 \wedge z1 \neq z2 \wedge z \neq z2$
proof (*cases z ≠ 0_h ∧ z ≠ 1_h*)
case *True*
thus *?thesis*
by (*rule-tac x=0_h in exI, rule-tac x=1_h in exI, auto*)
next
case *False*
hence $z = 0_h \vee z = 1_h$
by *simp*
thus *?thesis*
proof
assume $z = 0_h$
thus *?thesis*
by (*rule-tac x=∞_h in exI, rule-tac x=1_h in exI, auto*)
next
assume $z = 1_h$
thus *?thesis*
by (*rule-tac x=∞_h in exI, rule-tac x=0_h in exI, auto*)
qed
qed

Conversion from complex

definition *of-complex-coords where*
of-complex-coords z = Abs-homo-coords (z, 1)

lemma [*simp*]: *Rep-homo-coords (of-complex-coords z) = (z, 1)*
by (*simp add: of-complex-coords-def Abs-homo-coords-inverse*)

lift-definition *of-complex* :: *complex* \Rightarrow *complex-homo* **is** *of-complex-coords*
by (*simp del: homo-coords-eq-def*)

lemma *of-complex-inj*:
assumes *of-complex x = of-complex y*
shows *x = y*
using assms
by *transfer simp*

```

lemma of-complex-image-inj:
  assumes of-complex ‘A = of-complex ‘B
  shows A = B
  using assms
  using of-complex-inj
  by auto

lemma [simp]: of-complex x ≠ ∞h
  by transfer simp

lemma [simp]: ∞h ≠ of-complex x
  by transfer simp

lemma inf-homo-or-complex-homo:
  z = ∞h ∨ (∃ x. z = of-complex x)
  proof(transfer)
    fix z
    obtain a b where *: Rep-homo-coords z = (a, b)
      by (rule obtain-homo-coords)
    show z ≈ inf-homo-rep ∨ (∃ x. z ≈ of-complex-coords x)
      using * Rep-homo-coords[of z]
      by (cases b = 0) auto
  qed

lemma zero-of-complex [simp]: of-complex 0 = 0h
  by transfer simp

lemma one-of-complex [simp]: of-complex 1 = 1h
  by transfer simp

lemma
  [simp]: of-complex a = 0h ↔ a = 0
  by (subst zero-of-complex[symmetric]) (auto simp add: of-complex-inj)

lemma
  [simp]: of-complex a = 1h ↔ a = 1
  by (subst one-of-complex[symmetric]) (auto simp add: of-complex-inj)

Coercion to complex

definition to-complex-homo-coords :: homo-coords ⇒ complex where
  to-complex-homo-coords z = (let (z1, z2) = Rep-homo-coords z in z1/z2)

lift-definition to-complex :: complex-homo ⇒ complex is to-complex-homo-coords
proof-
  fix x y
  assume x ≈ y
  thus to-complex-homo-coords x = to-complex-homo-coords y
    by (auto simp add: to-complex-homo-coords-def split-def Let-def)

```

qed

lemma [simp]: *to-complex (of-complex z) = z*
by (transfer) (simp add: of-complex-coords-def to-complex-homo-coords-def Abs-homo-coords-inverse)

lemma [simp]: $z \neq \infty_h \implies (\text{of-complex}(\text{to-complex } z)) = z$

proof (transfer)

fix z

obtain $z1 z2$ where $zz: \text{Rep-homo-coords } z = (z1, z2)$

by (rule obtain-homo-coords)

assume $\neg z \approx \text{inf-homo-rep}$

hence $z2 \neq 0$

using $zz \text{ Rep-homo-coords}[of } z]$

by auto (erule-tac $x=1/z1$ in allE, simp)

thus $\text{of-complex-coords}(\text{to-complex-homo-coords } z) \approx z$

using zz

by (auto simp add: of-complex-coords-def to-complex-homo-coords-def Abs-homo-coords-inverse)

qed

Addition

definition add-homo-coords :: homo-coords \Rightarrow homo-coords \Rightarrow homo-coords (**infixl**
 $+_{hc}$ 100) **where**
 $z +_{hc} w = (\text{let } (z1, z2) = \text{Rep-homo-coords } z;$
 $(w1, w2) = \text{Rep-homo-coords } w \text{ in}$
 $\text{Abs-homo-coords } (z1*w2 + w1*z2, z2*w2))$

lemma add-homo-coords-Rep:

assumes $\text{Rep-homo-coords } z = (z1, z2) \text{ Rep-homo-coords } w = (w1, w2) z2 \neq 0$
 $\vee w2 \neq 0$

shows $\text{Rep-homo-coords}(z +_{hc} w) = (z1*w2 + w1*z2, z2*w2)$

proof –

from assms

have $(z1*w2 + w1*z2, z2*w2) \neq \text{vec-zero}$

using $\text{Rep-homo-coords}[of } z] \text{ Rep-homo-coords}[of } w]$

by auto

thus ?thesis

using assms(1–2)

by (auto simp add: add-homo-coords-def split-def Let-def Abs-homo-coords-inverse)

qed

lemma add-homo-coords-00:

assumes $\text{Rep-homo-coords } z = (z1, z2) \text{ Rep-homo-coords } w = (w1, w2) z2 = 0$
 $w2 = 0$

shows $z +_{hc} w = \text{Abs-homo-coords } (0, 0)$

using assms unfolding add-homo-coords-def

by simp

lemma add-coords-well-defined-lemma:

assumes $x \approx y$ $x' \approx y'$

```

shows  $x +_{hc} x' \approx y +_{hc} y'$ 
using assms
proof-
  obtain  $Ax Bx$  where  $xx: Rep\text{-homo-coords } x = (Ax, Bx)$ 
    by (rule obtain-homo-coords)
  obtain  $Ax' Bx'$  where  $xx': Rep\text{-homo-coords } x' = (Ax', Bx')$ 
    by (rule obtain-homo-coords)
  obtain  $Ay By$  where  $yy: Rep\text{-homo-coords } y = (Ay, By)$ 
    by (rule obtain-homo-coords)
  obtain  $Ay' By'$  where  $yy': Rep\text{-homo-coords } y' = (Ay', By')$ 
    by (rule obtain-homo-coords)
  from assms obtain  $k k'$  where
     $*: k \neq 0 \ Ay = k * Ax \ By = k * Bx \ k' \neq 0 \ Ay' = k' * Ax' \ By' = k' * Bx'$ 
    using  $xx \ xx' \ yy \ yy'$ 
    by auto
  show ?thesis
  proof (cases  $Bx = 0 \wedge Bx' = 0$ )
    case True
    thus ?thesis
      using add-homo-coords-00[of  $x Ax 0 x' Ax' 0$ ] add-homo-coords-00[of  $y Ay 0 y' Ay' 0$ ]  $xx \ yy \ xx' \ yy' *$ 
        by (auto, rule-tac  $x=1$  in exI, simp)
  next
    case False
    thus ?thesis
      using  $xx \ xx' \ yy \ yy' *$ 
      using Rep-homo-coords[of  $x$ ] Rep-homo-coords[of  $x$ ]  $\langle k \neq 0 \rangle \langle k' \neq 0 \rangle$ 
      using add-homo-coords-Rep[of  $x Ax Bx x' Ax' Bx'$ ] add-homo-coords-Rep[of  $y k * Ax k * Bx y' k' * Ax' k' * Bx'$ ]
        by simp (rule-tac  $x=k*k'$  in exI, auto simp add: field-simps)
  qed
qed

```

lift-definition $add\text{-homo} :: complex\text{-homo} \Rightarrow complex\text{-homo} \Rightarrow complex\text{-homo}$ (**infixl**
 $+_h 100$) **is** $add\text{-homo-coords}$
by (rule add-coords-well-defined-lemma, simp-all)

```

lemma  $add\text{-homo-commute}: x +_h y = y +_h x$ 
proof (transfer)
  fix  $x y$ 
  obtain  $Ax Bx$  where  $xx: Rep\text{-homo-coords } x = (Ax, Bx)$ 
    by (rule obtain-homo-coords)
  obtain  $Ay By$  where  $yy: Rep\text{-homo-coords } y = (Ay, By)$ 
    by (rule obtain-homo-coords)

  show  $x +_{hc} y \approx y +_{hc} x$ 
  proof (cases  $Bx \neq 0 \vee By \neq 0$ )
    case True
    thus ?thesis

```

```

using add-homo-coords-Rep[of x Ax Bx y Ay By, OF xx yy]
using add-homo-coords-Rep[of y Ay By x Ax Bx, OF yy xx]
by auto (rule-tac x=1 in exI, simp) +
next
  case False
  thus ?thesis
    using xx yy add-homo-coords-00
    by (auto, rule-tac x=1 in exI, simp)
qed
qed

lemma of-complex-add: (of-complex za) +h (of-complex zb) = of-complex (za + zb)
proof (transfer)
  fix za zb
  have Rep-homo-coords (Abs-homo-coords (za, 1)) = (za, 1) Rep-homo-coords
  (Abs-homo-coords (zb, 1)) = (zb, 1)
    by (auto simp add: Abs-homo-coords-inverse)
  thus of-complex-coords za +hc of-complex-coords zb ≈ of-complex-coords (za + zb)
    unfolding of-complex-coords-def
    using add-homo-coords-Rep[of Abs-homo-coords (za, 1) za 1 Abs-homo-coords
    (zb, 1) zb 1]
    by (simp add: Abs-homo-coords-inverse)
qed

lemma [simp]: (of-complex z) +h ∞h = ∞h
proof (transfer)
  fix z
  show of-complex-coords z +hc inf-homo-rep ≈ inf-homo-rep
    using add-homo-coords-Rep[of Abs-homo-coords (z, 1) z 1 Abs-homo-coords (1, 0) 1 0]
    unfolding of-complex-coords-def
    by (simp add: Abs-homo-coords-inverse)
qed

lemma [simp]: ∞h +h (of-complex z) = ∞h
by (subst add-homo-commute) simp

lemma add-homo-zero-right [simp]: z +h 0h = z
proof (transfer)
  fix z
  obtain z1 z2 where zz: Rep-homo-coords z = (z1, z2)
    by (rule obtain-homo-coords)
  thus z +hc zero-homo-rep ≈ z
    using add-homo-coords-Rep[of z z1 z2 zero-homo-rep 0 1]
    by auto (metis zero-neq-one)
qed

```

```

lemma add-homo-zero-left [simp]:  $0_h +_h z = z$ 
  by (subst add-homo-commute) simp

uminus

definition uminus-homo-coords where
  uminus-homo-coords  $z = (\text{let } (z1, z2) = \text{Rep-homo-coords } z \text{ in } \text{Abs-homo-coords } (-z1, z2))$ 

lemma uminus-homo-coords-Rep [simp]: Rep-homo-coords (uminus-homo-coords  $z) = (\text{let } (z1, z2) = \text{Rep-homo-coords } z \text{ in } (-z1, z2))$ 
  unfolding uminus-homo-coords-def Let-def
  apply (cases Rep-homo-coords  $z)$ 
  using Rep-homo-coords[of  $z)$ 
  by (auto simp add: Abs-homo-coords-inverse)

lift-definition uminus-homo :: complex-homo  $\Rightarrow$  complex-homo is uminus-homo-coords
  by (auto simp add: split-def Let-def)

lemma of-complex-uminus [simp]: uminus-homo (of-complex  $z) = \text{of-complex } (-z)$ 
  by (transfer) auto

Subtraction

definition minus-homo :: complex-homo  $\Rightarrow$  complex-homo  $\Rightarrow$  complex-homo (infixl
 $-_h 100$ ) where
   $z1 -_h z2 = z1 +_h (\text{uminus-homo } z2)$ 

lemma minus-homo-coords-Rep:
  assumes Rep-homo-coords  $z = (z1, z2)$  Rep-homo-coords  $w = (w1, w2)$   $z2 \neq 0$ 
   $\vee w2 \neq 0$ 
  shows Rep-homo-coords  $(z +_{hc} (\text{uminus-homo-coords } w)) = (z1*w2 - w1*z2,$ 
 $z2*w2)$ 
  using assms
  using add-homo-coords-Rep[of  $z z1 z2$  uminus-homo-coords  $w - w1 w2] uminus-homo-coords-Rep[of
 $w)$ 
  by simp

lemma of-complex-minus:
   $(\text{of-complex } z1) -_h (\text{of-complex } z2) = \text{of-complex } (z1 - z2)$ 
  unfolding minus-homo-def complex-diff-def
  by (simp add: of-complex-add)

lemma [simp]:
  assumes  $z \neq \infty_h$ 
  shows  $z -_h z = 0_h$ 
proof-
  from assms obtain  $z'$  where  $z = \text{of-complex } z'$ 
  using inf-homo-or-complex-homo[of  $z)$ 
  by auto
  thus ?thesis$ 
```

```

    by (simp add: of-complex-minus)
qed

lemma diff-zero-homo:
assumes z1 -_h z2 = 0_h z1 ≠ ∞_h ∨ z2 ≠ ∞_h
shows z1 = z2
using assms
unfolding minus-homo-def
proof transfer
fix z w
obtain z1 z2 where zz: Rep-homo-coords z = (z1, z2)
  by (rule obtain-homo-coords)
obtain w1 w2 where ww: Rep-homo-coords w = (w1, w2)
  by (rule obtain-homo-coords)
have mww: Rep-homo-coords (uminus-homo-coords w) = (-w1, w2)
  using ww
  by simp
assume *: z +_hc uminus-homo-coords w ≈ zero-homo-rep and
  ¬ z ≈ inf-homo-rep ∨ ¬ w ≈ inf-homo-rep
have z2 ≠ 0 ∨ w2 ≠ 0
  using Rep-homo-coords[of z] Rep-homo-coords[of w]
  using ¬ z ≈ inf-homo-rep ∨ ¬ w ≈ inf-homo-rep
  using inf-snd-0[of z] inf-snd-0[of w] zz ww
  by auto
thus z ≈ w
  using * zz ww
  apply simp
  apply (subst (asm) minus-homo-coords-Rep[of z z1 z2 w w1 w2])
  apply auto
  apply (rule-tac x=w2/z2 in exI, auto simp add: field-simps)
  apply (rule-tac x=w2/z2 in exI, auto)
done
qed

```

Multiplication

```

definition mult-homo-coords :: homo-coords ⇒ homo-coords ⇒ homo-coords (infixl
*_hc 100) where
x *_hc y = (let (x1, y1) = Rep-homo-coords x;
             (x2, y2) = Rep-homo-coords y in
             Abs-homo-coords (x1*x2, y1*y2))

lemma mult-homo-coords-Rep:
assumes Rep-homo-coords x = (Ax, Bx) Rep-homo-coords x' = (Ax', Bx') (Bx
≠ 0 ∨ Ax' ≠ 0) ∧ (Bx' ≠ 0 ∨ Ax ≠ 0)
shows Rep-homo-coords (x *_hc x') = (Ax*Ax', Bx*Bx')
using assms Rep-homo-coords[of x] Rep-homo-coords[of x']
by (auto simp add: mult-homo-coords-def split-def Let-def Abs-homo-coords-inverse)

lemma mult-homo-coords-00:

```

```

assumes Rep-homo-coords  $x = (Ax, Bx)$  Rep-homo-coords  $x' = (Ax', Bx')$  ( $Bx = 0 \wedge Ax' = 0$ )  $\vee (Bx' = 0 \wedge Ax = 0)$ 
shows  $x *_{hc} x' = \text{Abs-homo-coords } (0, 0)$ 
using assms unfolding mult-homo-coords-def
by auto

lemma mult-coords-well-defined-lemma:
assumes  $x \approx y$   $x' \approx y'$ 
shows  $x *_{hc} x' \approx y *_{hc} y'$ 
proof-
obtain Ax Bx where xx: Rep-homo-coords  $x = (Ax, Bx)$ 
by (rule obtain-homo-coords)
obtain Ax' Bx' where xx': Rep-homo-coords  $x' = (Ax', Bx')$ 
by (rule obtain-homo-coords)
obtain Ay By where yy: Rep-homo-coords  $y = (Ay, By)$ 
by (rule obtain-homo-coords)
obtain Ay' By' where yy': Rep-homo-coords  $y' = (Ay', By')$ 
by (rule obtain-homo-coords)
from assms obtain k k' where
*:  $k \neq 0$   $Ay = k * Ax$   $By = k * Bx$   $k' \neq 0$   $Ay' = k' * Ax'$   $By' = k' * Bx'$ 
using xx xx' yy yy'
by auto
show ?thesis
proof (cases ( $Bx \neq 0 \vee Ax' \neq 0$ )  $\wedge (Bx' \neq 0 \vee Ax \neq 0)$ )
case False
thus ?thesis
using mult-homo-coords-00[of x Ax Bx x' Ax' Bx'] mult-homo-coords-00[of y
Ay By y' Ay' By'] xx yy xx' yy' *
by auto (rule-tac x=1 in exI, simp) +
next
case True
thus ?thesis
using xx xx' yy yy' *
using Rep-homo-coords[of x] Rep-homo-coords[of x']  $\langle k \neq 0 \rangle$   $\langle k' \neq 0 \rangle$ 
using mult-homo-coords-Rep[of x Ax Bx x' Ax' Bx']
mult-homo-coords-Rep[of y k * Ax k * Bx y' k' * Ax' k' * Bx']
by simp (rule-tac x=k*k' in exI, auto simp add: field-simps)
qed
qed

lift-definition mult-homo :: complex-homo  $\Rightarrow$  complex-homo  $\Rightarrow$  complex-homo
(infixl  $*_h$  100) is mult-homo-coords
by (rule mult-coords-well-defined-lemma, simp-all)

lemma mult-of-complex:
shows (of-complex z1)  $*_h$  (of-complex z2) = of-complex (z1 * z2)
proof (transfer)
fix z1 z2
show of-complex-coords  $z1 *_{hc}$  of-complex-coords  $z2 \approx$  of-complex-coords (z1 *

```

```

z2)
  using mult-homo-coords-Rep[of of-complex-coords z1 - - of-complex-coords z2]
  by simp
qed

lemma mult-homo-commute:
  shows  $z1 *_h z2 = z2 *_h z1$ 
proof transfer
  fix  $z1 z2$ 
  obtain  $z11 z12$  where  $z1: \text{Rep-homo-coords } z1 = (z11, z12)$ 
    by (rule obtain-homo-coords)
  obtain  $z21 z22$  where  $z2: \text{Rep-homo-coords } z2 = (z21, z22)$ 
    by (rule obtain-homo-coords)
  show  $z1 *_{hc} z2 \approx z2 *_{hc} z1$ 
  proof (cases  $(z12 \neq 0 \vee z21 \neq 0) \wedge (z22 \neq 0 \vee z11 \neq 0)$ )
    case True
    thus ?thesis
      using mult-homo-coords-Rep[of  $z1 z11 z12 z2 z21 z22$ ]  $z1 z2$ 
      using mult-homo-coords-Rep[of  $z2 z21 z22 z1 z11 z12$ ]
      by simp (rule-tac  $x=1$  in exI, simp)
  next
    case False
    thus ?thesis
      using mult-homo-coords-00[of  $z1 z11 z12 z2 z21 z22$ ]  $z1 z2$ 
      using mult-homo-coords-00[of  $z2 z21 z22 z1 z11 z12$ ]
      by auto (rule-tac  $x=1$  in exI, simp)+
  qed
qed

lemma mult-homo-zero-left [simp]:
  assumes  $z \neq \infty_h$ 
  shows  $0_h *_h z = 0_h$ 
using assms
proof-
  obtain  $z'$  where  $z = \text{of-complex } z'$ 
    using inf-homo-or-complex-homo[of  $z$ ] assms
    by auto
  thus ?thesis
    using zero-of-complex
    using mult-of-complex[of  $0 z'$ ]
    by simp
qed

lemma mult-homo-zero-right [simp]:
  assumes  $z \neq \infty_h$ 
  shows  $z *_h 0_h = 0_h$ 
using mult-homo-zero-left[OF assms]
by (simp add: mult-homo-commute)

```

```

lemma mult-homo-inf-right [simp]:
  assumes  $z \neq 0_h$ 
  shows  $z *_h \infty_h = \infty_h$ 
  using assms
  proof (transfer)
    fix  $z$ 
    obtain  $z1 z2$  where  $Rep\text{-homo-coords } z = (z1, z2)$ 
      by (rule obtain-homo-coords)
    assume  $\neg z \approx zero\text{-homo-rep}$ 
    hence  $z1 \neq 0$ 
      using Rep-homo-coords[of  $z$ ]  $zz$ 
      by auto (metis divide-self-if eq-divide-eq mult-divide-mult-cancel-right)
    thus  $z *_{hc} inf\text{-homo-rep} \approx inf\text{-homo-rep}$ 
      using zz mult-homo-coords-Rep[of  $z z1 z2$  Abs-homo-coords (1, 0) 1 0]
        by auto
  qed

lemma mult-homo-inf-left [simp]:
  assumes  $z \neq 0_h$ 
  shows  $\infty_h *_h z = \infty_h$ 
  using mult-homo-inf-right[OF assms]
  by (simp add: mult-homo-commute)

lemma mult-homo-one-left [simp]:
  shows  $1_h *_h z = z$ 
  proof (transfer)
    fix  $z$ 
    obtain  $z1 z2$  where  $Rep\text{-homo-coords } z = (z1, z2)$ 
      by (rule obtain-homo-coords)
    thus one-homo-rep  $*_{hc} z \approx z$ 
      using mult-homo-coords-Rep[of Abs-homo-coords (1, 1) 1 1 z z1 z2]
        by auto (metis zero-neq-one)
  qed

lemma mult-homo-one-right [simp]:
  shows  $z *_h 1_h = z$ 
  using mult-homo-one-left[of  $z$ ]
  by (simp add: mult-homo-commute)

Reciprocal

definition reciprocal-homo-coords :: homo-coords  $\Rightarrow$  homo-coords where
  reciprocal-homo-coords  $x = (\text{let } (x1, y1) = Rep\text{-homo-coords } x \text{ in } Abs\text{-homo-coords } (y1, x1))$ 

lemma reciprocal-homo-coords-Rep:  $Rep\text{-homo-coords } (\text{reciprocal-homo-coords } x) = (\text{let } (x1, y1) = Rep\text{-homo-coords } x \text{ in } (y1, x1))$ 
  apply (cases Rep-homo-coords  $x$ )
  unfolding reciprocal-homo-coords-def Let-def
  using Rep-homo-coords[of  $x$ ]

```

```

by (auto simp add: Abs-homo-coords-inverse)

lift-definition reciprocal-homo :: complex-homo ⇒ complex-homo is reciprocal-homo-coords
proof-
  fix x y
  assume x ≈ y
  thus reciprocal-homo-coords x ≈ reciprocal-homo-coords y
    by (cases Rep-homo-coords x, cases Rep-homo-coords y) (auto simp add:
      reciprocal-homo-coords-Rep)
qed

lemma [simp]: reciprocal-homo-coords (reciprocal-homo-coords z) = z
  unfolding reciprocal-homo-coords-def [of reciprocal-homo-coords z]
  by (cases Rep-homo-coords z) (auto simp add: reciprocal-homo-coords-Rep, metis
    Rep-homo-coords-inverse)

lemma [simp]: reciprocal-homo (reciprocal-homo z) = z
by (transfer) (auto, rule-tac x=1 in exI, simp)

lemma [simp]: reciprocal-homo 0h = ∞h
by (transfer) (simp add: reciprocal-homo-coords-Rep)

lemma [simp]: reciprocal-homo ∞h = 0h
by (transfer) (simp add: reciprocal-homo-coords-Rep)

lemma [simp]: reciprocal-homo 1h = 1h
by (transfer) (simp add: reciprocal-homo-coords-Rep)

Division

definition divide-homo :: complex-homo ⇒ complex-homo ⇒ complex-homo (infixl
  :h 100) where
  x :h y = x *h (reciprocal-homo y)

lemma [simp]:
  assumes z ≠ 0h
  shows z :h 0h = ∞h
  using assms
  unfolding divide-homo-def
  by simp

lemma [simp]:
  assumes z ≠ ∞h
  shows z :h ∞h = 0h
  using assms
  unfolding divide-homo-def
  by simp

lemma [simp]: ∞h :h 0h = ∞h
  unfolding divide-homo-def

```

```

by (transfer) (simp add: reciprocal-homo-coords-def mult-homo-coords-def)

lemma [simp]:  $0_h :_h \infty_h = 0_h$ 
unfolding divide-homo-def
by (transfer) (simp add: mult-homo-coords-def reciprocal-homo-coords-def)

lemma divide-homo-one [simp]:
  shows  $z :_h 1_h = z$ 
unfolding divide-homo-def
by simp

lemma of-complex-divide:
  assumes  $z2 \neq 0$ 
  shows  $(\text{of-complex } z1) :_h (\text{of-complex } z2) = \text{of-complex } (z1 / z2)$ 
using assms
unfolding divide-homo-def
proof (transfer)
  fix  $z1 z2 :: \text{complex}$ 
  assume  $z2 \neq 0$ 
  thus  $\text{of-complex-coords } z1 *_{hc} \text{reciprocal-homo-coords } (\text{of-complex-coords } z2) \approx$ 
     $\text{of-complex-coords } (z1 / z2)$ 
  by (auto simp add: of-complex-coords-def Abs-homo-coords-inverse mult-homo-coords-def
  reciprocal-homo-coords-def)
    (rule-tac  $x=1/z2$  in exI, auto)
qed

```

```

lemma divide-homo-coords-Rep [simp]:
  assumes Rep-homo-coords  $z = (z1, z2)$  Rep-homo-coords  $w = (w1, w2)$ 
     $(z2 \neq 0 \vee w2 \neq 0) \wedge (w1 \neq 0 \vee z1 \neq 0)$ 
  shows Rep-homo-coords  $(z *_{hc} (\text{reciprocal-homo-coords } w)) = (z1*w2, z2*w1)$ 
using assms
using mult-homo-coords-Rep[of  $z z1 z2$  reciprocal-homo-coords  $w w2 w1$ ] reciprocal-homo-coords-Rep[of
 $w$ ]
by simp

```

Conjugate

```

definition cnj-homo-coords where
  cnj-homo-coords  $z = (\text{let } (z1, z2) = \text{Rep-homo-coords } z \text{ in } \text{Abs-homo-coords } (\text{cnj } z1, \text{cnj } z2))$ 

lemma [simp]: Rep-homo-coords (cnj-homo-coords  $z$ ) = vec-cnj (Rep-homo-coords  $z$ )
apply (cases Rep-homo-coords  $z$ )
using Rep-homo-coords[of  $z$ ]
by (simp add: cnj-homo-coords-def Abs-homo-coords-inverse vec-cnj-def)

```

```

lift-definition cnj-homo :: complex-homo  $\Rightarrow$  complex-homo is cnj-homo-coords
by auto

```

```

lemma cnj-homo (of-complex z) = of-complex (cnj z)
by (transfer) (simp add: vec-cnj-def)

lemma cnj-homo  $\infty_h = \infty_h$ 
by (transfer) (simp add: vec-cnj-def)

lemma cnj-homo-coords-involution [simp]:
  cnj-homo-coords (cnj-homo-coords z) = z
unfolding cnj-homo-coords-def[of cnj-homo-coords z] Let-def
by (cases Rep-homo-coords z, auto simp add: Let-def split-def vec-cnj-def) (metis
Rep-homo-coords-inverse)

lemma cnj-homo-involution [simp]: cnj-homo (cnj-homo z) = z
by (transfer) (auto, rule-tac x=1 in exI, simp)

lemma [simp]:
  cnj-homo  $\infty_h = \infty_h$ 
by (transfer) (auto simp add: vec-cnj-def)

lemma [simp]:
  cnj-homo  $\theta_h = \theta_h$ 
by (transfer) (auto simp add: vec-cnj-def)

Inversion

definition inversion-homo where
  inversion-homo = cnj-homo  $\circ$  reciprocal-homo

lemma inversion-homo-sym:
  inversion-homo = reciprocal-homo  $\circ$  cnj-homo
unfolding inversion-homo-def
by (rule ext, simp) (transfer, case-tac Rep-homo-coords x, auto simp add: reciprocal-homo-coords-Rep
split-def Let-def vec-cnj-def, metis zero-neq-one)

lemma inversion-homo-involution [simp]: inversion-homo (inversion-homo z) = z
proof-
  have *: cnj-homo  $\circ$  reciprocal-homo = reciprocal-homo  $\circ$  cnj-homo
  using inversion-homo-sym
  by (simp add: inversion-homo-def)
  show ?thesis
  unfolding inversion-homo-def
  by (subst *) simp
qed

lemma [simp]:
  inversion-homo  $\theta_h = \infty_h$ 
by (simp add: inversion-homo-def)

lemma [simp]:
  inversion-homo  $\infty_h = \theta_h$ 

```

by (simp add: inversion-homo-def)

8.1 Ratio and crossratio

definition ratio-rep **where**

```
ratio-rep z1 z2 z3 =
  (let (z1x, z1y) = Rep-homo-coords z1;
   (z2x, z2y) = Rep-homo-coords z2;
   (z3x, z3y) = Rep-homo-coords z3 in
  Abs-homo-coords ((z1x*z2y - z2x*z1y)*z3y, (z1x*z3y - z3x*z1y)*z2y))
```

lemma ratio-rep-Rep [simp]:

```
assumes ( $\neg z1 \approx z2 \wedge \neg z3 \approx \text{inf-homo-rep}$ )  $\vee (\neg z1 \approx z3 \wedge \neg z2 \approx \text{inf-homo-rep})$ 
shows Rep-homo-coords (ratio-rep z1 z2 z3) = (let (z1x, z1y) = Rep-homo-coords z1;
                                              (z2x, z2y) = Rep-homo-coords z2;
                                              (z3x, z3y) = Rep-homo-coords z3 in
                                              ((z1x*z2y - z2x*z1y)*z3y, (z1x*z3y - z3x*z1y)*z2y))
```

proof –

```
obtain z1' z1'' where zz1: Rep-homo-coords z1 = (z1', z1'')
  by (rule obtain-homo-coords)
obtain z2' z2'' where zz2: Rep-homo-coords z2 = (z2', z2'')
  by (rule obtain-homo-coords)
obtain z3' z3'' where zz3: Rep-homo-coords z3 = (z3', z3'')
  by (rule obtain-homo-coords)
have ((z1' * z2'' - z2' * z1'') * z3'', (z1' * z3'' - z3' * z1'') * z2'')  $\neq$  vec-zero
  using assms
  using homo-coords-eq-mix[OF zz1 zz2] homo-coords-eq-mix[OF zz3, of inf-homo-rep
1 0]
  using homo-coords-eq-mix[OF zz1 zz3] homo-coords-eq-mix[OF zz2, of inf-homo-rep
1 0]
  by auto
thus ?thesis
  using zz1 zz2 zz3
  unfolding ratio-rep-def Let-def
  by (simp add: Abs-homo-coords-inverse)
qed
```

lemma ratio-rep-Rep' [simp]:

```
assumes (z1  $\approx$  z2  $\vee$  z3  $\approx$  inf-homo-rep)  $\wedge$  (z1  $\approx$  z3  $\vee$  z2  $\approx$  inf-homo-rep)
shows ratio-rep z1 z2 z3 = Abs-homo-coords (0, 0)
```

using assms

unfolding ratio-rep-def

```
by (cases Rep-homo-coords z1, cases Rep-homo-coords z2, cases Rep-homo-coords z3) auto
```

lift-definition ratio :: complex-homo \Rightarrow complex-homo \Rightarrow complex-homo \Rightarrow complex-homo
is ratio-rep
proof –

```

fix z1 z2 z3 w1 w2 w3
assume *: z1 ≈ w1 z2 ≈ w2 z3 ≈ w3
obtain z1' z1'' where zz1: Rep-homo-coords z1 = (z1', z1'')
  by (rule obtain-homo-coords)
obtain z2' z2'' where zz2: Rep-homo-coords z2 = (z2', z2'')
  by (rule obtain-homo-coords)
obtain z3' z3'' where zz3: Rep-homo-coords z3 = (z3', z3'')
  by (rule obtain-homo-coords)
obtain w1' w1'' where ww1: Rep-homo-coords w1 = (w1', w1'')
  by (rule obtain-homo-coords)
obtain w2' w2'' where ww2: Rep-homo-coords w2 = (w2', w2'')
  by (rule obtain-homo-coords)
obtain w3' w3'' where ww3: Rep-homo-coords w3 = (w3', w3'')
  by (rule obtain-homo-coords)

show ratio-rep z1 z2 z3 ≈ ratio-rep w1 w2 w3
proof (cases ¬ z1 ≈ z2 ∧ ¬ z3 ≈ inf-homo-rep ∨ ¬ z1 ≈ z3 ∧ ¬ z2 ≈
inf-homo-rep)
  case True
  hence ¬ w1 ≈ w2 ∧ ¬ w3 ≈ inf-homo-rep ∨ ¬ w1 ≈ w3 ∧ ¬ w2 ≈ inf-homo-rep
    using * homo-coords-eq-sym homo-coords-eq-trans
    by metis
  thus ?thesis
    apply (subst homo-coords-eq-def, unfold Let-def)
    using ratio-rep-Rep[OF ← z1 ≈ z2 ∧ ¬ z3 ≈ inf-homo-rep ∨ ¬ z1 ≈ z3 ∧
¬ z2 ≈ inf-homo-rep]
    using ratio-rep-Rep[OF ← w1 ≈ w2 ∧ ¬ w3 ≈ inf-homo-rep ∨ ¬ w1 ≈ w3 ∧
¬ w2 ≈ inf-homo-rep]
    using zz1 zz2 zz3 ww1 ww2 ww3 *
    by (simp add: Let-def field-simps, (erule-tac exE)+) (rule-tac x=k*ka*kb in
exI, simp)
  next
  case False
  hence ¬ (¬ w1 ≈ w2 ∧ ¬ w3 ≈ inf-homo-rep ∨ ¬ w1 ≈ w3 ∧ ¬ w2 ≈
inf-homo-rep)
    using * homo-coords-eq-sym homo-coords-eq-trans
    by metis
  thus ?thesis
    using False
    by (simp del: homo-coords-eq-def)
qed
qed

lemma ratio-is-ratio:
assumes z1 ≠ z2 ∨ z1 ≠ z3 z1 ≠ ∞h z2 ≠ ∞h ∨ z3 ≠ ∞h
shows ratio z1 z2 z3 = (z1 -h z2) :h (z1 -h z3)
unfolding minus-homo-def divide-homo-def
using assms
proof transfer

```

```

fix z w v
obtain z1 z2 where zz: Rep-homo-coords z = (z1, z2)
  by (rule obtain-homo-coords)
obtain w1 w2 where ww: Rep-homo-coords w = (w1, w2)
  by (rule obtain-homo-coords)
obtain v1 v2 where vv: Rep-homo-coords v = (v1, v2)
  by (rule obtain-homo-coords)
assume *: ¬ z ≈ w ∨ ¬ z ≈ v ∨ ¬ w ≈ inf-homo-rep
  ¬ w ≈ inf-homo-rep ∨ ¬ v ≈ inf-homo-rep
hence **: ¬ z ≈ w ∧ ¬ v ≈ inf-homo-rep ∨ ¬ z ≈ v ∧ ¬ w ≈ inf-homo-rep
  by (metis homo-coords-eq-trans)
have z2 ≠ 0 w2 ≠ 0 ∨ v2 ≠ 0 z1*w2 ≠ z2*w1 ∨ z1*v2 ≠ z2*v1
  using zz vv ww not-inf-snd-not0[of v] not-inf-snd-not0[of z] not-inf-snd-not0[w]
    homo-coords-eq-mix[of z z1 z2 w w1 w2] homo-coords-eq-mix[of z z1 z2 v v1 v2]
*
  by auto
thus ratio-rep z w v ≈
  z +hc uminus-homo-coords w *hc
  reciprocal-homo-coords (z +hc uminus-homo-coords v)
using zz ww vv **
  using divide-homo-coords-Rep[of z +hc uminus-homo-coords w z1 * w2 + - w1 * z2 z2 * w2 (z +hc uminus-homo-coords v) z1 * v2 + - v1 * z2 z2 * v2 ]
  using minus-homo-coords-Rep[of z z1 z2 w w1 w2]
  using minus-homo-coords-Rep[of z z1 z2 v v1 v2]
  by (auto simp add: field-simps)
qed

```

```

lemma
assumes z2 ≠ ∞h z3 ≠ ∞h
shows ratio ∞h z2 z3 = 1h
using assms
proof transfer
fix z2 z3
obtain z2x z2y where zz2: Rep-homo-coords z2 = (z2x, z2y)
  by (rule obtain-homo-coords)
obtain z3x z3y where zz3: Rep-homo-coords z3 = (z3x, z3y)
  by (rule obtain-homo-coords)
assume ¬ z2 ≈ inf-homo-rep ¬ z3 ≈ inf-homo-rep
have z2y ≠ 0 z3y ≠ 0
  using not-inf-snd-not0[OF (¬ z2 ≈ inf-homo-rep)] zz2
  using not-inf-snd-not0[OF (¬ z3 ≈ inf-homo-rep)] zz3
  by auto
thus ratio-rep inf-homo-rep z2 z3 ≈ one-homo-rep
  using (¬ z2 ≈ inf-homo-rep) (¬ z3 ≈ inf-homo-rep) zz2 zz3
  by (subst homo-coords-eq-def, subst ratio-rep-Rep, simp-all) (rule-tac x=1/(z2y*z3y))
in exI, auto
qed

```

lemma

assumes $z1 \neq \infty_h z3 \neq \infty_h$
 shows ratio $z1 \infty_h z3 = \infty_h$
 using assms
proof transfer
 fix $z1 z3$
obtain $z1x z1y$ where $zz1: Rep\text{-homo-coords } z1 = (z1x, z1y)$
 by (rule obtain-homo-coords)
obtain $z3x z3y$ where $zz3: Rep\text{-homo-coords } z3 = (z3x, z3y)$
 by (rule obtain-homo-coords)
assume $\neg z1 \approx inf\text{-homo-rep} \neg z3 \approx inf\text{-homo-rep}$
have $z1y \neq 0 z3y \neq 0$
 using not-inf-snd-not0[$OF \neg z1 \approx inf\text{-homo-rep}] zz1$
 using not-inf-snd-not0[$OF \neg z3 \approx inf\text{-homo-rep}] zz3$
 by auto
 thus ratio-rep $z1 inf\text{-homo-rep} z3 \approx inf\text{-homo-rep}$
 using $\neg z1 \approx inf\text{-homo-rep} \neg z3 \approx inf\text{-homo-rep} zz1 zz3$
 by (subst homo-coords-eq-def, subst ratio-rep-Rep, simp-all) (rule-tac $x=-1/(z1y*z3y)$)
 in exI, auto
qed

lemma

assumes $z1 \neq \infty_h z2 \neq \infty_h$
 shows ratio $z1 z2 \infty_h = 0_h$
 using assms
proof transfer
 fix $z1 z2$
obtain $z1x z1y$ where $zz1: Rep\text{-homo-coords } z1 = (z1x, z1y)$
 by (rule obtain-homo-coords)
obtain $z2x z2y$ where $zz2: Rep\text{-homo-coords } z2 = (z2x, z2y)$
 by (rule obtain-homo-coords)
assume $\neg z1 \approx inf\text{-homo-rep} \neg z2 \approx inf\text{-homo-rep}$
have $z1y \neq 0 z2y \neq 0$
 using not-inf-snd-not0[$OF \neg z1 \approx inf\text{-homo-rep}] zz1$
 using not-inf-snd-not0[$OF \neg z2 \approx inf\text{-homo-rep}] zz2$
 by auto
 thus ratio-rep $z1 z2 inf\text{-homo-rep} \approx zero\text{-homo-rep}$
 using $\neg z1 \approx inf\text{-homo-rep} \neg z2 \approx inf\text{-homo-rep} zz1 zz2$
 by (subst homo-coords-eq-def, subst ratio-rep-Rep, simp-all) (rule-tac $x=-1/(z1y*z2y)$)
 in exI, auto
qed

lemma

assumes $z1 \neq z2 z1 \neq \infty_h$
 shows ratio $z1 z2 z1 = \infty_h$
proof–
have $z1 -_h z2 \neq 0_h$

```

using diff-zero-homo[of z1 z2]  $\langle z1 \neq z2 \rangle \langle z1 \neq \infty_h \rangle$ 
by auto
thus ?thesis
  using assms
  using ratio-is-ratio[of z1 z2 z1]
  by simp
qed

```

```

definition cross-ratio-rep where
cross-ratio-rep z u v w =
  (let (z', z'') = Rep-homo-coords z;
   (u', u'') = Rep-homo-coords u;
   (v', v'') = Rep-homo-coords v;
   (w', w'') = Rep-homo-coords w
   in Abs-homo-coords ((z'*u'' - u'*z'')*(v'*w'' - w'*v''),
                        (z'*w'' - w'*z'')*(v'*u'' - u'*v'')))

lemma cross-ratio-rep-Rep [simp]:
assumes ( $\neg z1 \approx z2 \wedge \neg z3 \approx z4$ )  $\vee$  ( $\neg z1 \approx z4 \wedge \neg z2 \approx z3$ )
shows Rep-homo-coords (cross-ratio-rep z1 z2 z3 z4) =
  (let (z1', z1'') = Rep-homo-coords z1;
   (z2', z2'') = Rep-homo-coords z2;
   (z3', z3'') = Rep-homo-coords z3;
   (z4', z4'') = Rep-homo-coords z4
   in ((z1'*z2'' - z2'*z1'')*(z3'*z4'' - z4'*z3''), (z1'*z4'' - z4'*z1'')*(z3'*z2'' - z2'*z3'')))

proof-
  obtain z1' z1'' where zz1: Rep-homo-coords z1 = (z1', z1'')
  by (rule obtain-homo-coords)
  obtain z2' z2'' where zz2: Rep-homo-coords z2 = (z2', z2'')
  by (rule obtain-homo-coords)
  obtain z3' z3'' where zz3: Rep-homo-coords z3 = (z3', z3'')
  by (rule obtain-homo-coords)
  obtain z4' z4'' where zz4: Rep-homo-coords z4 = (z4', z4'')
  by (rule obtain-homo-coords)
  show ?thesis
    using zz1 zz2 zz3 zz4
    using assms
    unfolding cross-ratio-rep-def Let-def
    using homo-coords-eq-mix[OF zz1 zz2] homo-coords-eq-mix[OF zz3 zz4]
    using homo-coords-eq-mix[OF zz1 zz4] homo-coords-eq-mix[OF zz2 zz3]
    by (auto simp add: Abs-homo-coords-inverse)
qed

lift-definition cross-ratio :: complex-homo  $\Rightarrow$  complex-homo  $\Rightarrow$  complex-homo  $\Rightarrow$ 
complex-homo  $\Rightarrow$  complex-homo is cross-ratio-rep
proof-

```

```

fix z1 z2 z3 z4 w1 w2 w3 w4
obtain z1' z1'' where zz1: Rep-homo-coords z1 = (z1', z1'')
  by (rule obtain-homo-coords)
obtain z2' z2'' where zz2: Rep-homo-coords z2 = (z2', z2'')
  by (rule obtain-homo-coords)
obtain z3' z3'' where zz3: Rep-homo-coords z3 = (z3', z3'')
  by (rule obtain-homo-coords)
obtain z4' z4'' where zz4: Rep-homo-coords z4 = (z4', z4'')
  by (rule obtain-homo-coords)
obtain w1' w1'' where ww1: Rep-homo-coords w1 = (w1', w1'')
  by (rule obtain-homo-coords)
obtain w2' w2'' where ww2: Rep-homo-coords w2 = (w2', w2'')
  by (rule obtain-homo-coords)
obtain w3' w3'' where ww3: Rep-homo-coords w3 = (w3', w3'')
  by (rule obtain-homo-coords)
obtain w4' w4'' where ww4: Rep-homo-coords w4 = (w4', w4'')
  by (rule obtain-homo-coords)
let ?w12 = w1' * w2'' - w2' * w1''
let ?w34 = w3' * w4'' - w4' * w3''
let ?w14 = w1' * w4'' - w4' * w1''
let ?w32 = w3' * w2'' - w2' * w3''
let ?z12 = z1' * z2'' - z2' * z1''
let ?z34 = z3' * z4'' - z4' * z3''
let ?z14 = z1' * z4'' - z4' * z1''
let ?z32 = z3' * z2'' - z2' * z3''

assume *: z1 ≈ w1 z2 ≈ w2 z3 ≈ w3 z4 ≈ w4
hence **:
  ?w12 * ?w34 = 0 ↔ ?z12 * ?z34 = 0 ?w14 * ?w32 = 0 ↔ ?z14 * ?z32
= 0
  using zz1 zz2 zz3 zz4 ww1 ww2 ww3 ww4
  by auto

show cross-ratio-rep z1 z2 z3 z4 ≈ cross-ratio-rep w1 w2 w3 w4
proof (cases ?z12 * ?z34 = 0 ∧ ?z14 * ?z32 = 0)
  case True
  thus ?thesis
    using zz1 zz2 zz3 zz4 ww1 ww2 ww3 ww4 @@
    by (simp add: cross-ratio-rep-def split-def Let-def) (rule-tac x=1 in exI, auto)
  next
    case False
    have ¬ z1 ≈ z2 ∧ ¬ z3 ≈ z4 ∨ ¬ z1 ≈ z4 ∧ ¬ z2 ≈ z3
    using False
    using homo-coords-eq-mix[OF zz1 zz2] homo-coords-eq-mix[OF zz3 zz4]
    using homo-coords-eq-mix[OF zz1 zz4] homo-coords-eq-mix[OF zz2 zz3]
    by (simp del: homo-coords-eq-def) metis
  moreover
  have ¬ w1 ≈ w2 ∧ ¬ w3 ≈ w4 ∨ ¬ w1 ≈ w4 ∧ ¬ w2 ≈ w3
  using @@ False

```

```

using homo-coords-eq-mix[OF ww1 ww2] homo-coords-eq-mix[OF ww3 ww4]
using homo-coords-eq-mix[OF ww1 ww4] homo-coords-eq-mix[OF ww2 ww3]
by (simp del: homo-coords-eq-def) metis
ultimately
show ?thesis
  using *
  using cross-ratio-rep-Rep[of z1 z2 z3 z4]
  using cross-ratio-rep-Rep[of w1 w2 w3 w4]
  using zz1 zz2 zz3 zz4 ww1 ww2 ww3 ww4
  apply simp
  apply (erule exE)+
  apply simp
  apply (rule-tac x=k*ka*kb*kc in exI)
  apply (simp add: field-simps)
  done
qed
qed

lemma cross-ratio z 0h 1h ∞h = z
proof (transfer)
fix z
have *: ¬ z ≈ zero-homo-rep ∧ ¬ one-homo-rep ≈ inf-homo-rep ∨ ¬ z ≈
inf-homo-rep ∧ ¬ zero-homo-rep ≈ one-homo-rep
by (cases Rep-homo-coords z) auto
show cross-ratio-rep z zero-homo-rep one-homo-rep inf-homo-rep ≈ z
using cross-ratio-rep-Rep[OF *]
by (simp add: split-def Let-def) (rule-tac x=-1 in exI, simp)
qed

lemma cross-ratio-0:
assumes z1 ≠ z2 z1 ≠ z3
shows cross-ratio z1 z1 z2 z3 = 0h
using assms
proof (transfer)
fix z1 z2 z3
let ?z1 = Rep-homo-coords z1 and ?z2 = Rep-homo-coords z2 and ?z3 =
Rep-homo-coords z3
assume ¬ z1 ≈ z2 ∨ z1 ≈ z3
thus cross-ratio-rep z1 z1 z2 z3 ≈ zero-homo-rep
using cross-ratio-rep-Rep[of z1 z1 z2 z3]
homo-coords-eq-mix[of z1 fst ?z1 snd ?z1 z2 fst ?z2 snd ?z2] homo-coords-eq-mix[of
z1 fst ?z1 snd ?z1 z3 fst ?z3 snd ?z3]
by (cases ?z1, cases ?z2, cases ?z3, simp add: split-def Let-def)
qed

lemma cross-ratio-1:
assumes z1 ≠ z2 z2 ≠ z3
shows cross-ratio z2 z1 z2 z3 = 1h
using assms

```

```

proof (transfer)
  fix  $z_1 z_2 z_3$ 
  obtain  $z_1' z_1''$  where  $zz1: Rep\text{-homo-coords } z_1 = (z_1', z_1'')$ 
    by (rule obtain-homo-coords)
  obtain  $z_2' z_2''$  where  $zz2: Rep\text{-homo-coords } z_2 = (z_2', z_2'')$ 
    by (rule obtain-homo-coords)
  obtain  $z_3' z_3''$  where  $zz3: Rep\text{-homo-coords } z_3 = (z_3', z_3'')$ 
    by (rule obtain-homo-coords)
  assume  $\neg z_1 \approx z_2 \neg z_2 \approx z_3$ 
  thus cross-ratio-rep  $z_2 z_1 z_2 z_3 \approx$  one-homo-rep
    using  $zz1 zz2 zz3$ 
    using homo-coords-eq-mix[of  $z_1 z_1' z_1'' z_2 z_2' z_2''$ ] homo-coords-eq-mix[of  $z_2 z_2' z_2'' z_3 z_3' z_3''$ ]
      by (auto simp add: cross-ratio-rep-def split-def Let-def Abs-homo-coords-inverse)
      (rule-tac  $x=1 / ((z_2'*z_3'' - z_3'*z_2'') * (z_2'*z_1'' - z_1'*z_2''))$  in exI, simp)
  qed

lemma cross-ratio-inf:
  assumes  $z_1 \neq z_3 z_2 \neq z_3$ 
  shows cross-ratio  $z_3 z_1 z_2 z_3 = \infty_h$ 
  using assms
  proof (transfer)
    fix  $z_1 z_2 z_3$ 
    obtain  $z_1' z_1''$  where  $zz1: Rep\text{-homo-coords } z_1 = (z_1', z_1'')$ 
      by (rule obtain-homo-coords)
    obtain  $z_2' z_2''$  where  $zz2: Rep\text{-homo-coords } z_2 = (z_2', z_2'')$ 
      by (rule obtain-homo-coords)
    obtain  $z_3' z_3''$  where  $zz3: Rep\text{-homo-coords } z_3 = (z_3', z_3'')$ 
      by (rule obtain-homo-coords)
    assume  $\neg z_1 \approx z_3 \neg z_2 \approx z_3$ 
    thus cross-ratio-rep  $z_3 z_1 z_2 z_3 \approx$  inf-homo-rep
      using  $zz1 zz2 zz3$ 
      using homo-coords-eq-mix[of  $z_1 z_1' z_1'' z_3 z_3' z_3''$ ] homo-coords-eq-mix[of  $z_2 z_2' z_2'' z_3 z_3' z_3''$ ]
        by (auto simp add: cross-ratio-rep-def split-def Let-def Abs-homo-coords-inverse)
    qed

lemma
  assumes  $(z \neq u \wedge v \neq w) \vee (z \neq w \wedge u \neq v) z \neq \infty_h u \neq \infty_h v \neq \infty_h w \neq \infty_h$ 
  shows cross-ratio  $z u v w = ((z -_h u) *_h (v -_h w)) :_h ((z -_h w) *_h (v -_h u))$ 
  using assms
  unfolding minus-homo-def divide-homo-def
  proof transfer
    fix  $z u v w$ 
    obtain  $z_1 z_2$  where  $zz: Rep\text{-homo-coords } z = (z_1, z_2)$ 
      by (rule obtain-homo-coords)
    obtain  $u_1 u_2$  where  $uu: Rep\text{-homo-coords } u = (u_1, u_2)$ 
      by (rule obtain-homo-coords)

```

```

obtain v1 v2 where vv: Rep-homo-coords v = (v1, v2)
  by (rule obtain-homo-coords)
obtain w1 w2 where ww: Rep-homo-coords w = (w1, w2)
  by (rule obtain-homo-coords)

assume *:  $\neg z \approx u \wedge \neg v \approx w \vee \neg z \approx w \wedge \neg u \approx v$  and
  **:  $\neg z \approx \text{inf-homo-rep} \wedge \neg u \approx \text{inf-homo-rep} \wedge \neg v \approx \text{inf-homo-rep} \wedge \neg w \approx \text{inf-homo-rep}$ 
have z2 ≠ 0 u2 ≠ 0 v2 ≠ 0 w2 ≠ 0
  using ** zz uu vv ww
  using not-inf-snd-not0[of z] not-inf-snd-not0[of u] not-inf-snd-not0[of v] not-inf-snd-not0[of w]
  by simp-all
moreover
  have ((z1*u2 - z2*u1 ≠ 0)  $\wedge$  (v1*w2 - v2*w1 ≠ 0))  $\vee$  ((z1*w2 - z2*w1 ≠ 0)  $\wedge$  (v1*u2 - v2*u1 ≠ 0))
  using *
  apply (subst (asm) homo-coords-eq-mix[OF zz uu])
  apply (subst (asm) homo-coords-eq-mix[OF vv ww])
  apply (subst (asm) homo-coords-eq-mix[OF zz ww])
  apply (subst (asm) homo-coords-eq-mix[OF uu vv])
  by (auto simp add: field-simps)
moreover
  hence z1 * w2 ≠ w1 * z2  $\wedge$  v1 * u2 ≠ u1 * v2  $\vee$  z1 * u2 ≠ u1 * z2  $\wedge$  v1 * w2 ≠ w1 * v2
  by auto
ultimately
  show cross-ratio-rep z u v w ≈
    z +hc uminus-homo-coords u *hc (v +hc uminus-homo-coords w) *hc
    reciprocal-homo-coords
    (z +hc uminus-homo-coords w *hc (v +hc uminus-homo-coords u))
  using uu vv ww zz *
  apply simp
  apply (subst divide-homo-coords-Rep[of (z +hc uminus-homo-coords u) *hc (v +hc uminus-homo-coords w)] (z1 * u2 - u1 * z2) * (v1 * w2 - w1 * v2) z2 * u2 * (v2 * w2) (z +hc uminus-homo-coords w) *hc (v +hc uminus-homo-coords u) (z1 * w2 - w1 * z2) * (v1 * u2 - u1 * v2) z2 * w2 * (v2 * u2))
  using mult-homo-coords-Rep[of z +hc uminus-homo-coords u z1 * u2 - u1 * z2 z2 * u2 v +hc uminus-homo-coords w v1 * w2 - w1 * v2 v2 * w2]
  using minus-homo-coords-Rep[of z z1 z2 u u1 u2]
  using minus-homo-coords-Rep[of v v1 v2 w w1 w2]
  using mult-homo-coords-Rep[of z +hc uminus-homo-coords w z1 * w2 - w1 * z2 z2 * w2 v +hc uminus-homo-coords u v1 * u2 - u1 * v2 v2 * u2]
  using minus-homo-coords-Rep[of z z1 z2 w w1 w2]
  using minus-homo-coords-Rep[of v v1 v2 u u1 u2]
  using mult-homo-coords-Rep[of z +hc uminus-homo-coords u z1 * u2 - u1 * z2 z2 * u2 v +hc uminus-homo-coords w v1 * w2 - w1 * v2 v2 * w2]
  using minus-homo-coords-Rep[of z z1 z2 u u1 u2]

```

```

using minus-homo-coords-Rep[of v v1 v2 w w1 w2]
by simp-all (rule-tac  $x=z2*u2*(v2*w2)$  in exI, simp)
qed

```

8.2 Distance

definition inprod-homo-rep **where**

```

inprod-homo-rep z w =
  (let (z1, z2) = Rep-homo-coords z;
   (w1, w2) = Rep-homo-coords w
   in vec-cnj (z1, z2) *vv (w1, w2))

```

syntax

```
-inprod-homo-rep :: homo-coords  $\Rightarrow$  homo-coords  $\Rightarrow$  complex ( $\langle \cdot, \cdot \rangle$ )

```

translations

```
 $\langle z, w \rangle == CONST$  inprod-homo-rep z w
```

lemma [simp]: *is-real* $\langle z, z \rangle$

unfolding inprod-homo-rep-def

by (cases Rep-homo-coords *z*, simp add: vec-cnj-def)

lemma [simp]: $Re \langle z, z \rangle \geq 0$

unfolding inprod-homo-rep-def

by (cases Rep-homo-coords *z*, simp add: vec-cnj-def)

lemma inprod-homo-bilinear1:

assumes Rep-homo-coords *z' = k *_{sv} Rep-homo-coords z*

shows $\langle z', w \rangle = cnj k * \langle z, w \rangle$

using assms

unfolding inprod-homo-rep-def Let-def

by (cases Rep-homo-coords *z*, cases Rep-homo-coords *z'*, cases Rep-homo-coords *w*)
 (simp add: vec-cnj-def complex-cnj field-simps)

lemma inprod-homo-bilinear2:

assumes Rep-homo-coords *w' = k *_{sv} Rep-homo-coords w*

shows $\langle z, w' \rangle = k * \langle z, w \rangle$

using assms

unfolding inprod-homo-rep-def Let-def

by (cases Rep-homo-coords *z*, cases Rep-homo-coords *z'*, cases Rep-homo-coords *w*)
 (simp add: vec-cnj-def complex-cnj field-simps)

definition norm-homo-rep **where**

norm-homo-rep *z* = sqrt ($Re \langle z, z \rangle$)

syntax

```
-norm-homo-rep :: homo-coords  $\Rightarrow$  complex ( $\langle \cdot \rangle$ )

```

translations

```
 $\langle z \rangle == CONST$  norm-homo-rep z
```

lemma

norm-homo-rep-square: $\langle z \rangle^2 = Re (\langle z, z \rangle)$

```

unfolding norm-homo-rep-def
by simp

lemma norm-homo-gt-0:  $\langle z \rangle > 0$ 
proof-
obtain z1 z2 where Rep-homo-coords z = (z1, z2)
by (rule obtain-homo-coords)
thus ?thesis
using complex-mult-cnj-cmod[of z1] complex-mult-cnj-cmod[of z2] Rep-homo-coords[of z]
unfolding norm-homo-rep-def inprod-homo-rep-def
by (simp add: vec-cnj-def split-def Let-def field-simps power2-eq-square) (metis
norm-eq-zero sum-squares-gt-zero-iff)
qed

lemma norm-homo-scale:
assumes Rep-homo-coords z' = k *sv Rep-homo-coords z
shows  $\langle z' \rangle^2 = \operatorname{Re}(\operatorname{cnj}(k * k) * \langle z \rangle^2)$ 
apply (subst norm-homo-rep-square)+
apply (subst inprod-homo-bilinear1[OF assms])
apply (subst inprod-homo-bilinear2[OF assms])
apply (simp add: field-simps)
done

definition dist-homo-rep where
dist-homo-rep z1 z2 =
(let (z1x, z1y) = Rep-homo-coords z1;
 (z2x, z2y) = Rep-homo-coords z2;
 num = (z1x*z2y - z2x*z1y) * (cnj z1x*cnj z2y - cnj z2x*cnj z1y);
 den = (z1x*cnj z1x + z1y*cnj z1y) * (z2x*cnj z2x + z2y*cnj z2y)
in 2*sqrt(Re num / Re den))

lemma dist-homo-rep-iff: dist-homo-rep z w = 2*sqrt(1 - (cmod(z,w))^2 / ( $\langle z \rangle^2$ 
*  $\langle w \rangle^2$ ))
proof-
obtain z1 z2 w1 w2 where *: Rep-homo-coords z = (z1, z2) Rep-homo-coords
w = (w1, w2)
by (cases Rep-homo-coords z, cases Rep-homo-coords w) auto
have 1:  $2\sqrt{1 - (\operatorname{cmod}(z,w))^2 / (\langle z \rangle^2 * \langle w \rangle^2)} = \sqrt{(\langle z \rangle^2 * \langle w \rangle^2 - (\operatorname{cmod}(z,w))^2) / (\langle z \rangle^2 * \langle w \rangle^2)}$ 
using norm-homo-gt-0[of z] norm-homo-gt-0[of w]
by (simp add: field-simps)

have 2:  $\langle z \rangle^2 * \langle w \rangle^2 = \operatorname{Re}((z1*cnj z1 + z2*cnj z2) * (w1*cnj w1 + w2*cnj w2))$ 
using *
by (simp add: norm-homo-rep-def inprod-homo-rep-def vec-cnj-def)

have 3:  $\langle z \rangle^2 * \langle w \rangle^2 - (\operatorname{cmod}(z,w))^2 = \operatorname{Re}((z1*w2 - w1*z2) * (cnj z1*cnj w2$ 

```

```

- cnj w1*cnj z2))
  apply (subst cmod-square, (subst norm-homo-rep-square)+)
  using *
  by (simp add: inprod-homo-rep-def vec-cnj-def field-simps)

thus ?thesis
  using 1 2 3
  using *
  unfolding dist-homo-rep-def Let-def
  by simp
qed

lift-definition dist-homo :: complex-homo ⇒ complex-homo ⇒ real is dist-homo-rep
proof-
  fix z1 z2 z1' z2'
  obtain z1x z1y z2x z2y z1'x z1'y z2'x z2'y where
    zz: Rep-homo-coords z1 = (z1x, z1y) Rep-homo-coords z2 = (z2x, z2y) Rep-homo-coords
    z1' = (z1'x, z1'y) Rep-homo-coords z2' = (z2'x, z2'y)
    by (cases Rep-homo-coords z1, cases Rep-homo-coords z2, cases Rep-homo-coords
    z1', cases Rep-homo-coords z2') blast

  assume z1 ≈ z1' z2 ≈ z2'
  then obtain k1 k2 where
    *: k1 ≠ 0 Rep-homo-coords z1' = k1 *sv Rep-homo-coords z1 and
    **: k2 ≠ 0 Rep-homo-coords z2' = k2 *sv Rep-homo-coords z2
    by auto
  have (cmod ⟨z1,z2⟩)2 / ((⟨z1⟩2 * ⟨z2⟩2) = (cmod ⟨z1',z2'⟩)2 / ((⟨z1'⟩2 * ⟨z2'⟩2)
  using ⟨k1 ≠ 0⟩ ⟨k2 ≠ 0⟩
  using cmod-square[symmetric, of k1] cmod-square[symmetric, of k2]
  apply (subst norm-homo-scale[OF *(2)])
  apply (subst norm-homo-scale[OF **(2)])
  apply (subst inprod-homo-bilinear1[OF *(2)])
  apply (subst inprod-homo-bilinear2[OF **(2)])
  by (simp add: power2-eq-square)
  thus dist-homo-rep z1 z2 = dist-homo-rep z1' z2'
    by (subst dist-homo-rep-iff)+
  qed

lemma dist-homo-finite:
  dist-homo (of-complex z1) (of-complex z2) = 2 * cmod(z1 - z2) / (sqrt (1 + (cmod
  z1)2) * sqrt (1 + (cmod z2)2))
  apply transfer
  apply (subst cmod-square)+
  apply (simp add: dist-homo-rep-def real-sqrt-divide cmod-def power2-eq-square)
  by (smt ab-diff-minus comm-semiring-1-class.normalize-semiring-rules(24) minus-diff-eq
  minus-mult-right real-sqrt-mult-distrib2)

lemma dist-homo-infinite1:
  dist-homo (of-complex z1) ∞h = 2 / sqrt (1 + (cmod z1)2)

```

```

by transfer (subst cmod-square, simp add: dist-homo-rep-def real-sqrt-divide)

lemma dist-homo-infinite2:
  dist-homo ∞h (of-complex z1) = 2 / sqrt (1+(cmod z1)2)
by transfer (subst cmod-square, simp add: dist-homo-rep-def real-sqrt-divide)

lemma dist-homo-rep-zero:
  dist-homo-rep z w = 0 ↔ (cmod ⟨z,w⟩)2 = (⟨z⟩2 * ⟨w⟩2)
using norm-homo-gt-0[of z] norm-homo-gt-0[of w]
by (subst dist-homo-rep-iff) auto

lemma dist-homo-zero1 [simp]: dist-homo z z = 0
by transfer (subst dist-homo-rep-zero, ((subst norm-homo-rep-square)+), subst cmod-square,
simp)

lemma dist-homo-zero2 [simp]:
assumes dist-homo z1 z2 = 0
shows z1 = z2
using assms
proof transfer
fix z w
obtain z1 z2 w1 w2 where *: Rep-homo-coords z = (z1, z2) Rep-homo-coords
w = (w1, w2)
  by (cases Rep-homo-coords z, cases Rep-homo-coords w, auto)
let ?x = (z1*w2 - w1*z2) * (cnj z1*cnj w2 - cnj w1*cnj z2)
assume dist-homo-rep z w = 0
hence (cmod ⟨z,w⟩)2 = (⟨z⟩2 * ⟨w⟩2)
  by (subst (asm) dist-homo-rep-zero)
hence Re ?x = 0
  using *
    by (subst (asm) cmod-square) ((subst (asm) norm-homo-rep-square)+, simp
add: inprod-homo-rep-def vec-cnj-def field-simps)
hence ?x = 0
  using complex-mult-cnj-cmod[of z1*w2 - w1*z2]
  by (subst complex-eq-if-Re-eq[of ?x 0]) (simp add: complex-cnj power2-eq-square,
auto)
thus z ≈ w
  using homo-coords-eq-mix[OF *]
  by (auto simp del: homo-coords-eq-def) (metis complex-cnj-cnj complex-cnj-mult)
qed

lemma dist-homo-sym [simp]:
shows dist-homo z1 z2 = dist-homo z2 z1
by transfer (simp add: dist-homo-rep-def split-def Let-def field-simps)

```

Triangle inequality

```

lemma dist-homo-triangle-finite: cmod(a - b) / (sqrt (1+(cmod a)2) * sqrt (1+(cmod
b)2)) ≤ cmod (a - c) / (sqrt (1+(cmod a)2) * sqrt (1+(cmod c)2)) + cmod (c -
b) / (sqrt (1+(cmod b)2) * sqrt (1+(cmod c)2))

```

proof–

```
let ?cc = 1+(cmod c)^2 and ?bb = 1+(cmod b)^2 and ?aa = 1+(cmod a)^2
have sqrt ?cc > 0 sqrt ?aa > 0 sqrt ?bb > 0
  by (auto simp add: power2-eq-square) (metis add-strict-increasing norm-ge-zero
norm-mult zero-less-one)+

have (a - b)*(1+cnj c*c) = (a-c)*(1+cnj c*b) + (c-b)*(1 + cnj c*a)
  by (simp add: field-simps)
moreover
have cmod ((a - b)*(1+cnj c*c)) = cmod(a - b) * (1+(cmod c)^2)
  using complex-mult-cnj-cmod[of cnj c]
  by (auto simp add: power2-eq-square) (metis abs-add-abs abs-one abs-power2
norm-of-real of-real-1 of-real-add of-real-mult power2-eq-square)
ultimately
have cmod(a - b) * (1+(cmod c)^2) ≤ cmod (a-c) * cmod (1+cnj c*b) + cmod
(c-b) * cmod(1 + cnj c*a)
  using complex-mod-triangle-ineq2[of (a-c)*(1+cnj c*b) (c-b)*(1 + cnj c*a)]
  by simp
moreover
have *: ∫ a b c d b' d'. [|b ≤ b'; d ≤ d'; a ≥ (0::real); c ≥ 0|] ==> a*b + c*d
≤ a*b' + c*d'
  by (metis add-mono comm-mult-left-mono)
have cmod (a-c) * cmod (1+cnj c*b) + cmod (c-b) * cmod(1 + cnj c*a) ≤
cmod (a - c) * (sqrt (1+(cmod c)^2) * sqrt (1+(cmod b)^2)) + cmod (c - b) *
(sqrt (1+(cmod c)^2) * sqrt (1+(cmod a)^2))
  using *[OF cmod-1-plus-mult-le[of cnj c b] cmod-1-plus-mult-le[of cnj c a], of
cmod (a-c) cmod (c-b)]
  by (simp add: field-simps real-sqrt-mult[symmetric])
ultimately
have cmod(a - b) * ?cc ≤ cmod (a - c) * sqrt ?cc * sqrt ?bb + cmod (c - b)
* sqrt ?cc * sqrt ?aa
  by simp
moreover
hence 0 ≤ ?cc * sqrt ?aa * sqrt ?bb
  using mult-right-mono[of 0 sqrt ?aa sqrt ?bb]
  using mult-right-mono[of 0 ?cc sqrt ?aa * sqrt ?bb]
  by simp
moreover
have sqrt ?cc / ?cc = 1 / sqrt ?cc
  using (sqrt ?cc > 0)
  by (simp add: field-simps) (metis abs-of-pos real-sqrt-abs2 real-sqrt-mult-distrib2)
hence sqrt ?cc / (?cc * sqrt ?aa) = 1 / (sqrt ?aa * sqrt ?cc)
  using times-divide-eq-right[of 1/sqrt ?aa sqrt ?cc ?cc]
  using (sqrt ?aa > 0)
  by simp
hence cmod (a - c) * sqrt ?cc / (?cc * sqrt ?aa) = cmod (a - c) / (sqrt ?aa
* sqrt ?cc)
  using times-divide-eq-right[of cmod (a - c) sqrt ?cc (?cc * sqrt ?aa)]
  by simp
```

```

moreover
have sqrt ?cc / ?cc = 1 / sqrt ?cc
  using (sqrt ?cc > 0)
  by (simp add: field-simps) (metis abs-of-pos real-sqrt-abs2 real-sqrt-mult-distrib2)
hence sqrt ?cc / (?cc * sqrt ?bb) = 1 / (sqrt ?bb * sqrt ?cc)
  using times-divide-eq-right[of 1/sqrt ?bb sqrt ?cc ?cc]
  using (sqrt ?bb > 0)
  by simp
hence cmod (c - b) * sqrt ?cc / (?cc * sqrt ?bb) = cmod (c - b) / (sqrt ?bb *
sqrt ?cc)
  using times-divide-eq-right[of cmod (c - b) sqrt ?cc ?cc * sqrt ?bb]
  by simp
ultimately
show ?thesis
  using divide-right-mono[of cmod (a - b) * ?cc cmod (a - c) * sqrt ?cc * sqrt
?bb + cmod (c - b) * sqrt ?cc * sqrt ?aa ?cc * sqrt ?aa * sqrt ?bb] (sqrt ?aa >
0) (sqrt ?bb > 0) (sqrt ?cc > 0)
  by (simp add: add-divide-distrib)
qed

lemma dist-homo-triangle-infinite1: 1 / sqrt(1 + (cmod b)2) ≤ 1 / sqrt(1 +
(cmod c)2) + cmod (b - c) / (sqrt(1 + (cmod b)2) * sqrt(1 + (cmod c)2))
proof-
  let ?bb = sqrt (1 + (cmod b)2) and ?cc = sqrt (1 + (cmod c)2)
  have ?bb > 0 ?cc > 0
    by (metis add-strict-increasing real-sqrt-gt-0-iff zero-le-power2 zero-less-one)+
  hence *: ?bb * ?cc ≥ 0
    by (metis one-power2 real-sqrt-mult-distrib2 real-sqrt-sum-squares-mult-ge-zero)
  have **: (?cc - ?bb) / (?bb * ?cc) = 1 / ?bb - 1 / ?cc
    using (sqrt (1 + (cmod b)2) > 0) (sqrt (1 + (cmod c)2) > 0)
    by (simp add: field-simps)
  show 1 / ?bb ≤ 1 / ?cc + cmod (b - c) / (?bb * ?cc)
    using divide-right-mono[OF cmod-diff-ge[of c b] *]
    by (subst (asm) **) (simp add: field-simps norm-minus-commute)
qed

lemma dist-homo-triangle-infinite2:
  1 / sqrt(1 + (cmod a)2) ≤ cmod (a - c) / (sqrt (1+(cmod a)2) * sqrt (1+(cmod
c)2)) + 1 / sqrt(1 + (cmod c)2)
using dist-homo-triangle-infinite1[of a c]
by simp

lemma dist-homo-triangle-infinite3:
  cmod(a - b) / (sqrt (1+(cmod a)2) * sqrt (1+(cmod b)2)) ≤ 1 / sqrt(1 + (cmod
a)2) + 1 / sqrt(1 + (cmod b)2)
proof-
  let ?aa = sqrt (1 + (cmod a)2) and ?bb = sqrt (1 + (cmod b)2)
  have ?aa > 0 ?bb > 0
    by (metis add-strict-increasing real-sqrt-gt-0-iff zero-le-power2 zero-less-one)+

```

```

hence *: ?aa * ?bb ≥ 0
  by (metis one-power2 real-sqrt-mult-distrib2 real-sqrt-sum-squares-mult-ge-zero)
have **: (?aa + ?bb) / (?aa * ?bb) = 1 / ?aa + 1 / ?bb
  using (?aa > 0) (?bb > 0)
  by (simp add: field-simps)
show cmod (a - b) / (?aa * ?bb) ≤ 1 / ?aa + 1 / ?bb
  using divide-right-mono[OF cmod-diff-le[of a b] *]
  by (subst (asm) **) (simp add: field-simps norm-minus-commute)
qed

lemma dist-homo-triangle:
  shows dist-homo A B ≤ dist-homo A C + dist-homo C B
proof (cases A = ∞h)
  case True
  show ?thesis
  proof (cases B = ∞h)
    case True
    show ?thesis
    proof (cases C = ∞h)
      case True
      show ?thesis
      using ⟨A = ∞h⟩ ⟨B = ∞h⟩ ⟨C = ∞h⟩
      by simp
    next
    case False
    then obtain c where C = of-complex c
      using inf-homo-or-complex-homo[of C]
      by auto
    show ?thesis
    using ⟨A = ∞h⟩ ⟨B = ∞h⟩ ⟨C = of-complex c⟩
    by (simp add: dist-homo-infinite2)
  qed
next
  case False
  then obtain b where B = of-complex b
    using inf-homo-or-complex-homo[of B]
    by auto
  show ?thesis
  proof (cases C = ∞h)
    case True
    show ?thesis
    using ⟨A = ∞h⟩ ⟨C = ∞h⟩ ⟨B = of-complex b⟩
    by simp
  next
    case False
    then obtain c where C = of-complex c
      using inf-homo-or-complex-homo[of C]
      by auto
    show ?thesis
  qed
qed

```

```

using ⟨A = ∞h⟩ ⟨B = of-complex b⟩ ⟨C = of-complex c⟩
using mult-left-mono[OF dist-homo-triangle-infinite1[of b c], of 2]
by (simp add: dist-homo-finite dist-homo-infinite1 dist-homo-infinite2)
qed
qed
next
case False
then obtain a where A = of-complex a
  using inf-homo-or-complex-homo[of A]
  by auto
show ?thesis
proof (cases B = ∞h)
  case True
  show ?thesis
  proof (cases C = ∞h)
    case True
    show ?thesis
    using ⟨B = ∞h⟩ ⟨C = ∞h⟩ ⟨A = of-complex a⟩
    by (simp add: dist-homo-infinite2)
  next
    case False
    then obtain c where C = of-complex c
      using inf-homo-or-complex-homo[of C]
      by auto
    show ?thesis
    using ⟨B = ∞h⟩ ⟨C = of-complex c⟩ ⟨A = of-complex a⟩
    using mult-left-mono[OF dist-homo-triangle-infinite2[of a c], of 2]
    by (simp add: dist-homo-finite dist-homo-infinite1 dist-homo-infinite2)
  qed
next
case False
then obtain b where B = of-complex b
  using inf-homo-or-complex-homo[of B]
  by auto
show ?thesis
proof (cases C = ∞h)
  case True
  thus ?thesis
    using ⟨C = ∞h⟩ ⟨B = of-complex b⟩ ⟨A = of-complex a⟩
    using mult-left-mono[OF dist-homo-triangle-infinite3[of a b], of 2]
    by (simp add: dist-homo-finite dist-homo-infinite1 dist-homo-infinite2)
next
case False
then obtain c where C = of-complex c
  using inf-homo-or-complex-homo[of C]
  by auto
show ?thesis
  using ⟨A = of-complex a⟩ ⟨B = of-complex b⟩ ⟨C = of-complex c⟩
  using mult-left-mono[OF dist-homo-triangle-finite[of a b c], of 2]

```

```

    by (simp add: dist-homo-finite norm-minus-commute)
qed
qed
qed

instantiation complex-homo :: metric-space
begin
definition dist-complex-homo = dist-homo
definition open-complex-homo S = ( $\forall x \in S. \exists e > 0. \forall y. dist\text{-homo } y x < e \longrightarrow y \in S$ )
instance
proof
fix x y :: complex-homo
show (dist x y = 0) = (x = y)
unfolding dist-complex-homo-def
using dist-homo-zero1[of x] dist-homo-zero2[of x y]
by auto
next
fix S :: complex-homo set
show open S = ( $\forall x \in S. \exists e > 0. \forall y. dist\text{ } y x < e \longrightarrow y \in S$ )
unfolding open-complex-homo-def dist-complex-homo-def
by simp
next
fix x y z :: complex-homo
show dist x y  $\leq$  dist x z + dist y z
unfolding dist-complex-homo-def
using dist-homo-triangle[of x y z]
by simp
qed
end

theory RiemannSphere
imports HomogeneousCoordinates  $\sim\sim$ /src/HOL/Library/Product-Vector
begin

lemma Lim-within: ( $f \dashrightarrow l$ ) (at a within S)  $\longleftrightarrow$ 
 $(\forall e > 0. \exists d > 0. \forall x \in S. 0 < dist\text{ } x a \wedge dist\text{ } x a < d \longrightarrow dist\text{ } (f x) l < e)$ 
by (auto simp add: tends-to-iff eventually-at dist-nz)

lemma continuous-on-iff:
continuous-on s f  $\longleftrightarrow$ 
 $(\forall x \in s. \forall e > 0. \exists d > 0. \forall x' \in s. dist\text{ } x' x < d \longrightarrow dist\text{ } (f x') (f x) < e)$ 
unfolding continuous-on-def Lim-within
apply (intro ball-cong [OF refl] all-cong ex-cong)
apply (rename-tac y, case-tac y = x)
apply simp

```

```

apply (simp add: dist-nz)
done

```

9 Riemann sphere

```

typedef riemann-sphere = {(x::real, y::real, z::real). x*x + y*y + z*z = 1}
by (rule-tac x=(1, 0, 0) in exI) simp

```

```

lemma sphere-bounds':
assumes x*x + y*y + z*z = (1::real)
shows -1 ≤ x ∧ x ≤ 1
proof-
from assms have x*x ≤ 1
  by (smt real-minus-mult-self-le)
hence x2 ≤ 12 (- x)2 ≤ 12
  by (auto simp add: power2-eq-square)
show -1 ≤ x ∧ x ≤ 1
proof (cases x ≥ 0)
  case True
  thus ?thesis
    using square-cancel[OF x*x ≤ 12]
    by simp
next
  case False
  thus ?thesis
    using square-cancel[OF (-x)2 ≤ 12]
    by simp
qed
qed

```

```

lemma sphere-bounds:
assumes x*x + y*y + z*z = (1::real)
shows -1 ≤ x ∧ x ≤ 1 -1 ≤ y ∧ y ≤ 1 -1 ≤ z ∧ z ≤ 1
using assms
using sphere-bounds'[of x y z] sphere-bounds'[of y x z] sphere-bounds'[of z x y]
by (auto simp add: field-simps)

```

Polar coords parametrization

```

lemma sphere-params-on-sphere:
assumes x = cos α * cos β y = cos α * sin β z = sin α
shows x*x + y*y + z*z = 1
proof-
have x*x + y*y = (cos α * cos α) * (cos β * cos β) + (cos α * cos α) * (sin
β * sin β)
  using assms
  by simp
hence x*x + y*y = cos α * cos α
  using sin-cos-squared-add3[of β]
  by (subst (asm) distrib-left[symmetric]) (simp add: field-simps)

```

```

thus ?thesis
  using assms
  using sin-cos-squared-add3[of α]
  by simp
qed

lemma sphere-params:
  assumes "x*x + y*y + z*z = 1"
  shows "x = cos (arcsin z) * cos (atan2 y x) ∧ y = cos (arcsin z) * sin (atan2 y
x) ∧ z = sin (arcsin z)"
  proof (cases "z=1 ∨ z = -1")
    case True
    hence "x = 0 ∧ y = 0"
      using assms
      by auto
    thus ?thesis
      using ⟨z = 1 ∨ z = -1⟩
      by (auto simp add: cos-arcsin)
  next
    case False
    hence "x ≠ 0 ∨ y ≠ 0"
      using assms
      by auto (metis minus-one square-eq-1-iff)
    thus ?thesis
      using sphere-bounds[OF assms] assms
      by (auto simp add: cos-arcsin cos-arctan sin-arctan power2-eq-square field-simps
real-sqrt-divide atan2-def cos-periodic-pi2 cos-periodic-pi3 sin-periodic-pi3) (smt real-sqrt-abs2)+
qed

```

```

lemma ex-sphere-params:
  assumes "x*x + y*y + z*z = 1"
  shows "∃ α β. x = cos α * cos β ∧ y = cos α * sin β ∧ z = sin α ∧ -pi / 2
≤ α ∧ α ≤ pi / 2 ∧ -pi ≤ β ∧ β < pi"
  using assms arcsin-bounded[of z] sphere-bounds[of x y z]
  by (rule-tac x=arcsin z in exI, rule-tac x=atan2 y x in exI) (simp add: sphere-params
arcsin-bounded atan2-bounded)

```

Stereographic and inverse stereographic projection

```

definition stereographic-coords :: riemann-sphere ⇒ homo-coords where
stereographic-coords M = (let (x, y, z) = Rep-riemann-sphere M in
  (if (x, y, z) ≠ (0, 0, 1) then
    Abs-homo-coords (Complex x y, complex-of-real (1 - z))
  else
    Abs-homo-coords (1, 0)
  ))

```

```

lemma stereographic-coords-rep:
  Rep-homo-coords (stereographic-coords M) = (let (x, y, z) = Rep-riemann-sphere
M in

```

```

(if (x, y, z) ≠ (0, 0, 1) then
    (Complex x y, complex-of-real (1 - z))
else
    (1, 0)
))
proof-
obtain x y z where MM: (x, y, z) = Rep-riemann-sphere M
by (cases Rep-riemann-sphere M) auto
show ?thesis
proof (cases (x, y, z) ≠ (0, 0, 1) )
case True
thus ?thesis
using MM[symmetric] Abs-homo-coords-inverse[of (Complex x y, 1 - cor z)]
using Rep-riemann-sphere[of M]
by (cases x = 0 ∧ y = 0, cases z=1) (auto simp add: stereographic-coords-def,
metis Complex-eq-1 complex-of-real-def)
next
case False
thus ?thesis
using MM
by (simp add: stereographic-coords-def)
qed
qed

```

lift-definition stereographic :: riemann-sphere ⇒ complex-homo **is** stereographic-coords
by (simp del: homo-coords-eq-def)

definition inv-stereographic-coords :: homo-coords ⇒ riemann-sphere **where**
inv-stereographic-coords z = (
let (z1, z2) = Rep-homo-coords z
in if z2 = 0 then
Abs-riemann-sphere (0, 0, 1)
else
let z = z1/z2;
X = Re (2*z / (1 + z*c conj z));
Y = Im (2*z / (1 + z*c conj z));
Z = ((cmod z)^2 - 1) / (1 + (cmod z)^2)
in Abs-riemann-sphere (X, Y, Z))

lift-definition inv-stereographic :: complex-homo ⇒ riemann-sphere **is** inv-stereographic-coords
by (auto simp add: inv-stereographic-coords-def split-def Let-def)

lemma one-plus-square-neq-zero [simp]:
fixes x :: real
shows 1 + (cor x)^2 ≠ 0
by (metis (hide-lams, no-types) of-real-1 of-real-add of-real-eq-0-iff of-real-power
power-one sum-power2-eq-zero-iff zero-neq-one)

lemma Re-stereographic: Re (2 * z / (1 + z * cnj z)) = 2 * Re z / (1 + (cmod

$z)^2)$
using one-plus-square-neq-zero
by (subst complex-mult-cnj-cmod, subst Re-divide-real) (auto simp add: power2-eq-square)

lemma Im-stereographic: $\text{Im} (2 * z / (1 + z * \text{cnj } z)) = 2 * \text{Im } z / (1 + (\text{cmod } z)^2)$
using one-plus-square-neq-zero
by (subst complex-mult-cnj-cmod, subst Im-divide-real) (auto simp add: power2-eq-square)

lemma inv-stereographic-on-sphere:
assumes $X = \text{Re} (2*z / (1 + z*\text{cnj } z))$ $Y = \text{Im} (2*z / (1 + z*\text{cnj } z))$ $Z = ((\text{cmod } z)^2 - 1) / (1 + (\text{cmod } z)^2)$
shows $X*X + Y*Y + Z*Z = 1$
proof-
have $1 + (\text{cmod } z)^2 \neq 0$
by (metis power-one realpow-two-sum-zero-iff zero-neq-one)
thus ?thesis
using assms
by (simp add: Re-stereographic Im-stereographic) (cases z, simp add: power2-eq-square real-sqrt-mult[symmetric] add-divide-distrib[symmetric], simp add: field-simps)
qed

lemma inv-stereographic-coords-Rep:
 $\text{Rep-riemann-sphere} (\text{inv-stereographic-coords } z) =$
 $(\text{let } (z1, z2) = \text{Rep-homo-coords } z$
 $\quad \text{in if } z2 = 0 \text{ then}$
 $\quad \quad (0, 0, 1)$
 $\quad \text{else}$
 $\quad \quad \text{let } z = z1/z2;$
 $\quad \quad X = \text{Re} (2*z / (1 + z*\text{cnj } z));$
 $\quad \quad Y = \text{Im} (2*z / (1 + z*\text{cnj } z));$
 $\quad \quad Z = ((\text{cmod } z)^2 - 1) / (1 + (\text{cmod } z)^2)$
 $\quad \text{in } (X, Y, Z))$
proof-
obtain z1 z2 **where** zz: Rep-homo-coords z = (z1, z2)
by (rule obtain-homo-coords)
show ?thesis
proof (cases z2 = 0)
case True
thus ?thesis
using zz
by (simp add: Let-def inv-stereographic-coords-def Abs-riemann-sphere-inverse)
next
case False
thus ?thesis
using inv-stereographic-on-sphere[of - z1/z2] zz
by (simp add: Let-def inv-stereographic-coords-def Abs-riemann-sphere-inverse)
qed

qed

definition [*simp*]: *North* = *Abs-riemann-sphere* (0, 0, 1)

lemma *stereographic-North*: *stereographic* $x = \infty_h \longleftrightarrow x = \text{North}$

proof (*transfer*)

fix x

show *stereographic-coords* $x \approx \text{inf-homo-rep} \longleftrightarrow x = \text{North}$

proof

assume $x = \text{North}$

thus *stereographic-coords* $x \approx \text{inf-homo-rep}$

by (*simp add: stereographic-coords-def Abs-riemann-sphere-inverse Abs-homo-coords-inverse*)

next

assume $*: \text{stereographic-coords } x \approx \text{inf-homo-rep}$

show $x = \text{North}$

proof (*cases Rep-riemann-sphere* $x = (0, 0, 1)$)

case *True*

thus *?thesis*

by *auto (metis Rep-riemann-sphere-inverse)*

next

case *False*

thus *?thesis*

using *

using *Rep-riemann-sphere*[*of x*]

by (*auto simp add: stereographic-coords-def split-def Let-def Abs-homo-coords-inverse complex-of-real-def split: split-if-asm*) (*metis pair-collapse*)

qed

qed

qed

lemma *stereographic-inv-stereographic'*:

assumes

$z: z = z1/z2 \text{ and } z2 \neq 0 \text{ and }$

$X: X = \text{Re}(2*z / (1 + z*cnj z)) \text{ and } Y: Y = \text{Im}(2*z / (1 + z*cnj z)) \text{ and }$

$Z: Z = ((cmod z)^2 - 1) / (1 + (cmod z)^2)$

shows $\exists k. k \neq 0 \wedge (\text{Complex } X Y, \text{complex-of-real } (1 - Z)) = k *_{sv} (z1, z2)$

proof-

have $1 + (cmod z)^2 \neq 0$

by (*metis one-power2 sum-power2-eq-zero-iff zero-neq-one*)

hence $\text{cor}(1 - Z) = 2 / \text{cor}(1 + (cmod z)^2)$

using *Z*

by (*simp add: field-simps complex-of-real-def*)

moreover

have $X = 2 * \text{Re}(z) / (1 + (cmod z)^2)$

using *X*

by (*simp add: Re-stereographic*)

have $Y = 2 * \text{Im}(z) / (1 + (cmod z)^2)$

using *Y*

by (*simp add: Im-stereographic*)

```

have Complex X Y = 2 * z / cor (1 + (cmod z)2)
  using <1 + (cmod z)2 ≠ 0>
  by (subst <X = 2*Re(z) / (1 + (cmod z)2)>, subst <Y = 2*Im(z) / (1 +
(cmod z)2)>, simp add: Complex-scale4 Complex-scale1 of-real-numeral)
moreover
have 1 + (cor (cmod (z1 / z2)))2 ≠ 0
  by (rule one-plus-square-neq-zero)
ultimately
show ?thesis
  using <z2 ≠ 0> <1 + (cmod z)2 ≠ 0>
  by (simp, subst z)+
    (rule-tac x=(2 / (1 + (cor (cmod (z1 / z2)))2)) / z2 in exI, auto)
qed

lemma stereographic-inv-stereographic:
  stereographic (inv-stereographic z) = z
proof transfer
  fix z
  obtain z1 z2 where zz: Rep-homo-coords z = (z1, z2)
    by (rule obtain-homo-coords)
  have z ≈ stereographic-coords (inv-stereographic-coords z)
  proof (cases z2 = 0)
    case True
    thus ?thesis
      using zz Rep-homo-coords[of z]
      by (simp add: stereographic-coords-def inv-stereographic-coords-Rep)
  next
    case False
    thus ?thesis
      using zz stereographic-inv-stereographic'[of z1/z2 z1 z2]
      by (simp add: stereographic-coords-rep inv-stereographic-coords-Rep Let-def)
  qed
  thus stereographic-coords (inv-stereographic-coords z) ≈ z
    by (rule homo-coords-eq-sym)
qed

lemma bij-stereographic: bij stereographic
unfolding bij-def inj-on-def surj-def
proof (safe)
  fix x y
  assume stereographic x = stereographic y
  thus x = y
  proof (transfer)
    fix a b
    assume *: stereographic-coords a ≈ stereographic-coords b
    obtain xa ya za xb yb zb where **: Rep-riemann-sphere a = (xa, ya, za)
      Rep-riemann-sphere b = (xb, yb, zb)
      by (metis prod-cases3)

```

```

show a = b
proof (subst Rep-riemann-sphere-inject[symmetric])
  show Rep-riemann-sphere a = Rep-riemann-sphere b
  proof (cases Rep-riemann-sphere a = (0, 0, 1))
    case True
    thus ?thesis
      using * ** Rep-riemann-sphere[of b]
      unfolding stereographic-coords-def
      by (cases zb=1) (auto simp add: Abs-homo-coords-inverse complex-of-real-def)
next
  {
    fix k
    assume xa * xa + (ya * ya + za * za) = 1
    zb * zb + (k * (k * (xa * xa)) + k * (k * (ya * ya))) = 1
    zb ≠ 1 za ≠ 1 k ≠ 0 1 + k * za = k + zb k ≠ 1
    hence False
    by algebra
  } note *** = this

case False
thus ?thesis
  using * ** Rep-riemann-sphere[of a] Rep-riemann-sphere[of b]
  unfolding stereographic-coords-def
  apply (case-tac[!] zb = 1, case-tac[!] za = 1)
  apply (auto simp add: Abs-homo-coords-inverse complex-of-real-def)
  apply (case-tac[!] k)
  using ***
  apply (auto simp add: field-simps)
  apply (case-tac real1 = 1)
  by auto
qed
qed
qed
next
  fix a
  show ∃ b. a = stereographic b
  by (rule-tac x=inv-stereographic a in exI) (simp add: stereographic-inv-stereographic)
qed

lemma inv-stereographic-stereographic:
  inv-stereographic (stereographic x) = x
using stereographic-inv-stereographic[of stereographic x]
using bij-stereographic
unfolding bij-def inj-on-def
by simp

lemma inv-stereographic-is-inv:
  inv-stereographic = inv stereographic

```

```

by (rule inv-equality[symmetric], simp-all add: inv-stereographic-stereographic stereographic-inv-stereographic)

Circles on the sphere

type-synonym real-vec-4 = real × real × real × real

fun mult-sv :: real ⇒ real-vec-4 ⇒ real-vec-4 (infixl *sv4 100) where
  k *sv4 (a, b, c, d) = (k*a, k*b, k*c, k*d)

typedef plane-vec = {(a::real, b::real, c::real, d::real). a ≠ 0 ∨ b ≠ 0 ∨ c ≠ 0 ∨
d ≠ 0}
by (rule-tac x=(1, 1, 1, 1) in exI) simp

definition plane-vec-eq where
  plane-vec-eq v1 v2 ←→ (∃ k. k ≠ 0 ∧ Rep-plane-vec v2 = k *sv4 Rep-plane-vec
v1)

lemma [simp]: 1 *sv4 x = x
by (cases x) simp

lemma [simp]: x *sv4 (y *sv4 v) = (x*y) *sv4 v
by (cases v) simp

quotient-type plane = plane-vec / plane-vec-eq
proof (rule equivpI)
  show reflp plane-vec-eq
    unfolding reflp-def
    by (auto simp add: plane-vec-eq-def) (rule-tac x=1 in exI, simp)
next
  show symp plane-vec-eq
    unfolding symp-def
    by (auto simp add: plane-vec-eq-def) (rule-tac x=1/k in exI, simp)
next
  show transp plane-vec-eq
    unfolding transp-def
    by (auto simp add: plane-vec-eq-def) (rule-tac x=ka*k in exI, simp)
qed

definition on-sphere-circle-rep where
  on-sphere-circle-rep α A ←→
    (let (X, Y, Z) = Rep-riemann-sphere A;
     (a, b, c, d) = Rep-plane-vec α
     in a*X + b*Y + c*Z + d = 0)

lift-definition on-sphere-circle :: plane ⇒ riemann-sphere ⇒ bool is on-sphere-circle-rep
proof –
  fix v1 v2
  obtain a1 b1 c1 d1 where vv1: Rep-plane-vec v1 = (a1, b1, c1, d1)
    by (cases Rep-plane-vec v1) auto
  obtain a2 b2 c2 d2 where vv2: Rep-plane-vec v2 = (a2, b2, c2, d2)

```

```

by (cases Rep-plane-vec v2) auto
assume plane-vec-eq v1 v2
then obtain k where *: a2 = k*a1 b2 = k*b1 c2 = k*c1 d2 = k*d1 k ≠ 0
  using vv1 vv2
    by (auto simp add: plane-vec-eq-def)
show on-sphere-circle-rep v1 = on-sphere-circle-rep v2
proof (rule ext)
  fix M
  obtain x y z where MM: Rep-riemann-sphere M = (x, y, z)
    by (cases Rep-riemann-sphere M) auto
  have k * a1 * x + k * b1 * y + k * c1 * z + k * d1 = k*(a1*x + b1*y +
  c1*z + d1)
    by (simp add: field-simps)
  thus on-sphere-circle-rep v1 M = on-sphere-circle-rep v2 M
    using vv1 vv2 MM *
      by (auto simp add: plane-vec-eq-def on-sphere-circle-rep-def split-def Let-def)
  qed
qed

```

definition sphere-circle-set **where**

$$\text{sphere-circle-set } \alpha = \{A. \text{on-sphere-circle } \alpha \ A\}$$

Distance on the Riemann sphere

```

definition dist-riemann-sphere' where
dist-riemann-sphere' M1 M2 =
  (let (x1, y1, z1) = Rep-riemann-sphere M1;
   (x2, y2, z2) = Rep-riemann-sphere M2
   in norm (x1 - x2, y1 - y2, z1 - z2))

```

```

lemma dist-riemann-sphere'-inner:
  (dist-riemann-sphere' M1 M2)2 = 2 - 2 * inner (Rep-riemann-sphere M1)
  (Rep-riemann-sphere M2)
using Rep-riemann-sphere[of M1] Rep-riemann-sphere[of M2]
unfolding dist-riemann-sphere'-def
by (auto simp add: norm-prod-def) (simp add: power2-eq-square field-simps)

```

```

lemma xxx [simp]:
  Re (2 * m1 / (1 + cor ((cmod m1)2))) = 2 * Re m1 / (1 + (cmod m1)2)
apply (subst Re-divide-real)
apply (simp add: power2-eq-square)
apply (metis numeral_One of-real_1 of-real-add of-real-eq_0 iff power-one sum-power2-eq-zero-iff
zero-neq-numeral)
apply (simp add: power2-eq-square)
done

```

```

lemma yyy [simp]:
  Im (2 * m1 / (1 + cor ((cmod m1)2))) = 2 * Im m1 / (1 + (cmod m1)2)
apply (subst Im-divide-real)
apply (simp add: power2-eq-square)

```

```

apply (metis numeral_One of-real_1 of-real-add of-real-eq_0 iff power-one sum-power2_eq_zero_if
zero_neq_numeral)
apply (simp add: power2_eq_square)
done

lemma dist_riemann_sphere'_ge_0 [simp]: dist_riemann_sphere' M1 M2 ≥ 0
using norm_ge_zero
unfolding dist_riemann_sphere'_def
by (simp add: split_def Let_def)

lemma dist_homo_stereographic_finite:
assumes stereographic M1 = of_complex m1 stereographic M2 = of_complex m2
shows dist_riemann_sphere' M1 M2 = 2 * cmod (m1 - m2) / (sqrt (1 + (cmod m1)^2) * sqrt (1 + (cmod m2)^2))
proof-
obtain x1 y1 z1 x2 y2 z2 where MM: (x1, y1, z1) = Rep_riemann_sphere M1
(x2, y2, z2) = Rep_riemann_sphere M2
by (cases Rep_riemann_sphere M1, cases Rep_riemann_sphere M2, auto, blast)
have *: M1 = inv_stereographic (of_complex m1) M2 = inv_stereographic (of_complex m2)
using inv_stereographic_is_inv_assms
by (metis inv_stereographic_stereographic)+
have (1 + (cmod m1)^2) ≠ 0 (1 + (cmod m2)^2) ≠ 0
by (metis power_one realpow_two_sum_zero_iff zero_neq_one)+
have (1 + (cmod m1)^2) > 0 (1 + (cmod m2)^2) > 0
by (smt realpow_square_minus_le)+
hence (1 + (cmod m1)^2) * (1 + (cmod m2)^2) > 0
by (metis norm_mult_less norm_zero power2_eq_square zero_power2)
hence sqrt ((1 + cmod m1 * cmod m1) * (1 + cmod m2 * cmod m2)) > 0
using real_sqrt_gt_0_iff
by (simp add: power2_eq_square)
hence **: (2 * cmod (m1 - m2) / sqrt ((1 + cmod m1 * cmod m1) * (1 + cmod m2 * cmod m2))) ≥ 0 ↔ cmod (m1 - m2) ≥ 0
by (metis diff_self_divide_nonneg_pos mult_2 norm_ge_zero norm_triangle_ineq4
norm_zero)

have (dist_riemann_sphere' M1 M2)^2 * (1 + (cmod m1)^2) * (1 + (cmod m2)^2)
= 4 * (cmod (m1 - m2))^2
apply (subst *)+
proof transfer
fix m1 m2
have (1 + (cmod m1)^2) ≠ 0 (1 + (cmod m2)^2) ≠ 0
by (metis power_one realpow_two_sum_zero_iff zero_neq_one)+
thus (dist_riemann_sphere' (inv_stereographic_coords (of_complex_coords m1))
(inv_stereographic_coords (of_complex_coords m2)))^2 * (1 + (cmod m1)^2) * (1 + (cmod m2)^2) = 4 * (cmod (m1 - m2))^2
apply (simp add: dist_riemann_sphere'_inner inv_stereographic_coords_Rep
complex_mult_cnj_cmod)
apply (subst cor_squared)+
```

```

apply (subst xxx)+
apply (subst yyy)+
apply (subst left-diff-distrib[of 2])
apply (subst left-diff-distrib[of 2*(1+(cmod m1)^2)])
apply (subst distrib-right[of -- (1 + (cmod m1)^2)])
apply (subst distrib-right[of -- (1 + (cmod m1)^2)])
apply (subst distrib-right[of 2 * (2 * Re m1 / (1 + (cmod m1)^2) * (2 * Re
m2 / (1 + (cmod m2)^2))) * (1 + (cmod m1)^2) - (1 + (cmod m2)^2)])
apply (subst distrib-right[of 2 * (2 * Im m1 / (1 + (cmod m1)^2) * (2 * Im
m2 / (1 + (cmod m2)^2))) * (1 + (cmod m1)^2) - (1 + (cmod m2)^2)])
apply simp
apply (subst (asm) cmod-square)+
apply (subst cmod-square)+
apply (simp add: field-simps)
done
qed
hence (dist-riemann-sphere' M1 M2)^2 = 4 * (cmod (m1 - m2))^2 / ((1 + (cmod
m1)^2) * (1 + (cmod m2)^2))
  using <(1 + (cmod m1)^2) ≠ 0> <(1 + (cmod m2)^2) ≠ 0>
  using eq-divide-imp[of (1 + (cmod m1)^2) * (1 + (cmod m2)^2) (dist-riemann-sphere'
M1 M2)^2 4 * (cmod (m1 - m2))^2]
  by simp
thus dist-riemann-sphere' M1 M2 = 2 * cmod (m1 - m2) / (sqrt (1 + (cmod
m1)^2) * sqrt (1 + (cmod m2)^2))
  using power2-eq-iff[of dist-riemann-sphere' M1 M2 2 * (cmod (m1 - m2)) /
sqrt ((1 + (cmod m1)^2) * (1 + (cmod m2)^2))]
  using <(1 + (cmod m1)^2) * (1 + (cmod m2)^2) > 0> <(1 + (cmod m1)^2) > 0>
<(1 + (cmod m2)^2) > 0>
  apply (auto simp add: power2-eq-square real-sqrt-mult[symmetric])
  using dist-riemann-sphere'-ge-0[of M1 M2] **
  by simp
qed

```

lemma dist-homo-stereographic-infinite:

assumes stereographic M1 = ∞_h stereographic M2 = of-complex m2
shows dist-riemann-sphere' M1 M2 = 2 / sqrt (1 + (cmod m2)^2)

proof –

obtain x2 y2 z2 where MM: (0, 0, 1) = Rep-riemann-sphere M1 (x2, y2, z2)
= Rep-riemann-sphere M2
 using stereographic M1 = ∞_h
 using stereographic-North[of M1]
 by (cases Rep-riemann-sphere M2, auto simp add: Abs-riemann-sphere-inverse)
have *: M1 = inv-stereographic ∞_h M2 = inv-stereographic (of-complex m2)
 using inv-stereographic-is-inv assms
 by (metis inv-stereographic-stereographic)+
have (1 + (cmod m2)^2) ≠ 0
 by (metis power-one realpow-two-sum-zero-iff zero-neq-one)+
have (1 + (cmod m2)^2) > 0
 by (smt realpow-square-minus-le)+

```

hence sqrt (1 + cmod m2 * cmod m2) > 0
  using real-sqrt-gt-0-iff
  by (simp add: power2-eq-square)
hence **: 2 / sqrt (1 + cmod m2 * cmod m2) > 0
  by simp

have (dist-riemann-sphere' M1 M2)2 * (1 + (cmod m2)2) = 4
  apply (subst *)+
proof transfer
  fix m2
  have (1 + (cmod m2)2) ≠ 0
    by (metis power-one realpow-two-sum-zero-iff zero-neq-one)
  thus (dist-riemann-sphere' (inv-stereographic-coords inf-homo-rep) (inv-stereographic-coords
  (of-complex-coords m2)))2 * (1 + (cmod m2)2) = 4
    by (simp add: dist-riemann-sphere'-inner inv-stereographic-coords-Rep complex-mult-cnj-cmod)
      (subst left-diff-distrib[of 2], simp)
qed
hence (dist-riemann-sphere' M1 M2)2 = 4 / (1 + (cmod m2)2)
  using ‹(1 + (cmod m2)2) ≠ 0›
  by (simp add: field-simps)
thus dist-riemann-sphere' M1 M2 = 2 / sqrt (1 + (cmod m2)2)
  using power2-eq-iff[of dist-riemann-sphere' M1 M2 2 / sqrt (1 + (cmod m2)2)]
  using ‹(1 + (cmod m2)2) > 0›
  apply (auto simp add: power2-eq-square real-sqrt-mult[symmetric])
  using dist-riemann-sphere'-ge-0[of M1 M2] **
  by simp
qed

lemma dist-riemann-sphere'-sym: dist-riemann-sphere' M1 M2 = dist-riemann-sphere'
M2 M1
proof-
  obtain x1 y1 z1 x2 y2 z2 where MM: (x1, y1, z1) = Rep-riemann-sphere M1
  (x2, y2, z2) = Rep-riemann-sphere M2
    by (cases Rep-riemann-sphere M1, cases Rep-riemann-sphere M2, auto, blast)
  show ?thesis
    unfolding dist-riemann-sphere'-def
    using norm-minus-cancel[of (x1 - x2, y1 - y2, z1 - z2)] MM[symmetric]
    by simp
qed

lemma dist-homo-stereographic: dist-riemann-sphere' M1 M2 = dist-homo (stereographic
M1) (stereographic M2)
proof (cases M1 = North)
  case True
  hence stereographic M1 = ∞h
    by (simp add: stereographic-North)
  show ?thesis
proof (cases M2 = North)
  case True

```

```

show ?thesis
  using ⟨M1 = North⟩ ⟨M2 = North⟩
    by (auto simp add: Abs-riemann-sphere-inverse dist-riemann-sphere'-def
norm-prod-def)
next
  case False
  hence stereographic M2 ≠ ∞h
    using stereographic-North[of M2]
    by simp
  then obtain m2 where stereographic M2 = of-complex m2
    using inf-homo-or-complex-homo[of stereographic M2]
    by auto
  show ?thesis
    using ⟨stereographic M2 = of-complex m2⟩ ⟨stereographic M1 = ∞h⟩
    using dist-homo-infinite1 dist-homo-stereographic-infinite
    by simp
qed
next
  case False
  hence stereographic M1 ≠ ∞h
    by (simp add: stereographic-North)
  then obtain m1 where stereographic M1 = of-complex m1
    using inf-homo-or-complex-homo[of stereographic M1]
    by auto
  show ?thesis
proof (cases M2 = North)
  case True
  hence stereographic M2 = ∞h
    by (simp add: stereographic-North)
  show ?thesis
    using ⟨stereographic M1 = of-complex m1⟩ ⟨stereographic M2 = ∞h⟩
    using dist-homo-infinite2 dist-homo-stereographic-infinite
    by (subst dist-riemann-sphere'-sym, simp)
next
  case False
  hence stereographic M2 ≠ ∞h
    by (simp add: stereographic-North)
  then obtain m2 where stereographic M2 = of-complex m2
    using inf-homo-or-complex-homo[of stereographic M2]
    by auto
  show ?thesis
    using ⟨stereographic M1 = of-complex m1⟩ ⟨stereographic M2 = of-complex
m2⟩
    using dist-homo-finite dist-homo-stereographic-finite
    by simp
qed
qed

```

lemma dist-homo-stereographic':

```

dist-homo A B = dist-riemann-sphere' (inv-stereographic A) (inv-stereographic B)
by (subst dist-homo-stereographic) (metis stereographic-inv-stereographic)

instantiation riemann-sphere :: metric-space
begin
definition dist-riemann-sphere = dist-riemann-sphere'
definition open-riemann-sphere S = ( $\forall x \in S. \exists e > 0. \forall y. dist\text{-riemann-sphere}' y x < e \rightarrow y \in S$ )
instance
proof
  fix x y :: riemann-sphere
  show (dist x y = 0) = (x = y)
  proof-
    obtain x1 y1 z1 x2 y2 z2 where MM: (x1, y1, z1) = Rep-riemann-sphere x
    (x2, y2, z2) = Rep-riemann-sphere y
    by (cases Rep-riemann-sphere x, cases Rep-riemann-sphere y, auto, blast)
    show ?thesis
    unfolding dist-riemann-sphere-def
    using norm-eq-zero[of (x1 - y2, y1 - y2, z1 - z2)] MM[symmetric]
    Rep-riemann-sphere-inject[of x y]
    by (simp add: dist-riemann-sphere'-def) (smt prod.inject zero-prod-def)
  qed
next
  fix S :: riemann-sphere set
  show open S = ( $\forall x \in S. \exists e > 0. \forall y. dist\text{ }y\text{ }x < e \rightarrow y \in S$ )
  unfolding open-riemann-sphere-def dist-riemann-sphere-def
  by simp
next
  fix x y z :: riemann-sphere
  show dist x y ≤ dist x z + dist y z
  proof-
    obtain x1 y1 z1 x2 y2 z2 x3 y3 z3 where MM: (x1, y1, z1) = Rep-riemann-sphere
    x (x2, y2, z2) = Rep-riemann-sphere y (x3, y3, z3) = Rep-riemann-sphere z
    by (cases Rep-riemann-sphere x, cases Rep-riemann-sphere y, cases Rep-riemann-sphere z, auto, blast)
    show ?thesis
    unfolding dist-riemann-sphere-def
    using MM[symmetric] norm-minus-cancel[of (x3 - x2, y3 - y2, z3 - z2)]
    norm-triangle-ineq[of (x1 - x3, y1 - y3, z1 - z3) (x3 - x2, y3 - y2, z3 - z2)]
    by (simp add: dist-riemann-sphere'-def field-simps)
  qed
qed
end

lemma ex-cos-gt':
assumes a ≥ 0 a < 1 -pi/2 ≤ α ∧ α ≤ pi/2
shows ∃ α'. -pi/2 ≤ α' ∧ α' ≤ pi/2 ∧ α' ≠ α ∧ cos(α - α') = a

```

```

proof-
  have arccos a > 0 arccos a ≤ pi/2
    using ⟨a ≥ 0⟩ ⟨a < 1⟩
    using arccos-lt-bounded arccos-le-pi2
    by auto

  show ?thesis
  proof (cases α = arccos a ≥ -pi/2)
    case True
    thus ?thesis
      using assms ⟨arccos a > 0⟩ ⟨arccos a ≤ pi/2⟩
      by (rule-tac x = α - arccos a in exI) auto
  next
    case False
    thus ?thesis
      using assms ⟨arccos a > 0⟩ ⟨arccos a ≤ pi/2⟩
      by (rule-tac x = α + arccos a in exI) auto
  qed
qed

lemma ex-cos-gt:
  assumes a < 1 -pi/2 ≤ α ∧ α ≤ pi/2
  shows ∃ α'. -pi/2 ≤ α' ∧ α' ≤ pi/2 ∧ α' ≠ α ∧ cos(α - α') > a
proof-
  have ∃ a'. a' ≥ 0 ∧ a' > a ∧ a' < 1
    using ⟨a < 1⟩
    using divide-strict-right-mono[of 2*a + (1 - a) 2 2]
    by (rule-tac x=if a < 0 then 0 else a + (1-a)/2 in exI) (auto simp add:
  field-simps)
  then obtain a' where a' ≥ 0 a' > a a' < 1
    by auto
  thus ?thesis
    using ex-cos-gt'[of a' α] assms
    by auto
qed

instantiation riemann-sphere :: perfect-space
begin
instance proof
  fix M :: riemann-sphere
  obtain x y z where MM: Rep-riemann-sphere M = (x, y, z)
    by (cases Rep-riemann-sphere M) auto
  then obtain α β where *: x = cos α * cos β y = cos α * sin β z = sin α -pi
  / 2 ≤ α ∧ α ≤ pi / 2
    using Rep-riemann-sphere[of M]
    using ex-sphere-params[of x y z]
    by auto
  show ⊢ open {M}
    unfolding open-riemann-sphere-def

```

```

proof auto
fix e :: real
assume e > 0
then obtain α' where 1 - (e*e/2) < cos (α - α') α ≠ α' -pi/2 ≤ α' α' ≤
pi/2
  using ex-cos-gt[of 1 - (e*e/2) α] ← pi / 2 ≤ α ∧ α ≤ pi / 2
  by (auto simp add: mult-pos-pos)
hence sin α ≠ sin α'
  using ‹-pi / 2 ≤ α ∧ α ≤ pi / 2› sin-inj[of α α']
  by auto

have 2 - 2 * cos (α - α') < e*e
  using mult-strict-right-mono[OF ‹1 - (e*e/2) < cos (α - α')›, of 2]
  by (simp add: field-simps)
have 2 - 2 * cos (α - α') ≥ 0
  using cos-le-one[of α - α']
  by (simp add: sign-simps)
let ?M' = Abs-riemann-sphere (cos α' * cos β, cos α' * sin β, sin α')
  have dist-riemann-sphere' M ?M' = sqrt ((cos α - cos α')² + (sin α - sin
α')²)
    using MM * sphere-params-on-sphere[of - α' β]
    using sin-cos-squared-add[of β]
apply (simp add: dist-riemann-sphere'-def Abs-riemann-sphere-inverse norm-prod-def)
  apply (subst left-diff-distrib[symmetric])+
  apply (subst power-mult-distrib)+
  apply (subst distrib-left[symmetric])
  apply simp
done
also have ... = sqrt (2 - 2*cos (α - α'))
  by (simp add: power2-eq-square field-simps cos-diff)
finally
have (dist-riemann-sphere' M ?M')² = 2 - 2*cos (α - α')
  using ‹2 - 2 * cos (α - α') ≥ 0›
  by simp
hence (dist-riemann-sphere' M ?M')² < e²
  using ‹2 - 2 * cos (α - α') < e*e›
  by (simp add: power2-eq-square)
hence dist-riemann-sphere' M ?M' < e
  apply (rule power2-less-imp-less)
  using ‹e > 0›
  by simp
moreover
have M ≠ ?M'
  apply (subst Rep-riemann-sphere-inverse[symmetric])
  using Abs-riemann-sphere-inject[of Rep-riemann-sphere M (cos α' * cos β,
cos α' * sin β, sin α') ]
  using MM MM[symmetric] * sphere-params-on-sphere[of - α' β] Rep-riemann-sphere[of
M] ‹sin α ≠ sin α'›
  by (simp add: Abs-riemann-sphere-inverse)

```

```

ultimately
show  $\exists y. \text{dist-riemann-sphere}' y M < e \wedge y \neq M$ 
    by (rule-tac  $x=?M'$  in exI) (simp add: dist-riemann-sphere'-sym)
qed
qed
end

instantiation complex-homo :: perfect-space
begin
instance proof
    fix  $x:\text{complex-homo}$ 
    show  $\neg \text{open } \{x\}$ 
        unfolding open-complex-homo-def[ $\{x\}$ ]
        proof (auto)
            fix  $e:\text{real}$ 
            assume  $e > 0$ 
            thus  $\exists y. \text{dist-homo } y x < e \wedge y \neq x$ 
                using not-open-singleton[ $\{ \text{inv-stereographic } x \}$ ]
                unfolding open-riemann-sphere-def[ $\{ \text{inv-stereographic } x \}$ ]
                apply (subst dist-homo-stereographic', auto)
                apply (erule-tac  $x=e$  in allE, auto)
                apply (rule-tac  $x=\text{stereographic}$   $y$  in exI, auto simp add: inv-stereographic-stereographic)
                done
            qed
        qed
    end

lemma continuous-on UNIV stereographic
unfolding continuous-on-iff
unfolding dist-complex-homo-def dist-riemann-sphere-def
by (subst dist-homo-stereographic', auto simp add: inv-stereographic-stereographic)

lemma continuous-on UNIV inv-stereographic
unfolding continuous-on-iff
unfolding dist-complex-homo-def dist-riemann-sphere-def
by (subst dist-homo-stereographic) (auto simp add: stereographic-inv-stereographic)

end

```

10 Moebius transformations

```

theory Moebius
imports HomogeneousCoordinates
begin

typedef moebius-mat = { $M:\text{complex-mat}. \text{mat-det } M \neq 0$ }
by (rule-tac  $x=\text{eye}$  in exI, simp)

```

```

definition moebius-mat-eq where
  [simp]: moebius-mat-eq A B  $\longleftrightarrow$  ( $\exists$  k::complex. k  $\neq$  0  $\wedge$  Rep-moebius-mat B = k *sm (Rep-moebius-mat A))

lemma [simp]: moebius-mat-eq x x
  by (simp, rule-tac x=1 in exI, simp)

quotient-type moebius = moebius-mat / moebius-mat-eq
proof (rule equivpI)
  show reflp moebius-mat-eq
    by (auto simp add: reflp-def, rule-tac x=1 in exI, simp)
next
  show symp moebius-mat-eq
    by (auto simp add: symp-def, rule-tac x=1/k in exI, simp)
next
  show transp moebius-mat-eq
    by (auto simp add: transp-def, rule-tac x=ka*k in exI, simp)
qed

definition mk-moebius-rep where
  mk-moebius-rep a b c d = Abs-moebius-mat (a, b, c, d)

lift-definition mk-moebius :: complex  $\Rightarrow$  complex  $\Rightarrow$  complex  $\Rightarrow$  complex  $\Rightarrow$  moebius is mk-moebius-rep
  by (simp del: moebius-mat-eq-def)

lemma mk-moebius-rep-Rep:
  assumes mat-det (a, b, c, d)  $\neq$  0
  shows Rep-moebius-mat (mk-moebius-rep a b c d) = (a, b, c, d)
  using assms
  by (simp add: mk-moebius-rep-def Abs-moebius-mat-inverse)

lemma ex-mk-moebius:
  shows  $\exists$  a b c d. M = mk-moebius a b c d  $\wedge$  mat-det (a, b, c, d)  $\neq$  0
proof transfer
  fix M
  obtain a b c d where Rep-moebius-mat M = (a, b, c, d)
    by (cases Rep-moebius-mat M) auto
  hence moebius-mat-eq M (mk-moebius-rep a b c d)  $\wedge$  mat-det (a, b, c, d)  $\neq$  0
    using Rep-moebius-mat[of M]
    by (simp add: mk-moebius-rep-Rep, rule-tac x=1 in exI, simp)
    thus  $\exists$  a b c d. moebius-mat-eq M (mk-moebius-rep a b c d)  $\wedge$  mat-det (a, b, c, d)  $\neq$  0
      by blast
  qed

```

10.1 Action on points

definition *moebius-pt-rep* :: *moebius-mat* \Rightarrow *homo-coords* \Rightarrow *homo-coords* **where**

```
moebius-pt-rep M z =
  (let z = Rep-homo-coords z;
   M = Rep-moebius-mat M
   in Abs-homo-coords (M *mv z))
```

lemma [*simp*]: *Rep-homo-coords* (*Abs-homo-coords* (*Rep-moebius-mat* $M *_{mv}$ *Rep-homo-coords* x)) = *Rep-moebius-mat* $M *_{mv}$ *Rep-homo-coords* x
using *Rep-moebius-mat*[of *M*] *Rep-homo-coords*[of *x*] *mult-mv-nonzero*[of *Rep-homo-coords* x *Rep-moebius-mat* M]
by (*simp add: Abs-homo-coords-inverse*)

lemma [*simp*]: *Rep-homo-coords* (*moebius-pt-rep* $M z$) = *Rep-moebius-mat* $M *_{mv}$ *Rep-homo-coords* z
by (*simp add: moebius-pt-rep-def*)

lift-definition *moebius-pt* :: *moebius* \Rightarrow *complex-homo* \Rightarrow *complex-homo* **is** *moebius-pt-rep*
proof -
fix *M M' x x'*
assume *moebius-mat-eq* $M M' x \approx x'$
thus *moebius-pt-rep* $M x \approx moebius-pt-rep M' x'$
by (*cases Rep-moebius-mat M, cases Rep-homo-coords x, auto simp add: field-simps*) (*rule-tac x=k*ka in exI, simp*)
qed

lemma *bij-moebius-pt*:
shows *bij* (*moebius-pt* *M*)
unfolding *bij-def inj-on-def surj-def*
proof (*simp, transfer, safe*)
fix *M x y*
assume *moebius-pt-rep* $M x \approx moebius-pt-rep M y$
thus $x \approx y$
using *Rep-moebius-mat*[of *M*]
apply *auto*
apply (*subst (asm) mult-sv-mv*)
using *mult-mv-cancel-l*
by *blast*
next
fix *M y*
let *?M = Rep-moebius-mat M*
let *?iM = mat-inv ?M*
let *?y = Rep-homo-coords y*
show $\exists x. y \approx moebius-pt-rep M x$
using *Rep-moebius-mat*[of *M*] *mat-det-inv*[of *?M*] *Rep-homo-coords*[of *y*] *mult-mv-nonzero*[of *?y ?iM*]
using *mat-inv-r*[of *?M*] *eye-mv-l*[of *?y*]
by (*auto, rule-tac x=Abs-homo-coords ((mat-inv (Rep-moebius-mat M)) *_{mv}*

```

Rep-homo-coords y) in exI, rule-tac x=1 in exI)
  (auto simp add: Abs-homo-coords-inverse)
qed

definition is-moebius where
  is-moebius f  $\longleftrightarrow$  ( $\exists M$ . f = moebius-pt M)

Bilinear and linear expressions

lemma moebius-bilinear:
  assumes mat-det (a, b, c, d)  $\neq 0$ 
  shows moebius-pt (mk-moebius a b c d) z =
    (if z  $\neq \infty_h$  then
      ((of-complex a) *h z +h (of-complex b)) :h
      ((of-complex c) *h z +h (of-complex d)))
    else
      (of-complex a) :h
      (of-complex c))
unfolding divide-homo-def
using assms
proof (transfer)
  fix a b c d :: complex and z
  obtain z1 z2 where zz: Rep-homo-coords z = (z1, z2)
    by (rule obtain-homo-coords)
  assume *: mat-det (a, b, c, d)  $\neq 0$ 
  let ?oc = of-complex-coords
  show moebius-pt-rep (mk-moebius-rep a b c d) z  $\approx$ 
    (if  $\neg z \approx \text{inf-homo-rep}$ 
      then ?oc a *hc z +hc ?oc b *hc
        reciprocal-homo-coords (?oc c *hc z +hc ?oc d)
      else ?oc a *hc
        reciprocal-homo-coords (of-complex-coords c))
  proof (cases z  $\approx$  inf-homo-rep)
    case True
    thus ?thesis
      using zz *
      using mult-homo-coords-Rep[of ?oc a a 1 reciprocal-homo-coords (?oc c) 1 c]
      using reciprocal-homo-coords-Rep[of ?oc c]
      by (force simp add: mk-moebius-rep-Rep field-simps)
    next
      case False
      hence z2  $\neq 0$ 
        using zz Rep-homo-coords[of z]
      by auto (metis mult.commute complex-divide-def mult-zero-right right-inverse-eq)
      thus ?thesis
        using zz * False
        using regular-homogenous-system[of a d b c z1 z2]
        apply simp
        apply (subst mult-homo-coords-Rep[of ?oc a *hc z +hc ?oc b a*z1+b*z2 z2
          reciprocal-homo-coords (?oc c *hc z +hc ?oc d) z2 c*z1+d*z2])

```

```

using add-homo-coords-Rep[of ?oc a *hc z a*z1 z2 ?oc b b 1]
using mult-homo-coords-Rep[of ?oc a a 1 z z1 z2]
using reciprocal-homo-coords-Rep[of ?oc c *hc z +hc ?oc d]
using add-homo-coords-Rep[of ?oc c *hc z c*z1 z2 ?oc d d 1]
using mult-homo-coords-Rep[of ?oc c c 1 z z1 z2]
by (auto simp add: mk-moebius-rep-Rep)
qed
qed

```

10.2 Moebius group

definition moebius-inv-rep **where**

```

moebius-inv-rep M =
  (let M = Rep-moebius-mat M
    in Abs-moebius-mat (mat-inv M))

```

lemma [simp]: Rep-moebius-mat (Abs-moebius-mat (mat-inv (Rep-moebius-mat M))) = mat-inv (Rep-moebius-mat M)
using Rep-moebius-mat[of M] mat-det-inv[of Rep-moebius-mat M]
by (auto simp add: Abs-moebius-mat-inverse)

lemma [simp]: Rep-moebius-mat (moebius-inv-rep M) = mat-inv (Rep-moebius-mat M)
by (simp add: moebius-inv-rep-def)

lift-definition moebius-inv :: moebius ⇒ moebius **is** moebius-inv-rep
proof –

```

fix x y
assume moebius-mat-eq x y
thus moebius-mat-eq (moebius-inv-rep x) (moebius-inv-rep y)
  by (auto simp add: mat-inv-mult-sm) (rule-tac x=1/k in exI, simp)
qed

```

lemma moebius-inv: moebius-pt (moebius-inv M) = inv (moebius-pt M)
proof (rule inv-equality[symmetric])

```

fix x
show moebius-pt (moebius-inv M) (moebius-pt M x) = x
proof (transfer)
  fix M x
  show moebius-pt-rep (moebius-inv-rep M) (moebius-pt-rep M x) ≈ x
    using Rep-moebius-mat[of M] Rep-homo-coords[of x] eye-mv-l
    by (simp add: mat-inv-l) (rule-tac x=1 in exI, simp)
qed

```

next

```

fix y
show moebius-pt M (moebius-pt (moebius-inv M) y) = y
proof (transfer)
  fix M y
  show moebius-pt-rep M (moebius-pt-rep (moebius-inv-rep M) y) ≈ y

```

```

using Rep-moebius-mat[of M] eye-mv-l
by (simp add: mat-inv-r) (rule-tac x=1 in exI, simp)
qed
qed

lemma is-moebius-inv:
assumes is-moebius m
shows is-moebius (inv m)
using assms
unfolding is-moebius-def
using moebius-inv[symmetric]
by auto

definition moebius-comp-rep where
moebius-comp-rep M1 M2 =
(let M1 = Rep-moebius-mat M1;
M2 = Rep-moebius-mat M2 in
Abs-moebius-mat (M1 *mm M2))

lemma [simp]: Rep-moebius-mat (Abs-moebius-mat ((Rep-moebius-mat M1) *mm
(Rep-moebius-mat M2))) = (Rep-moebius-mat M1) *mm (Rep-moebius-mat M2)
using Rep-moebius-mat[of M1] Rep-moebius-mat[of M2]
by (simp add: Abs-moebius-mat-inverse)

lemma [simp]: Rep-moebius-mat (moebius-comp-rep M1 M2) = (Rep-moebius-mat
M1) *mm (Rep-moebius-mat M2)
by (simp add: moebius-comp-rep-def)

lift-definition moebius-comp :: moebius ⇒ moebius ⇒ moebius is moebius-comp-rep
by auto (rule-tac x=ka*k in exI, simp)

lemma moebius-comp: moebius-pt M1 ∘ moebius-pt M2 = moebius-pt (moebius-comp
M1 M2)
unfolding comp-def
by (rule ext, transfer) (simp, rule-tac x=1 in exI, simp)

lemma is-moebius-comp:
assumes is-moebius m1 is-moebius m2
shows is-moebius (m1 ∘ m2)
using assms
unfolding is-moebius-def
using moebius-comp
by auto

definition [simp]: id-moebius-rep = Abs-moebius-mat eye

lift-definition id-moebius :: moebius is id-moebius-rep
done

```

```

lemma [simp]: Rep-moebius-mat (Abs-moebius-mat (1, 0, 0, 1)) = eye
by (simp add: Abs-moebius-mat-inverse)

lemma [simp]: Rep-moebius-mat (id-moebius-rep) = eye
by simp

lemma moebius-pt id-moebius = id
unfolding id-def
apply (rule ext, transfer)
using eye-mv-l
by simp (rule-tac x=1 in exI, simp)

instantiation moebius :: group-add
begin
definition plus-moebius :: moebius ⇒ moebius ⇒ moebius where
  [simp]: plus-moebius = moebius-comp

definition uminus-moebius :: moebius ⇒ moebius where
  [simp]: uminus-moebius = moebius-inv

definition zero-moebius :: moebius where
  [simp]: zero-moebius = id-moebius

definition minus-moebius :: moebius ⇒ moebius ⇒ moebius where
  [simp]: minus-moebius A B = A + (-B)

instance proof
  fix a b c :: moebius
  show a + b + c = a + (b + c)
    unfolding plus-moebius-def
    proof (transfer)
      fix a b c
      show moebius-mat-eq (moebius-comp-rep (moebius-comp-rep a b) c) (moebius-comp-rep
        a (moebius-comp-rep b c))
        using Rep-moebius-mat[of a] Rep-moebius-mat[of b] Rep-moebius-mat[of c]
        by simp (rule-tac x=1 in exI, simp add: mult-mm-assoc)
    qed
  next
    fix a :: moebius
    show a + 0 = a
      unfolding plus-moebius-def zero-moebius-def
      proof (transfer)
        fix A
        show moebius-mat-eq (moebius-comp-rep A id-moebius-rep) A
          using mat-eye-r
          by simp (rule-tac x=1 in exI, simp)
    qed
  next

```

```

fix a :: moebius
show 0 + a = a
  unfolding plus-moebius-def zero-moebius-def
proof (transfer)
  fix A
  show moebius-mat-eq (moebius-comp-rep id-moebius-rep A) A
    using mat-eye-l
    by simp (rule-tac x=1 in exI, simp)
qed
next
  fix a :: moebius
  show - a + a = 0
    unfolding plus-moebius-def uminus-moebius-def zero-moebius-def
proof (transfer)
  fix a
  show moebius-mat-eq (moebius-comp-rep (moebius-inv-rep a) a) id-moebius-rep
    using Rep-moebius-mat[of a]
    by (simp add: mat-inv-l)
qed
next
  fix a b :: moebius
  show a - b = a + - b
    unfolding minus-moebius-def
    by simp
qed
end

lemma [simp]: moebius-comp (moebius-inv M) M = id-moebius
by (metis left-minus plus-moebius-def uminus-moebius-def zero-moebius-def)

lemma [simp]: moebius-comp M (moebius-inv M) = id-moebius
by (metis right-minus plus-moebius-def uminus-moebius-def zero-moebius-def)

lemma moebius-pt-moebius-id [simp]: moebius-pt (id-moebius) = id
by (rule ext) (transfer, case-tac Rep-homo-coords x, auto, rule-tac x=1 in exI,
  simp)

lemma [simp]: moebius-pt (moebius-inv M) (moebius-pt M z) = z
proof-
  have moebius-pt (moebius-inv M) (moebius-pt M z) = (moebius-pt (moebius-inv
  M) o moebius-pt M) z
    by simp
  thus ?thesis
    using moebius-comp[of moebius-inv M M]
    by simp
qed

lemma moebius-pt-invert:
assumes w = moebius-pt M z

```

```

shows  $z = \text{moebius-pt}(\text{moebius-inv } M) w$ 
using assms
by auto

```

10.3 Special kinds of Moebius transformations

Reciprocal ($1/z$) as a moebius transformation

```

definition reciprocal-moebius :: moebius where
  reciprocal-moebius = mk-moebius 0 1 1 0

```

```

lemma [simp]: Rep-moebius-mat (Abs-moebius-mat (0, 1, 1, 0)) = (0, 1, 1, 0)
by (simp add: Abs-moebius-mat-inverse)

```

```

lemma [simp]: Rep-moebius-mat (mk-moebius-rep 0 1 1 0) = (0, 1, 1, 0)
by (simp add: mk-moebius-rep-def)

```

```

lemma [simp]: Rep-homo-coords (reciprocal-homo-coords z) = (let (x, y) = Rep-homo-coords
z in (y, x))
unfolding reciprocal-homo-coords-def Let-def
apply (cases Rep-homo-coords z)
using Rep-homo-coords[of z]
by (auto simp add: Abs-homo-coords-inverse)

```

```

lemma reciprocal-moebius:
  reciprocal-homo = moebius-pt reciprocal-moebius
unfolding reciprocal-moebius-def
by (rule ext, transfer) (auto simp add: split-def Let-def, case-tac Rep-homo-coords
x, rule-tac x=1 in exI, auto)

```

```

lemma reciprocal-moebius-inv [simp]:
  moebius-inv reciprocal-moebius = reciprocal-moebius
unfolding reciprocal-moebius-def
by transfer simp

```

```

lemma reciprocal-homo-only-0-to-inf:
  assumes reciprocal-homo  $z = \infty_h$ 
  shows  $z = 0_h$ 
using assms
unfolding reciprocal-moebius
using moebius-pt-invert[of  $\infty_h$  reciprocal-moebius z]
by (simp add: reciprocal-moebius[symmetric])

```

```

lemma reciprocal-homo-only-inf-to-0:
  assumes reciprocal-homo  $z = 0_h$ 
  shows  $z = \infty_h$ 
using assms
unfolding reciprocal-moebius
using moebius-pt-invert[of  $0_h$  reciprocal-moebius z]
by (simp add: reciprocal-moebius[symmetric])

```

Euclidean similarity as a Moebius transform

```

definition similarity-moebius :: complex ⇒ complex ⇒ moebius where
  similarity-moebius a b = mk-moebius a b 0 1

lemma moebius-similarity-linear:
  assumes a ≠ 0
  shows moebius-pt (similarity-moebius a b) z = (of-complex a) *h z +h (of-complex
  b)
  unfolding similarity-moebius-def
  using assms
  using mult-homo-inf-right[of of-complex a]
  by (subst moebius-bilinear, auto)

lemma moebius-similarity':
  assumes a ≠ 0
  shows moebius-pt (similarity-moebius a b) = (λ z. (of-complex a) *h z +h
  (of-complex b))
  using moebius-similarity-linear[OF assms, symmetric]
  by simp

lemma is-moebius-similarity':
  assumes a ≠ 0h a ≠ ∞h b ≠ ∞h
  shows (λ z. a *h z +h b) = moebius-pt (similarity-moebius (to-complex a)
  (to-complex b))
  proof-
    obtain ka kb where *: a = of-complex ka ka ≠ 0 b = of-complex kb
    using assms
    using inf-homo-or-complex-homo[of a] inf-homo-or-complex-homo[of b]
    by auto
    thus ?thesis
      unfolding is-moebius-def
      using moebius-similarity'[of ka kb]
      by simp
  qed

lemma is-moebius-similarity:
  assumes a ≠ 0h a ≠ ∞h b ≠ ∞h
  shows is-moebius (λ z. a *h z +h b)
  using is-moebius-similarity'[OF assms]
  unfolding is-moebius-def
  by auto

lemma similarity-moebius-comp:
  assumes a ≠ 0 c ≠ 0
  shows similarity-moebius a b + similarity-moebius c d = similarity-moebius
  (a*c) (a*d+b)
  using assms
  unfolding similarity-moebius-def plus-moebius-def
  by transfer (simp add: mk-moebius-rep-def Abs-moebius-mat-inverse)

```

```

lemma similarity-moebius-inv:
  assumes a ≠ 0
  shows – similarity-moebius a b = similarity-moebius (1/a) (–b/a)
  using assms
  unfolding similarity-moebius-def uminus-moebius-def
  by transfer (simp add: mk-moebius-rep-def Abs-moebius-mat-inverse)

lemma similarity-moebius-id: id-moebius = similarity-moebius 1 0
unfolding similarity-moebius-def
by transfer (simp add: mk-moebius-rep-def)

lemma similarity-inf-fixed:
  assumes a ≠ 0
  shows moebius-pt (similarity-moebius a b) ∞h = ∞h
  using assms
  unfolding similarity-moebius-def
  by transfer (simp add: mk-moebius-rep-def Abs-moebius-mat-inverse)

lemma similarity-only-inf-to-inf:
  assumes a ≠ 0 moebius-pt (similarity-moebius a b) z = ∞h
  shows z = ∞h
  using assms moebius-pt-invert[of ∞h similarity-moebius a b z] similarity-inf-fixed[of
  1/a –b/a]
  using similarity-moebius-inv[of a b]
  by simp

lemma inf-fixed-similarity:
  assumes moebius-pt M ∞h = ∞h
  shows ∃ a b. a ≠ 0 ∧ M = similarity-moebius a b
  using assms
  unfolding similarity-moebius-def
  proof transfer
    fix M
    obtain a b c d where MM: Rep-moebius-mat M = (a, b, c, d)
      by (cases M) (auto simp add: Abs-moebius-mat-inverse)
    assume moebius-pt-rep M inf-homo-rep ≈ inf-homo-rep
    hence c = 0
      using MM
      by auto
    hence *: a ≠ 0 ∧ d ≠ 0
      using Rep-moebius-mat[of M] MM
      by auto
    show ∃ a b. a ≠ 0 ∧ moebius-mat-eq M (mk-moebius-rep a b 0 1)
      proof (rule-tac x=a/d in exI, rule-tac x=b/d in exI)
        show a/d ≠ 0 ∧ moebius-mat-eq M (mk-moebius-rep (a / d) (b / d) 0 1)
          using MM ‹c = 0› ‹a ≠ 0 ∧ d ≠ 0›
        by simp (rule-tac x=1/d in exI, simp add: mk-moebius-rep-def Abs-moebius-mat-inverse)
      qed

```

qed

Translation

definition *translation-moebius* **where**

translation-moebius v = similarity-moebius 1 v

lemma *translation-moebius-comp*:

$(\text{translation-moebius } v1) + (\text{translation-moebius } v2) = \text{translation-moebius } (v1 + v2)$

unfolding *translation-moebius-def similarity-moebius-def plus-moebius-def*
by (*transfer*) (*auto simp add: mk-moebius-rep-Rep*)

lemma *translation-moebius-zero*:

translation-moebius 0 = id-moebius

unfolding *translation-moebius-def similarity-moebius-def*
by (*transfer*) (*auto simp add: mk-moebius-rep-Rep*)

lemma *moebius-translation-inv*:

$-(\text{translation-moebius } v1) = \text{translation-moebius } (-v1)$

using *translation-moebius-comp[of v1 -v1] translation-moebius-zero uminus-moebius-def*
using *equals-zero-I[of translation-moebius v1 translation-moebius (-v1)]*

by *simp*

lemma *moebius-pt-translation [simp]: moebius-pt (translation-moebius v) (of-complex z) = of-complex (v + z)*

unfolding *translation-moebius-def similarity-moebius-def*
by *transfer (simp add: mk-moebius-rep-Rep)*

Rotation

definition *rotation-moebius* **where**

rotation-moebius φ = similarity-moebius (cis φ) 0

lemma *rotation-moebius-comp*:

$(\text{rotation-moebius } \varphi1) + (\text{rotation-moebius } \varphi2) = \text{rotation-moebius } (\varphi1 + \varphi2)$

unfolding *rotation-moebius-def similarity-moebius-def plus-moebius-def*
by (*transfer (simp add: mk-moebius-rep-Rep cis-mult)*)

lemma *rotation-moebius-zero*:

rotation-moebius 0 = id-moebius

unfolding *rotation-moebius-def similarity-moebius-def*
by (*transfer (simp add: mk-moebius-rep-Rep)*)

lemma *rotation-moebius-inverse*:

$-(\text{rotation-moebius } \varphi) = \text{rotation-moebius } (-\varphi)$

using *rotation-moebius-comp[of φ -φ] rotation-moebius-zero*

using *equals-zero-I[of rotation-moebius φ rotation-moebius (-φ)]*

by *simp*

lemma *moebius-pt-rotation [simp]: moebius-pt (rotation-moebius φ) (of-complex*

$z) = \text{of-complex} (\text{cis } \varphi * z)$
unfolding *rotation-moebius-def similarity-moebius-def*
by transfer (*simp add: mk-moebius-rep-Rep*)

Dilatation

definition *dilatation-moebius* **where**
dilatation-moebius a = similarity-moebius (cor a) 0

lemma *dilatation-moebius-comp*:
assumes $a_1 > 0$ $a_2 > 0$
shows $(\text{dilatation-moebius } a_1) + (\text{dilatation-moebius } a_2) = \text{dilatation-moebius}$
 $(a_1 * a_2)$
using assms
unfolding *dilatation-moebius-def similarity-moebius-def plus-moebius-def*
by transfer (*simp add: mk-moebius-rep-def Abs-moebius-mat-inverse*)

lemma *dilatation-moebius-zero*:
dilatation-moebius 1 = id-moebius
unfolding *dilatation-moebius-def similarity-moebius-def*
by transfer (*simp add: mk-moebius-rep-Rep*)

lemma *dilatation-moebius-inverse*:
assumes $a > 0$
shows $-(\text{dilatation-moebius } a) = \text{dilatation-moebius } (1/a)$
using assms
using *dilatation-moebius-comp[of a 1/a] dilatation-moebius-zero*
using *equals-zero-I[of dilatation-moebius a dilatation-moebius (1/a)]*
by simp

lemma *moebius-pt-dilatation* [*simp*]: $a \neq 0 \implies \text{moebius-pt } (\text{dilatation-moebius } a)$
 $(\text{of-complex } z) = \text{of-complex } (\text{cor } a * z)$
unfolding *dilatation-moebius-def similarity-moebius-def*
by transfer (*simp add: mk-moebius-rep-Rep*)

rotation-dilation-moebius

definition *rotation-dilatation-moebius* **where**
rotation-dilatation-moebius a = similarity-moebius a 0

lemma *rot-dil*:
assumes $a \neq 0$
shows $\text{rotation-dilatation-moebius } a = \text{rotation-moebius } (\arg a) + \text{dilatation-moebius}$
 $(\text{cmod } a)$
using assms
unfolding *rotation-dilatation-moebius-def rotation-moebius-def dilatation-moebius-def*
similarity-moebius-def plus-moebius-def
by transfer (*simp add: mk-moebius-rep-Rep*)

10.4 Decomposition

```

lemma similarity-decomposition:
  assumes a ≠ 0
  shows similarity-moebius a b = (translation-moebius b) + (rotation-moebius (arg
  a)) + (dilatation-moebius (cmod a))
proof-
  have similarity-moebius a b = (translation-moebius b) + rotation-dilatation-moebius
  a
    unfolding rotation-dilatation-moebius-def translation-moebius-def similarity-moebius-def
    plus-moebius-def
    using assms
    by transfer (simp add: mk-moebius-rep-Rep)
    thus ?thesis
      using rot-dil[OF assms]
      by (auto simp add: add-assoc simp del: plus-moebius-def)
  qed

lemma moebius-decomposition:
  assumes c ≠ 0 a*d - b*c ≠ 0
  shows mk-moebius a b c d =
    translation-moebius (a/c) +
    rotation-dilatation-moebius ((b*c - a*d)/(c*c)) +
    reciprocal-moebius +
    translation-moebius (d/c)
  using assms
  unfolding rotation-dilatation-moebius-def translation-moebius-def similarity-moebius-def
  plus-moebius-def reciprocal-moebius-def
  by transfer (simp add: mk-moebius-rep-Rep, rule-tac x=1/c in exI, simp add:
  field-simps)

lemma wlog-moebius-decomposition:
  assumes
    trans:  $\bigwedge v. P(\text{translation-moebius } v)$  and rot:  $\bigwedge \alpha. P(\text{rotation-moebius } \alpha)$  and
    dil:  $\bigwedge k. P(\text{dilatation-moebius } k)$  and recip:  $P(\text{reciprocal-moebius})$  and
    comp:  $\bigwedge M1 M2. [P M1; P M2] \implies P(M1 + M2)$ 
  shows P M
proof-
  obtain a b c d where M = mk-moebius a b c d mat-det (a, b, c, d) ≠ 0
  using ex-mk-moebius[of M]
  by auto
  show ?thesis
  proof (cases c = 0)
    case False
    show ?thesis
      using moebius-decomposition[of c a d b] (mat-det (a, b, c, d) ≠ 0) (c ≠ 0)
      M = mk-moebius a b c d
      using rot-dil[of (b*c - a*d) / (c*c)]
      using trans[of a/c] rot[of arg ((b*c - a*d) / (c*c))] dil[of cmod ((b*c -
      a*d) / (c*c))] recip

```

```

using comp
by simp (metis trans)
next
case True
hence M = similarity-moebius (a/d) (b/d)
  using ⟨M = mk-moebius a b c d⟩ ⟨mat-det (a, b, c, d) ≠ 0⟩
  unfolding similarity-moebius-def
    by transfer (auto simp add: mk-moebius-rep-Rep, rule-tac x=k/d in exI,
      case-tac Rep-moebius-mat M, simp)
  thus ?thesis
    using ⟨c = 0⟩ ⟨mat-det (a, b, c, d) ≠ 0⟩
    using similarity-decomposition[of a/d b/d]
    using trans[of b/d] rot[of arg (a/d)] dil[of cmod (a/d)] comp
    by simp
qed
qed

```

10.5 Cross ratio and moebius existence

```

lemma is-moebius-cross-ratio:
  assumes z1 ≠ z2 z2 ≠ z3 z1 ≠ z3
  shows is-moebius (λ z. cross-ratio z z1 z2 z3)
proof –
  have ∃ M. ∀ z. cross-ratio z z1 z2 z3 = moebius-pt M z
    using assms
  proof (transfer)
    fix z1 z2 z3
    obtain z1' z1'' where zz1: Rep-homo-coords z1 = (z1', z1'')
      by (rule obtain-homo-coords)
    obtain z2' z2'' where zz2: Rep-homo-coords z2 = (z2', z2'')
      by (rule obtain-homo-coords)
    obtain z3' z3'' where zz3: Rep-homo-coords z3 = (z3', z3'')
      by (rule obtain-homo-coords)

    let ?m23 = z2'*z3'' - z3'*z2''
    let ?m21 = z2'*z1'' - z1'*z2''
    let ?m13 = z1'*z3'' - z3'*z1''
    let ?M = (z1''*?m23, -z1'*?m23, z3''*?m21, -z3'*?m21)
    assume ¬ z1 ≈ z2 ∨ z2 ≈ z3 ∨ z1 ≈ z3
    hence *: ?m23 ≠ 0 ∨ ?m21 ≠ 0 ∨ ?m13 ≠ 0
      using zz1 zz2 zz3
      using homo-coords-eq-mix[of z1 z1' z1'' z2 z2' z2''] homo-coords-eq-mix[of z1' z1'' z3 z3' z3''] homo-coords-eq-mix[of z2 z2' z2'' z3 z3' z3''] by auto
    have mat-det ?M = ?m21*?m23*?m13
      by (simp add: field-simps)
    hence mat-det ?M ≠ 0
      using *
      by simp

```

```

show  $\exists M. \forall z. \text{cross-ratio-rep } z z1 z2 z3 \approx \text{moebius-pt-rep } M z$ 
proof (rule-tac x=Abs-moebius-mat ?M in exI, rule)
  fix z
  obtain z' z'' where zz: Rep-homo-coords z = (z', z'')
    by (rule obtain-homo-coords)

  let ?m01 = z'*z1'' - z1'*z''
  let ?m03 = z'*z3'' - z3'*z''

  have ?m01 ≠ 0 ∨ ?m03 ≠ 0
    using * Rep-homo-coords[of z] zz
    apply (cases z'' = 0 ∨ z1'' = 0 ∨ z3'' = 0)
    apply (auto simp add: field-simps)
    apply (subgoal-tac z1'/z1'' = z3'/z3'')
    by (simp add: field-simps) (metis eq-divide-imp mult-divide-mult-cancel-left
times-divide-eq-right times-divide-times-eq)
  note * = * this

  show cross-ratio-rep z z1 z2 z3 ≈ moebius-pt-rep (Abs-moebius-mat ?M) z
    using zz1 zz2 zz3 zz * Rep-homo-coords[of z] mult-mv-nonzero[of Rep-homo-coords
z ?M] (mat-det ?M ≠ 0)
    by (simp add: cross-ratio-rep-def moebius-pt-rep-def split-def Let-def Abs-moebius-mat-inverse
Abs-homo-coords-inverse)
      (rule-tac x=1 in exI, simp add: field-simps)
  qed
  qed
  thus ?thesis
    by (auto simp add: is-moebius-def)
  qed

lemma ex-moebius-01inf:
  assumes z1 ≠ z2 z1 ≠ z3 z2 ≠ z3
  shows  $\exists M. ((\text{moebius-pt } M z1 = 0_h) \wedge (\text{moebius-pt } M z2 = 1_h) \wedge (\text{moebius-pt } M z3 = \infty_h))$ 
  using assms
  using is-moebius-cross-ratio[OF ⟨z1 ≠ z2⟩ ⟨z2 ≠ z3⟩ ⟨z1 ≠ z3⟩]
  using cross-ratio-0[OF ⟨z1 ≠ z2⟩ ⟨z1 ≠ z3⟩] cross-ratio-1[OF ⟨z1 ≠ z2⟩ ⟨z2 ≠ z3⟩] cross-ratio-inf[OF ⟨z1 ≠ z3⟩ ⟨z2 ≠ z3⟩]
  by (auto simp add: is-moebius-def) metis

lemma ex-moebius:
  assumes z1 ≠ z2 z1 ≠ z3 z2 ≠ z3 w1 ≠ w2 w1 ≠ w3 w2 ≠ w3
  shows  $\exists M. ((\text{moebius-pt } M z1 = w1) \wedge (\text{moebius-pt } M z2 = w2) \wedge (\text{moebius-pt } M z3 = w3))$ 
  proof-
    obtain M1 where *: moebius-pt M1 z1 = 0_h ∧ moebius-pt M1 z2 = 1_h ∧
moebius-pt M1 z3 = ∞_h
    using ex-moebius-01inf[OF assms(1-3)]

```

```

by auto
obtain M2 where **: moebius-pt M2 w1 = 0_h ∧ moebius-pt M2 w2 = 1_h ∧
moebius-pt M2 w3 = ∞_h
  using ex-moebius-01inf[OF assms(4–6)]
  by auto
let ?M = moebius-comp (moebius-inv M2) M1
show ?thesis
  using * ** bij-moebius-pt[of M2]
  by (rule-tac x=?M in exI, (subst moebius-comp[symmetric])+, (subst moebius-inv)+)
(simp add: bij-def inv-f-eq)
qed

lemma ex-moebius-1:
  shows ∃ M. moebius-pt M z1 = w1
proof –
  obtain z2 z3 where z1 ≠ z2 z1 ≠ z3 z2 ≠ z3
    using ex-3-different-points[of z1]
    by auto
  moreover
  obtain w2 w3 where w1 ≠ w2 w1 ≠ w3 w2 ≠ w3
    using ex-3-different-points[of w1]
    by auto
  ultimately
  show ?thesis
    using ex-moebius[of z1 z2 z3 w1 w2 w3]
    by auto
qed

lemma wlog-moebius-01inf:
  fixes M::moebius
  assumes P 0_h 1_h ∞_h z1 ≠ z2 z2 ≠ z3 z1 ≠ z3
    ∧ M a b c. P a b c ⟹ P (moebius-pt M a) (moebius-pt M b) (moebius-pt M c)
  shows P z1 z2 z3
proof –
  from assms obtain M where *:
    moebius-pt M z1 = 0_h moebius-pt M z2 = 1_h moebius-pt M z3 = ∞_h
    using ex-moebius-01inf[of z1 z2 z3]
    by auto
  have **: moebius-pt (moebius-inv M) 0_h = z1 moebius-pt (moebius-inv M) 1_h
    = z2 moebius-pt (moebius-inv M) ∞_h = z3
    by (subst *[symmetric], simp) +
  thus ?thesis
    using assms
    by auto
qed

```

10.6 Fixed points and moebius uniqueness

lemma three-fixed-points-01inf:

```

assumes moebius-pt M 0h = 0h moebius-pt M 1h = 1h moebius-pt M ∞h =
∞h
shows M = id-moebius
using assms
by transfer (case-tac Rep-moebius-mat M, auto)

```

lemma three-fixed-points:

```

assumes z1 ≠ z2 z1 ≠ z3 z2 ≠ z3
assumes moebius-pt M z1 = z1 moebius-pt M z2 = z2 moebius-pt M z3 = z3
shows M = id-moebius

```

proof –

```

from assms obtain M' where *: moebius-pt M' z1 = 0h moebius-pt M' z2 =
1h moebius-pt M' z3 = ∞h

```

```

using ex-moebius-01inf[of z1 z2 z3]
by auto

```

```

have **: moebius-pt (moebius-inv M') 0h = z1 moebius-pt (moebius-inv M') 1h
= z2 moebius-pt (moebius-inv M') ∞h = z3

```

```

by (subst *[symmetric], simp) +

```

```

have M' + M + (-M') = 0

```

```

unfoldng zero-moebius-def

```

```

apply (rule three-fixed-points-01inf)

```

```

using * ** assms

```

```

by (simp add: moebius-comp[symmetric]) +

```

```

thus ?thesis

```

```

by (metis eq-neg-iff-add-eq-0 minus-add-cancel zero-moebius-def)

```

qed

lemma unique-moebius-three-points:

```

assumes z1 ≠ z2 z1 ≠ z3 z2 ≠ z3

```

```

assumes moebius-pt M1 z1 = w1 moebius-pt M1 z2 = w2 moebius-pt M1 z3 =
w3

```

```

moebius-pt M2 z1 = w1 moebius-pt M2 z2 = w2 moebius-pt M2 z3 = w3

```

```

shows M1 = M2

```

proof –

```

let ?M = moebius-comp (moebius-inv M2) M1

```

```

have moebius-pt ?M z1 = z1

```

```

using ⟨moebius-pt M1 z1 = w1⟩ ⟨moebius-pt M2 z1 = w1⟩

```

```

using bij-moebius-pt[of M2]

```

```

by (subst moebius-comp[symmetric], subst moebius-inv, simp add: bij-def inv-f-eq)

```

moreover

```

have moebius-pt ?M z2 = z2

```

```

using ⟨moebius-pt M1 z2 = w2⟩ ⟨moebius-pt M2 z2 = w2⟩

```

```

using bij-moebius-pt[of M2]

```

```

by (subst moebius-comp[symmetric], subst moebius-inv, simp add: bij-def inv-f-eq)

```

moreover

```

have moebius-pt ?M z3 = z3

```

```

using ⟨moebius-pt M1 z3 = w3⟩ ⟨moebius-pt M2 z3 = w3⟩

```

```

using bij-moebius-pt[of M2]

```

```

by (subst moebius-comp[symmetric], subst moebius-inv, simp add: bij-def inv-f-eq)
ultimately
have ?M = id-moebius
  using assms three-fixed-points
  by auto
thus ?thesis
  by (metis add-minus-cancel left-minus plus-moebius-def uminus-moebius-def
zero-moebius-def)
qed

lemma ex-unique-moebius-three-points:
assumes z1 ≠ z2 z1 ≠ z3 z2 ≠ z3 w1 ≠ w2 w1 ≠ w3 w2 ≠ w3
shows ∃! M. ((moebius-pt M z1 = w1) ∧ (moebius-pt M z2 = w2) ∧ (moebius-pt
M z3 = w3))
proof-
  obtain M where *: moebius-pt M z1 = w1 ∧ moebius-pt M z2 = w2 ∧ moebius-pt
M z3 = w3
    using ex-moebius[OF assms]
    by auto
  show ?thesis
    unfolding Ex1-def
    proof (rule-tac x=M in exI, rule)
      show ∀ y. moebius-pt y z1 = w1 ∧ moebius-pt y z2 = w2 ∧ moebius-pt y z3 =
w3 → y = M
        using *
        using unique-moebius-three-points[OF assms(1-3)]
        by simp
    qed (simp add: *)
  qed
lemma ex-unique-moebius-three-points-fun:
assumes z1 ≠ z2 z1 ≠ z3 z2 ≠ z3 w1 ≠ w2 w1 ≠ w3 w2 ≠ w3
shows ∃! f. is-moebius f ∧ (f z1 = w1) ∧ (f z2 = w2) ∧ (f z3 = w3)
proof-
  obtain M where moebius-pt M z1 = w1 moebius-pt M z2 = w2 moebius-pt M
z3 = w3
    using ex-unique-moebius-three-points[OF assms]
    by auto
  thus ?thesis
    using ex-unique-moebius-three-points[OF assms]
    unfolding Ex1-def
    by (rule-tac x=moebius-pt M in exI) (auto simp add: is-moebius-def)
qed

lemma is-cross-ratio-01inf:
assumes z1 ≠ z2 z1 ≠ z3 z2 ≠ z3 is-moebius f
assumes f z1 = 0h f z2 = 1h f z3 = ∞h
shows f = (λ z. cross-ratio z z1 z2 z3)
using assms

```

```

using cross-ratio-0[OF ⟨z1 ≠ z2⟩ ⟨z1 ≠ z3⟩] cross-ratio-1[OF ⟨z1 ≠ z2⟩ ⟨z2 ≠
z3⟩] cross-ratio-inf[OF ⟨z1 ≠ z3⟩ ⟨z2 ≠ z3⟩]
using is-moebius-cross-ratio[OF ⟨z1 ≠ z2⟩ ⟨z2 ≠ z3⟩ ⟨z1 ≠ z3⟩]
using ex-unique-moebius-three-points-fun[OF ⟨z1 ≠ z2⟩ ⟨z1 ≠ z3⟩ ⟨z2 ≠ z3⟩, of
0h 1h ∞h]
by auto

```

lemma moebius-preserve-cross-ratio:

assumes $z1 \neq z2 \ z1 \neq z3 \ z2 \neq z3$

shows $\text{cross-ratio } z \ z1 \ z2 \ z3 = \text{cross-ratio } (\text{moebius-pt } M \ z) (\text{moebius-pt } M \ z1)$
 $(\text{moebius-pt } M \ z2) (\text{moebius-pt } M \ z3)$

proof –

let $?f = \lambda z. \text{cross-ratio } z \ z1 \ z2 \ z3$

let $?M = \text{moebius-pt } M$

let $?iM = \text{inv } ?M$

have $(?f \circ ?iM) (?M \ z1) = 0_h$

using bij-moebius-pt[of M] cross-ratio-0[OF ⟨z1 ≠ z2⟩ ⟨z1 ≠ z3⟩]
by (simp add: bij-def)

moreover

have $(?f \circ ?iM) (?M \ z2) = 1_h$

using bij-moebius-pt[of M] cross-ratio-1[OF ⟨z1 ≠ z2⟩ ⟨z2 ≠ z3⟩]
by (simp add: bij-def)

moreover

have $(?f \circ ?iM) (?M \ z3) = \infty_h$

using bij-moebius-pt[of M] cross-ratio-inf[OF ⟨z1 ≠ z3⟩ ⟨z2 ≠ z3⟩]
by (simp add: bij-def)

moreover

have is-moebius (?f ∘ ?iM)
by (rule is-moebius-comp, rule is-moebius-cross-ratio[OF ⟨z1 ≠ z2⟩ ⟨z2 ≠ z3⟩
⟨z1 ≠ z3⟩], rule is-moebius-inv, auto simp add: is-moebius-def)

moreover

have $?M \ z1 \neq ?M \ z2 \ ?M \ z1 \neq ?M \ z3 \ ?M \ z2 \neq ?M \ z3$

using assms
using bij-moebius-pt[of M]
unfolding bij-def inj-on-def
by blast+

ultimately

have $?f \circ ?iM = (\lambda z. \text{cross-ratio } z (?M \ z1) (?M \ z2) (?M \ z3))$

using assms
using is-cross-ratio-01inf[of ?M \ z1 ?M \ z2 ?M \ z3 ?f ∘ ?iM]
by simp

moreover

have $(?f \circ ?iM) (?M \ z) = \text{cross-ratio } z \ z1 \ z2 \ z3$

using bij-moebius-pt[of M]
by (simp add: bij-def)

moreover

have $(\lambda z. \text{cross-ratio } z (?M \ z1) (?M \ z2) (?M \ z3)) (?M \ z) = \text{cross-ratio } (?M \ z)$
 $(?M \ z1) (?M \ z2) (?M \ z3)$

```

by simp
ultimately
show ?thesis
  by simp
qed

lemma fixed-points-0inf':
  assumes moebius-pt M 0h = 0h moebius-pt M ∞h = ∞h
  shows ∃ k::complex-homo. (k ≠ 0h ∧ k ≠ ∞h) ∧ (∀ z. moebius-pt M z = k *h z)
using assms
proof (transfer)
  fix M
  obtain a b c d where MM: Rep-moebius-mat M = (a, b, c, d)
    by (cases M) (auto simp add: Abs-moebius-mat-inverse)
  assume moebius-pt-rep M zero-homo-rep ≈ zero-homo-rep moebius-pt-rep M
  inf-homo-rep ≈ inf-homo-rep
  hence b = 0 c = 0
    using MM
    by auto
  hence *: a ≠ 0 ∧ d ≠ 0
    using Rep-moebius-mat[of M] MM
    by auto
  show ∃ k. (¬ k ≈ zero-homo-rep ∧ ¬ k ≈ inf-homo-rep) ∧ (∀ z. moebius-pt-rep M z ≈ k *hc z)
    proof (rule-tac x=Abs-homo-coords (a, d) in exI, rule conjI)
      show ¬ Abs-homo-coords (a, d) ≈ zero-homo-rep ∧ ¬ Abs-homo-coords (a, d)
        ≈ inf-homo-rep
        using *
        by (auto simp add: Abs-homo-coords-inverse)
    next
      show ∀ z. moebius-pt-rep M z ≈ Abs-homo-coords (a, d) *hc z
      proof
        fix z
        obtain z1 z2 where zz: Rep-homo-coords z = (z1, z2)
          by (rule obtain-homo-coords)
        thus moebius-pt-rep M z ≈ Abs-homo-coords (a, d) *hc z
          using MM * ⟨b = 0⟩ ⟨c = 0⟩ mult-homo-coords-Rep[Abs-homo-coords (a, d) a d z z1 z2] Rep-homo-coords[of z]
            by (simp add: Abs-homo-coords-inverse) (rule-tac x=1 in exI, simp)
      qed
    qed
  qed

lemma fixed-points-0inf:
  assumes moebius-pt M 0h = 0h moebius-pt M ∞h = ∞h
  shows ∃ k::complex-homo. (k ≠ 0h ∧ k ≠ ∞h) ∧ moebius-pt M = (λ z. k *h z)
using fixed-points-0inf'[OF assms]
by auto

```

10.7 Pole

definition *is-pole* **where**

$$\text{is-pole } M z \longleftrightarrow \text{moebius-pt } M z = \infty_h$$

lemma *ex1-pole*:

$$\exists! z. \text{is-pole } M z$$

using *bij-moebius-pt*[*of M*]

unfolding *is-pole-def* *bij-def* *inj-on-def* *surj-def*

unfolding *Ex1-def*

by (*metis UNIV-I*)

definition *pole* **where** *pole M* = (*THE z. is-pole M z*)

lemma *pole-mk-moebius*:

assumes *is-pole* (*mk-moebius a b c d*) $z c \neq 0$ $a*d - b*c \neq 0$

shows *z = of-complex* ($-d/c$)

proof-

let *?t1 = translation-moebius* (*a / c*)

let *?rd = rotation-dilatation-moebius* (($b * c - a * d$) / ($c * c$))

let *?r = reciprocal-moebius*

let *?t2 = translation-moebius* (*d / c*)

have *moebius-pt* (*?rd + ?r + ?t2*) $z = \infty_h$

using *assms*

unfolding *is-pole-def*

apply (*subst (asm) moebius-decomposition*)

apply (*auto simp add: moebius-comp[symmetric] translation-moebius-def*)

apply (*subst similarity-only-inf-to-inf[of 1 a/c], auto*)

done

hence *moebius-pt* (*?r + ?t2*) $z = \infty_h$

using $\langle a*d - b*c \neq 0 \rangle \langle c \neq 0 \rangle$

unfolding *rotation-dilatation-moebius-def*

apply (*simp add: moebius-comp[symmetric]*)

apply (*subst similarity-only-inf-to-inf[of (b*c-a*d)/(c*c) 0], auto*)

done

hence *moebius-pt* *?t2* $z = 0_h$

apply (*simp add: moebius-comp[symmetric]*)

apply (*subst (asm) reciprocal-moebius[symmetric]*)

apply (*subst reciprocal-homo-only-0-to-inf, auto*)

done

thus *?thesis*

using *moebius-pt-invert[of 0_h ?t2 z]* *moebius-translation-inv[of d/c]*

by *simp (subst zero-of-complex[symmetric], simp del: zero-of-complex)*

qed

lemma *pole-similarity*:

assumes *is-pole* (*similarity-moebius a b*) $z a \neq 0$

shows *z = infinity_h*

using *assms*

unfolding *is-pole-def*

```
using similarity-only-inf-to-inf[of a b z]
by simp
```

10.8 Antihomographies

```
definition is-antihomography where
is-antihomography f  $\longleftrightarrow$  ( $\exists f'. is\text{-moebius } f' \wedge f = f' \circ cnj\text{-homo}$ )
```

```
lemma is-antihomography inversion-homo
using reciprocal-moebius
unfolding inversion-homo-sym is-antihomography-def
by (auto simp add: is-moebius-def)
```

10.9 Classification

```
lemma similarity-scale-1:
assumes k  $\neq 0$ 
shows similarity (k *sm I) M = similarity I M
using assms
unfolding similarity-def
using mat-inv-mult-sm[of k I]
by simp
```

```
lemma similarity-scale-2:
shows similarity I (k *sm M) = k *sm (similarity I M)
unfolding similarity-def
by auto
```

```
lemma [simp]: mat-trace (k *sm M) = k * mat-trace M
by (cases M) (simp add: field-simps)
```

```
definition moebius-mb-rep where
moebius-mb-rep I M = Abs-moebius-mat (similarity (Rep-moebius-mat I) (Rep-moebius-mat M))
```

```
lemma moebius-mb-rep-Rep [simp]:
Rep-moebius-mat (moebius-mb-rep I M) = similarity (Rep-moebius-mat I) (Rep-moebius-mat M)
using Rep-moebius-mat[of I] Rep-moebius-mat[of M]
unfolding moebius-mb-rep-def
by (simp add: mat-det-similarity Abs-moebius-mat-inverse)
```

```
lift-definition moebius-mb :: moebius  $\Rightarrow$  moebius  $\Rightarrow$  moebius is moebius-mb-rep
proof-
fix M M' I I'
assume moebius-mat-eq M M' moebius-mat-eq I I'
thus moebius-mat-eq (moebius-mb-rep I M) (moebius-mb-rep I' M')
by (auto simp add: similarity-scale-1 similarity-scale-2)
qed
```

```

definition similarity-invar-rep where
  similarity-invar-rep M =
    (let M = Rep-moebius-mat M
     in (mat-trace M)2 / mat-det M - 4)

lift-definition similarity-invar :: moebius  $\Rightarrow$  complex is similarity-invar-rep
by (auto simp add: similarity-invar-rep-def Let-def power2-eq-square)

lemma
  similarity-invar (moebius-mb I M) = similarity-invar M
proof transfer
  fix I M
  show similarity-invar-rep (moebius-mb-rep I M) = similarity-invar-rep M
  using Rep-moebius-mat[of I] Rep-moebius-mat[of M]
  by (simp add: similarity-invar-rep-def Let-def mat-trace-similarity mat-det-similarity)
qed

definition similar where
  similar M1 M2  $\longleftrightarrow$  ( $\exists$  I. moebius-mb I M1 = M2)

lemma [simp]: similarity eye M = M
unfolding similarity-def
by simp (metis eye-def mat-eye-l mat-eye-r)

lemma [simp]: similarity (1, 0, 0, 1) M = M
unfolding eye-def[symmetric]
by (simp del: eye-def)

lemma similarity-comp:
  assumes mat-det I1  $\neq$  0 mat-det I2  $\neq$  0
  shows similarity I1 (similarity I2 M) = similarity (I2 *mm I1) M
using assms
unfolding similarity-def
by (simp add: mult-mm-assoc mat-inv-mult-mm)

lemma similarity-inv:
  assumes similarity I M1 = M2 mat-det I  $\neq$  0
  shows similarity (mat-inv I) M2 = M1
using assms
unfolding similarity-def
by simp (metis mat-eye-l mult-mm-assoc mult-mm-inv-r)

lemma similar-refl [simp]: similar M M
unfolding similar-def
by (rule-tac x=id-moebius in exI) (transfer, simp, rule-tac x=1 in exI, auto)

lemma similar-sym:
  assumes similar M1 M2
  shows similar M2 M1

```

```

proof-
  from assms obtain I where  $M2 = \text{moebius-mb } I M1$ 
    unfolding similar-def
    by auto
  hence  $M1 = \text{moebius-mb } (\text{moebius-inv } I) M2$ 
  proof transfer
    fix  $M2 I M1$ 
    assume  $\text{moebius-mat-eq } M2 (\text{moebius-mb-rep } I M1)$ 
    then obtain  $k$  where  $k \neq 0$  similarity (Rep-moebius-mat I) (Rep-moebius-mat  $M1) = k *_{sm} \text{Rep-moebius-mat } M2$ 
      by auto
      thus  $\text{moebius-mat-eq } M1 (\text{moebius-mb-rep } (\text{moebius-inv-rep } I) M2)$ 
      using Rep-moebius-mat[of I] similarity-inv[of Rep-moebius-mat I Rep-moebius-mat  $M1$   $k *_{sm} \text{Rep-moebius-mat } M2]$ 
        by (auto simp add: similarity-scale-2) (rule-tac  $x=1/k$  in exI, simp, metis mult-sm-inv-l)
      qed
      thus ?thesis
        unfolding similar-def
        by auto
    qed

lemma similar-trans:
  assumes  $\text{similar } M1 M2 \text{ similar } M2 M3$ 
  shows  $\text{similar } M1 M3$ 
proof-
  obtain  $I1 I2$  where  $\text{moebius-mb } I1 M1 = M2 \text{ moebius-mb } I2 M2 = M3$ 
    using assms
    by (auto simp add: similar-def)
  thus ?thesis
    unfolding similar-def
    proof (rule-tac  $x=\text{moebius-comp } I1 I2$  in exI, transfer)
      fix  $I1 I2 M1 M2 M3$ 
      assume  $\text{moebius-mat-eq } (\text{moebius-mb-rep } I1 M1) M2$ 
         $\text{moebius-mat-eq } (\text{moebius-mb-rep } I2 M2) M3$ 
      thus  $\text{moebius-mat-eq } (\text{moebius-mb-rep } (\text{moebius-comp-rep } I1 I2) M1) M3$ 
        using Rep-moebius-mat[of I1] Rep-moebius-mat[of I2]
        by (auto simp add: similarity-scale-2 similarity-comp) (rule-tac  $x=ka*k$  in exI, simp)
      qed
    qed

end

```

11 Circline

theory *Circline*

```

imports Moebius HermiteanMatrices ElementaryComplexGeometry RiemannSphere
Angles
begin

```

11.1 Circline definition

```

typedef circline-mat = {H. hermitean H ∧ H ≠ mat-zero}
by (rule-tac x=eye in exI) (auto simp add: hermitean-def mat-adj-def mat-cnj-def)

lemma circline-mat-mult-sm-Rep [simp]:
  assumes k ≠ 0
  shows Rep-circline-mat (Abs-circline-mat ((cor k) *sm (Rep-circline-mat H)))
= (cor k) *sm (Rep-circline-mat H)
using assms Rep-circline-mat[of H]
using hermitean-mult-real[of Rep-circline-mat H k] nonzero-mult-real[of Rep-circline-mat
H cor k]
by (simp add: Abs-circline-mat-inverse)

definition circline-mat-eq where
  [simp]: circline-mat-eq A B ↔ (∃ k::real. k ≠ 0 ∧ Rep-circline-mat B =
complex-of-real k *sm (Rep-circline-mat A))

lemma [simp]: circline-mat-eq H H
  by (simp, rule-tac x=1 in exI, simp)

quotient-type circline = circline-mat / circline-mat-eq
proof (rule equivpI)
  show reflp circline-mat-eq
    unfolding reflp-def
    by (auto, rule-tac x=1 in exI, simp)
next
  show symp circline-mat-eq
    unfolding symp-def
    by (auto, rule-tac x=1/k in exI, simp)
next
  show transp circline-mat-eq
    unfolding transp-def
    by (auto, rule-tac x=ka*k in exI, simp)
qed

Circline with specified matrix

definition mk-circline-rep where
  mk-circline-rep A B C D = Abs-circline-mat (A, B, C, D)

lift-definition mk-circline :: complex ⇒ complex ⇒ complex ⇒ complex ⇒ cir-
cline is mk-circline-rep
by (simp del: circline-mat-eq-def)

lemma ex-mk-circline:

```

```

shows  $\exists A B C D. H = \text{mk-circline } A B C D \wedge \text{hermitean } (A, B, C, D) \wedge (A, B, C, D) \neq \text{mat-zero}$ 
proof transfer
fix  $H$ 
obtain  $A B C D$  where  $\text{Rep-circline-mat } H = (A, B, C, D)$ 
by (cases Rep-circline-mat  $H$ , auto)
hence  $\text{circline-mat-eq } H (\text{mk-circline-rep } A B C D) \wedge \text{hermitean } (A, B, C, D)$ 
 $\wedge (A, B, C, D) \neq \text{mat-zero}$ 
using Rep-circline-mat[of  $H$ ]
by (auto simp add: mk-circline-rep-def Abs-circline-mat-inverse) (rule-tac  $x=1$ 
in exI, simp)+
thus  $\exists A B C D. \text{circline-mat-eq } H (\text{mk-circline-rep } A B C D) \wedge \text{hermitean } (A, B, C, D) \wedge (A, B, C, D) \neq \text{mat-zero}$ 
by blast
qed

circline type

definition circline-type-rep where
circline-type-rep  $H = \text{sgn}(\text{Re}(\text{mat-det}(\text{Rep-circline-mat } H)))$ 

lift-definition circline-type :: circline  $\Rightarrow$  real is circline-type-rep
proof-
fix  $H H'$ 
assume circline-mat-eq  $H H'$ 
thus circline-type-rep  $H = \text{circline-type-rep } H'$ 
by (auto simp add: circline-type-rep-def sgn-mult) (metis not-real-square-gt-zero
real-sgn-pos sgn-mult)
qed

lemma circline-type: circline-type  $H = -1 \vee \text{circline-type } H = 0 \vee \text{circline-type }$ 
 $H = 1$ 
proof transfer
fix  $H$ 
show circline-type-rep  $H = -1 \vee \text{circline-type-rep } H = 0 \vee \text{circline-type-rep } H$ 
= 1
unfolding circline-type-rep-def
using Rep-circline-mat[of  $H$ ]
by (metis linorder-cases real-sgn-neg real-sgn-pos sgn-zero-iff)
qed

on-circline, circline-set

definition on-circline-rep where
on-circline-rep  $H z \leftrightarrow$ 
(let  $z = \text{Rep-homo-coords } z;$ 
 $H = \text{Rep-circline-mat } H$ 
in quad-form  $z H = 0$ )

```

lift-definition on-circline :: circline \Rightarrow complex-homo \Rightarrow bool is on-circline-rep

```
by (auto simp add: on-circline-rep-def quad-form-scale-m quad-form-scale-v Let-def
      simp del: vec-cnj-sv quad-form-def)
```

```
definition circline-set :: circline ⇒ complex-homo set where
  circline-set H = {z. on-circline H z}
```

Circlines trough 0 and inf

```
definition circline-A0-rep where
  circline-A0-rep H ↔
    (let (A, B, C, D) = Rep-circline-mat H in A = 0)
```

```
lift-definition circline-A0 :: circline ⇒ bool is circline-A0-rep
by (auto simp add: circline-A0-rep-def)
```

```
definition circline-D0-rep where
  circline-D0-rep H ↔
    (let (A, B, C, D) = Rep-circline-mat H in D = 0)
```

```
abbreviation is-line where
  is-line H ≡ circline-A0 H
```

```
abbreviation is-circle where
  is-circle H ≡ ¬ circline-A0 H
```

```
lift-definition circline-D0 :: circline ⇒ bool is circline-D0-rep
by (auto simp add: circline-D0-rep-def)
```

```
lemma inf-on-circline-rep: on-circline-rep H inf-homo-rep ↔ circline-A0-rep H
by (simp add: on-circline-rep-def Let-def circline-A0-rep-def split-def) (cases Rep-circline-mat
H, simp add: vec-cnj-def)
```

```
lemma
  inf-in-circline-set: ∞_h ∈ circline-set H ↔ is-line H
unfolding circline-set-def
apply simp
apply (transfer)
using inf-on-circline-rep
by simp
```

```
lemma zero-on-circline-rep: on-circline-rep H zero-homo-rep ↔ circline-D0-rep
H
using Rep-circline-mat[of H]
by (simp add: circline-D0-rep-def on-circline-rep-def split-def Let-def Abs-homo-coords-inverse
Abs-circline-mat-inverse vec-cnj-def) (cases Rep-circline-mat H, simp)
```

```
lemma zero-in-circline-set: 0_h ∈ circline-set H ↔ circline-D0 H
unfolding circline-set-def
apply simp
apply (transfer)
```

```
using zero-on-circline-rep
by simp
```

Connection with circlines in classic complex plane

```
lemma classic-circline:
assumes H = mk-circline A B C D hermitean (A, B, C, D) ∧ (A, B, C, D) ≠ mat-zero
shows circline-set H - {∞h} = of-complex ` circline (Re A) B (Re D)
using assms
unfolding circline-set-def
proof (safe)
fix z
assume hermitean (A, B, C, D) (A, B, C, D) ≠ mat-zero z ∈ circline (Re A)
B (Re D)
thus on-circline (mk-circline A B C D) (of-complex z)
using hermitean-elems[of A B C D]
by (transfer) (simp del: mat-zero-def add: on-circline-rep-def Let-def mk-circline-rep-def
Abs-circline-mat-inverse circline-def vec-cnj-def field-simps complex-of-real-Re)
next
fix z
assume of-complex z = ∞h
thus False
by simp
next
fix z
assume hermitean (A, B, C, D) (A, B, C, D) ≠ mat-zero on-circline (mk-circline
A B C D) z z ∉ of-complex ` circline (Re A) B (Re D)
moreover
have z ≠ ∞h → z ∈ of-complex ` circline (Re A) B (Re D)
proof
assume z ≠ ∞h
show z ∈ of-complex ` circline (Re A) B (Re D)
proof
show z = of-complex (to-complex z)
using ⟨z ≠ ∞h⟩
by simp
next
show to-complex z ∈ circline (Re A) B (Re D)
using ⟨on-circline (mk-circline A B C D) z⟩ ⟨z ≠ ∞h⟩ ⟨hermitean (A, B,
C, D)⟩ ⟨(A, B, C, D) ≠ mat-zero⟩
proof (transfer)
fix A B C D z
obtain z1 z2 where zz: Rep-homo-coords z = (z1, z2)
by (rule obtain-homo-coords)
assume *: ¬ z ≈ inf-homo-rep on-circline-rep (mk-circline-rep A B C D) z
hermitean (A, B, C, D) (A, B, C, D) ≠ mat-zero
have z2 ≠ 0
using ⟨¬ z ≈ inf-homo-rep⟩ Rep-homo-coords[of z] zz
by auto (erule-tac x=1/z1 in allE, simp)
```

```

thus to-complex-homo-coords  $z \in \text{circline}(\text{Re } A) B (\text{Re } D)$ 
  using *
  using zz
  using hermitean-elems[of A B C D]
  by (simp add: mk-circline-rep-def on-circline-rep-def to-complex-homo-coords-def
    Let-def Abs-circline-mat-inverse vec-cnj-def complex-cnj circline-def complex-of-real-Re
    field-simps del: mat-zero-def)
qed
qed
qed
ultimately
show  $z = \infty_h$ 
  by simp
qed

definition mk-circle-rep where
mk-circle-rep a r = Abs-circline-mat (1, -a, -cnj a, a*cnj a - cor r*cor r)
lift-definition mk-circle :: complex ⇒ real ⇒ circline is mk-circle-rep
by (simp del: circline-mat-eq-def)

lemma mk-circle-rep-Rep
[simp]: Rep-circline-mat (mk-circle-rep a r) = (1, -a, -cnj a, a*cnj a - cor r*cor r)
by (simp add: mk-circle-rep-def Abs-circline-mat-inverse hermitean-def mat-adj-def
  mat-cnj-def complex-cnj)

lemma is-circle-mk-circle: is-circle (mk-circle a r)
by transfer (simp add: circline-A0-rep-def)

lemma
assumes  $r \geq 0$ 
shows circline-set (mk-circle a r) = of-complex ` {z. cmod(z - a) = r}
proof-
let ?A = 1 and ?B = -a and ?C = -cnj a and ?D = a*cnj a - cor r*cor r
have *: (?A, ?B, ?C, ?D) ∈ {H. hermitean H ∧ H ≠ mat-zero}
  by (simp add: hermitean-def mat-adj-def mat-cnj-def complex-cnj)
have mk-circle a r = mk-circline ?A ?B ?C ?D
using *
  by transfer (simp add: mk-circline-rep-def Abs-circline-mat-inverse, rule-tac
    x=1 in exI, simp)
hence circline-set (mk-circle a r) - {∞_h} = of-complex ` circline ?A ?B (Re ?D)
using classic-circline[of mk-circle a r ?A ?B ?C ?D] *
  by simp
moreover
have circline ?A ?B (Re ?D) = circle a r
  by (rule circline-circle[of ?A Re ?D ?B circline ?A ?B (Re ?D) a r*r r], simp-all
    add: cmod-square ⟨r ≥ 0⟩)
moreover
have ∞_h ∉ circline-set (mk-circle a r)

```

```

using inf-in-circline-set[of mk-circle a r] is-circle-mk-circle[of a r]
by auto
ultimately
show ?thesis
  unfolding circle-def
  by simp
qed

definition mk-line-rep where mk-line-rep z1 z2 =
  (let B = ii*(z2-z1) in Abs-circline-mat (0, B, cnj B, -cnj-mix B z1))
lift-definition mk-line :: complex  $\Rightarrow$  complex  $\Rightarrow$  circline is mk-line-rep
by (simp del: circline-mat-eq-def)

lemma mk-line-rep-Rep [simp]:
assumes z1  $\neq$  z2
shows Rep-circline-mat (mk-line-rep z1 z2) =
  (let B = ii*(z2-z1) in (0, B, cnj B, -cnj-mix B z1))
using assms
by (simp add: mk-line-rep-def Let-def Abs-circline-mat-inverse hermitean-def mat-adj-def
mat-cnj-def complex-cnj)

lemma circline-line':
assumes z1  $\neq$  z2
shows circline 0 (i * (z2 - z1)) (Re (- cnj-mix (i * (z2 - z1)) z1)) = line z1
z2
proof-
  let ?B = ii * (z2 - z1)
  let ?D = Re (- cnj-mix ?B z1)
  have circline 0 ?B ?D = {z. cnj ?B*z + ?B*cnj z + complex-of-real ?D = 0}
    using assms
    by (simp add: circline-def)
  moreover
  have is-real (- cnj-mix (i * (z2 - z1)) z1)
    using cnj-mix-real[of ?B z1]
    by auto
  hence {z. cnj ?B*z + ?B*cnj z + complex-of-real ?D = 0} =
    {z. cnj ?B*z + ?B*cnj z - (cnj ?B*z1 + ?B*cnj z1) = 0}
    by (subst complex-of-real-Re, simp, simp add: complex-diff-def)
  moreover
  have line z1 z2 = {z. cnj-mix (i * (z2 - z1)) z - cnj-mix (i * (z2 - z1)) z1 = 0}
    using line-equation[of z1 z2 ?B] assms
    unfolding rot90-ii
    by simp
  ultimately
  show ?thesis
    by simp
qed

```

```

lemma
  assumes  $z1 \neq z2$ 
  shows circline-set (mk-line  $z1\ z2$ ) -  $\{\infty_h\}$  = of-complex ‘line  $z1\ z2$ 

proof-
  let  $?A = 0$  and  $?B = ii*(z2 - z1)$ 
  let  $?C = cnj\ ?B$  and  $?D = -cnj\cdot mix\ ?B\ z1$ 
  have  $*: (?A, ?B, ?C, ?D) \in \{H. hermitean\ H \wedge H \neq mat\cdot zero\}$ 
  using assms
  by (simp add: hermitean-def mat-adj-def mat-cnj-def complex-cnj)
  have mk-line  $z1\ z2$  = mk-circline  $?A\ ?B\ ?C\ ?D$ 
  using * assms
  by transfer (simp add: mk-circline-rep-def Abs-circline-mat-inverse Let-def,
  rule-tac x=1 in exI, simp)
  hence circline-set (mk-line  $z1\ z2$ ) -  $\{\infty_h\}$  = of-complex ‘circline  $?A\ ?B$  (Re  $?D$ )
  using classic-circline[of mk-line  $z1\ z2\ ?A\ ?B\ ?C\ ?D$ ] *
  by simp
  moreover
  have circline  $?A\ ?B$  (Re  $?D$ ) = line  $z1\ z2$ 
  using  $(z1 \neq z2)$ 
  using circline-line'
  by simp
  ultimately
  show thesis
  by simp
qed

definition euclidean-circle-rep where
  euclidean-circle-rep  $H = (\text{let } (A, B, C, D) = Rep\cdot circline\cdot mat\ H \text{ in } (-B/A, \sqrt{Re\ ((B*C - A*D)/(A*A))}))$ 

lift-definition euclidean-circle :: circline  $\Rightarrow$  complex  $\times$  real is euclidean-circle-rep
proof-
  fix  $H1\ H2$ 
  obtain  $A1\ B1\ C1\ D1$  where  $HH1: Rep\cdot circline\cdot mat\ H1 = (A1, B1, C1, D1)$ 
  by (cases Rep-circline-mat  $H1$ ) auto
  obtain  $A2\ B2\ C2\ D2$  where  $HH2: Rep\cdot circline\cdot mat\ H2 = (A2, B2, C2, D2)$ 
  by (cases Rep-circline-mat  $H2$ ) auto
  assume circline-mat-eq  $H1\ H2$ 
  then obtain  $k$  where  $k \neq 0$  and  $*: A2 = cor\ k * A1\ B2 = cor\ k * B1\ C2 = cor\ k * C1\ D2 = cor\ k * D1$ 
  using  $HH1\ HH2$ 
  by auto
  have  $(cor\ k * B1 * (cor\ k * C1) - cor\ k * A1 * (cor\ k * D1)) = (cor\ k)^2 * (B1*C1 - A1*D1)$ 
   $(cor\ k * A1 * (cor\ k * A1)) = (cor\ k)^2 * (A1*A1)$ 
  by (auto simp add: field-simps power2-eq-square)
  hence  $(cor\ k * B1 * (cor\ k * C1) - cor\ k * A1 * (cor\ k * D1)) / (cor\ k * A1 * (cor\ k * A1)) = (B1*C1 - A1*D1) / (A1*A1)$ 

```

```

using ⟨ $k \neq 0$ ⟩
by (simp add: power2-eq-square)
thus euclidean-circle-rep H1 = euclidean-circle-rep H2
    using HH1 HH2 * Rep-circline-mat[of H2]
    by (auto simp add: euclidean-circle-rep-def)
qed

lemma classic-circle:
assumes is-circle H (a, r) = euclidean-circle H circline-type H ≤ 0
shows circline-set H = of-complex ‘circle a r
proof –
  obtain A B C D where *: H = mk-circline A B C D hermitean (A, B, C, D)
  (A, B, C, D) ≠ mat-zero
    using ex-mk-circline[of H]
    by auto
  have is-real A is-real D C = cnj B
    using * hermitean-elems
    by auto

  have Re (A*D - B*C) ≤ 0
    using ⟨circline-type H ≤ 0⟩ *
    by simp (transfer, simp add: circline-type-rep-def mk-circline-rep-def Abs-circline-mat-inverse,
      smt real-sgn-pos)

  hence **: Re A * Re D ≤ (cmod B)2
    using ⟨is-real A⟩ ⟨is-real D⟩ ⟨C = cnj B⟩
    by (simp add: cmod-square)

  have A ≠ 0
    using ⟨is-circle H⟩ * ⟨is-real A⟩
    by simp (transfer, simp add: circline-A0-rep-def mk-circline-rep-def Abs-circline-mat-inverse)
  hence Re A ≠ 0
    using ⟨is-real A⟩
    by (cases A, simp)

  have ***: ∞h ∉ circline-set H
    using * inf-in-circline-set[of H] ⟨is-circle H⟩
    by simp

  let ?a = -B/A
  let ?r2 = ((cmod B)2 - Re A * Re D) / (Re A)2
  let ?r = sqrt ?r2

  have ?a = a ∧ ?r = r
    using ⟨(a, r) = euclidean-circle H⟩
    using * ⟨is-real A⟩ ⟨is-real D⟩ ⟨C = cnj B⟩ ⟨A ≠ 0⟩
    apply simp
    apply transfer
    apply (simp add: euclidean-circle-rep-def mk-circline-rep-def Abs-circline-mat-inverse)

```

```

apply (subst Re-divide-real)
apply (simp-all add: cmod-square, simp add: power2-eq-square)
done

show ?thesis
  using *** (Re A ≠ 0) (is-real A) (C = cnj B) (?a = a ∧ ?r = r)
  using classic-circline[of H A B C D] assms circline-circle[of Re A Re D B
circline (Re A) B (Re D) ?a ?r2 ?r]
    by (simp add: complex-of-real-Re circle-def)
qed

definition
euclidean-line-rep H =
  (let (A, B, C, D) = Rep-circline-mat H;
   z1 = -(D*B)/(2*B*C);
   z2 = z1 + ii*sgn (if arg B > 0 then -B else B)
   in (z1, z2))

lift-definition euclidean-line :: circline ⇒ complex × complex is euclidean-line-rep
proof –
fix H1 H2
obtain A1 B1 C1 D1 where HH1: Rep-circline-mat H1 = (A1, B1, C1, D1)
  by (cases Rep-circline-mat H1) auto
obtain A2 B2 C2 D2 where HH2: Rep-circline-mat H2 = (A2, B2, C2, D2)
  by (cases Rep-circline-mat H2) auto
assume circline-mat-eq H1 H2
then obtain k where k ≠ 0 and *: A2 = cor k * A1 B2 = cor k * B1 C2 =
cor k * C1 D2 = cor k * D1
  using HH1 HH2
  by auto
have 1: B1 ≠ 0 ∧ 0 < arg B1 → 0 < arg (-B1)
  using MoreComplex.canon-ang-plus-pi1[of arg B1] arg-bounded[of B1]
  by (auto simp add: arg-uminus)
have 2: B1 ≠ 0 ∧ 0 < arg B1 → 0 < arg (-B1)
  using MoreComplex.canon-ang-plus-pi2[of arg B1] arg-bounded[of B1]
  by (auto simp add: arg-uminus)

show euclidean-line-rep H1 = euclidean-line-rep H2
  using HH1 HH2 * (k ≠ 0)
  by (cases k > 0) (auto simp add: euclidean-line-rep-def Let-def, simp-all add:
sgn-eq arg-mult-real-positive arg-mult-real-negative 1 2)
qed

lemma classic-line:
assumes is-line H (z1, z2) = euclidean-line H circline-type H < 0
shows circline-set H - {∞h} = of-complex ‘line z1 z2
proof –
obtain A B C D where *: H = mk-circline A B C D hermitean (A, B, C, D)
(A, B, C, D) ≠ mat-zero

```

```

using ex-mk-circline[of H]
by auto
have is-real A is-real D C = cnj B
  using * hermitean-elems
  by auto
have Re A = 0
  using ⟨is-line H⟩* ⟨is-real A⟩ ⟨is-real D⟩ ⟨C = cnj B⟩
  by transfer (auto simp add: circline-A0-rep-def mk-circline-rep-def Abs-circline-mat-inverse)
have B ≠ 0
  using ⟨Re A = 0⟩ ⟨is-real A⟩ ⟨is-real D⟩ ⟨C = cnj B⟩ * ⟨circline-type H < 0⟩
  by transfer (auto simp add: circline-type-rep-def mk-circline-rep-def Abs-circline-mat-inverse,
(case-tac Rep-circline-mat H, simp)+)

let ?z1 = - cor (Re D) * B / (2 * B * cnj B)
let ?z2 = ?z1 + i * sgn (if 0 < arg B then - B else B)
have z1 = ?z1 ∧ z2 = ?z2
  using ⟨(z1, z2) = euclidean-line H⟩ * ⟨is-real A⟩ ⟨is-real D⟩ ⟨C = cnj B⟩
  by simp (transfer, simp add: euclidean-line-rep-def mk-circline-rep-def Abs-circline-mat-inverse
Let-def complex-of-real-Re)
thus ?thesis
  using *
  using classic-circline[of H A B C D] circline-line[of Re A B circline (Re A) B
(Re D) Re D ?z1 ?z2] ⟨Re A = 0⟩ ⟨B ≠ 0⟩
  by simp
qed

```

11.2 Connections with circles on the Riemann sphere

```

definition inv-stereographic-circline-rep where
inv-stereographic-circline-rep H =
(let (A, B, C, D) = Rep-circline-mat H in
Abs-plane-vec (Re (B+C), Re(ii*(C-B)), Re(A-D), Re(D+A)))

lemma inv-stereographic-circline-rep-Rep [simp]:
Rep-plane-vec (inv-stereographic-circline-rep H) =
(let (A, B, C, D) = Rep-circline-mat H in (Re (B+C), Re(ii*(C-B)),
Re(A-D), Re(D+A)))
proof-
obtain A B C D where HH: Rep-circline-mat H = (A, B, C, D)
by (cases Rep-circline-mat H) auto
have *: is-real A is-real D C = cnj B
  using HH Rep-circline-mat[of H] hermitean-elems[of A B C D]
  by auto
have Re B + Re C = 0 ∧ Im B - Im C = 0 ∧ Re A - Re D = 0 ∧ Re A +
Re D = 0 → (A, B, C, D) = mat-zero
  using *
  by auto (metis complex-of-real-Re of-real-0)+
hence **: Re B + Re C ≠ 0 ∨ Im B - Im C ≠ 0 ∨ Re A - Re D ≠ 0 ∨ Re
D + Re A ≠ 0

```

```

using Rep-circline-mat[of H] HH
by auto
thus ?thesis
using HH
by (simp add: Abs-plane-vec-inverse inv-stereographic-circline-rep-def)
qed

lift-definition inv-stereographic-circline :: circline  $\Rightarrow$  plane is inv-stereographic-circline-rep
proof-
  fix H1 H2
  obtain A1 B1 C1 D1 where HH1: Rep-circline-mat H1 = (A1, B1, C1, D1)
    by (cases Rep-circline-mat H1) auto
  obtain A2 B2 C2 D2 where HH2: Rep-circline-mat H2 = (A2, B2, C2, D2)
    by (cases Rep-circline-mat H2) auto
  have *: is-real A1 is-real A2 is-real D1 is-real D2 C1 = cnj B1 C2 = cnj B2
  using HH1 HH2 Rep-circline-mat[of H1] Rep-circline-mat[of H2] hermitean-elems[of
  A1 B1 C1 D1] hermitean-elems[of A2 B2 C2 D2]
  by auto

  assume circline-mat-eq H1 H2
  thus plane-vec-eq (inv-stereographic-circline-rep H1) (inv-stereographic-circline-rep
  H2)
    using HH1 HH2 *
    by (simp add: plane-vec-eq-def) (erule exE, rule-tac x=k in exI, simp add:
  field-simps)
  qed

definition stereographic-circline-rep where
stereographic-circline-rep  $\alpha$  =
  (let (a, b, c, d) = Rep-plane-vec  $\alpha$  in
    Abs-circline-mat (cor ((c+d)/2), ((cor a+ii* cor b)/2), ((cor a-ii*cor
    b)/2), cor ((d-c)/2)))

lemma stereographic-circline-rep-Rep:
  Rep-circline-mat (stereographic-circline-rep  $\alpha$ ) =
  (let (a, b, c, d) = Rep-plane-vec  $\alpha$  in
    (cor ((c+d)/2), ((cor a+ii* cor b)/2), ((cor a-ii*cor b)/2), cor
    ((d-c)/2)))
proof-
  obtain a b c d where AA: (a, b, c, d) = Rep-plane-vec  $\alpha$ 
  by (cases Rep-plane-vec  $\alpha$ ) auto
  let ?M = (cor ((c+d)/2), ((cor a+ii* cor b)/2), ((cor a-ii*cor b)/2), cor
  ((d-c)/2))
  have ?M  $\in$  {M. hermitean M  $\wedge$  M  $\neq$  mat-zero}
  using Rep-plane-vec[of  $\alpha$ ] AA
  by (auto simp add: hermitean-def mat-adj-def mat-cnj-def complex-cnj complex-of-real-def)
  thus ?thesis
  using AA[symmetric]
  by (simp add: Abs-circline-mat-inverse stereographic-circline-rep-def)

```

qed

```
lift-definition stereographic-circline :: plane ⇒ circline is stereographic-circline-rep
proof-
  fix α1 α2
  assume plane-vec-eq α1 α2
  thus circline-mat-eq (stereographic-circline-rep α1) (stereographic-circline-rep α2)
    apply (cases Rep-plane-vec α2, cases Rep-plane-vec α1)
    apply (auto simp add: plane-vec-eq-def stereographic-circline-rep-Rep)
    apply (rule-tac x=k in exI, simp add: field-simps)
    by (metis (hide-lams, mono-tags) comm-semiring-1-class.normalize-semiring-rules(19)
      complex-of-real-mult-Complex mult-zero-right)
qed

lemma stereographic-circline-inv-stereographic-circline:
  stereographic-circline ∘ inv-stereographic-circline = id
proof (rule ext, simp)
  fix H
  show stereographic-circline (inv-stereographic-circline H) = H
  proof transfer
    fix H
    obtain A B C D where HH: Rep-circline-mat H = (A, B, C, D)
      by (cases Rep-circline-mat H) auto
    have is-real A is-real D C = conj B
      using HH Rep-circline-mat[of H] hermitean-elems[of A B C D]
      by auto
    thus circline-mat-eq (stereographic-circline-rep (inv-stereographic-circline-rep
      H)) H
      using HH
      apply (simp add: stereographic-circline-rep-Rep)
      apply (rule-tac x=1 in exI)
      apply (auto simp add: complex-of-real-Re of-real-numeral)
      apply (cases B, simp)
      apply (cases B, simp add: complex-of-real-def, metis Im.simps Re.simps
        comm-semiring-1-class.normalize-semiring-rules(4) complex-diff-def complex-minus-def
        complex-of-real-add-Complex complex-of-real-def minus-zero monoid-add-class.add.right-neutral
        one-add-one)
      done
  qed
qed

lemma [simp]: Im (z / 2) = Im z / 2
by (subst Im-divide-real, auto)

lemma [simp]: (Complex a b) / 2 = Complex (a/2) (b/2)
by (subst complex-eq-iff) auto

lemma [simp]: Complex 2 0 = 2
```

```

by simp

lemma inv-stereographic-circline-stereographic-circline:
  inv-stereographic-circline ∘ stereographic-circline = id
proof (rule ext, simp)
  fix α
  show inv-stereographic-circline (stereographic-circline α) = α
  proof transfer
    fix α
    obtain a b c d where AA: Rep-plane-vec α = (a, b, c, d)
      by (cases Rep-plane-vec α) auto
    thus plane-vec-eq (inv-stereographic-circline-rep (stereographic-circline-rep α))
      α
      using AA
      by (simp add: plane-vec-eq-def stereographic-circline-rep-Rep) (rule-tac x=1
      in exI, auto simp add: field-simps complex-of-real-def)
    qed
  qed

lemma stereographic-sphere-circle-set'':
  on-sphere-circle (inv-stereographic-circline H) z ↔ on-circline H (stereographic
  z)
proof
  assume on-sphere-circle (inv-stereographic-circline H) z
  thus on-circline H (stereographic z)
  proof transfer
    fix M H
    obtain A B C D where HH: Rep-circline-mat H = (A, B, C, D)
      by (cases Rep-circline-mat H) auto
    have *: is-real A is-real D C = cnj B
      using Rep-circline-mat[of H] HH hermitean-elems[of A B C D]
      by auto
    obtain x y z where MM: Rep-riemann-sphere M = (x, y, z)
      by (cases Rep-riemann-sphere M) auto
    assume **: on-sphere-circle-rep (inv-stereographic-circline-rep H) M
    show on-circline-rep H (stereographic-coords M)
    proof (cases z=1)
      case True
      hence x = 0 y = 0
        using MM Rep-riemann-sphere[of M]
        by auto
      thus ?thesis
        using *** HH MM ⟨z=1⟩
        by (cases A, simp add: on-circline-rep-def stereographic-coords-rep on-sphere-circle-rep-def
        vec-cnj-def Let-def)
    next
      case False
      hence Re A*(1+z) + 2*Re B*x + 2*Im B*y + Re D*(1-z) = 0
        using *** HH MM
    qed
  qed
qed

```

```

by (simp add: on-sphere-circle-rep-def Let-def field-simps)
hence (Re A*(1+z) + 2*Re B*x + 2*Im B*y + Re D*(1-z))*(1-z) = 0
  by simp
  hence Re A*(1+z)*(1-z) + 2*Re B*x*(1-z) + 2*Im B*y*(1-z) + Re
D*(1-z)*(1-z) = 0
  by (simp add: field-simps)
moreover
have x*x+y*y = (1+z)*(1-z)
  using MM Rep-riemann-sphere[of M]
  by (simp add: field-simps)
ultimately
have Re A*(x*x+y*y) + 2*Re B*x*(1-z) + 2*Im B*y*(1-z) + Re
D*(1-z)*(1-z) = 0
  by simp
hence (x * Re A + (1 - z) * Re B) * x - (- (y * Re A) + - ((1 - z) *
Im B)) * y + (x * Re B + y * Im B + (1 - z) * Re D) * (1 - z) = 0
  by (simp add: field-simps)
thus ?thesis
  using `z ≠ 1` HH MM *(Re A*(1+z) + 2*Re B*x + 2*Im B*y + Re
D*(1-z) = 0)
  apply (simp add: on-circline-rep-def stereographic-coords-rep Let-def vec-cnj-def
complex-cnj)
    apply (subst complex-eq-iff)
    apply (simp add: field-simps)
    done
qed
qed
next
assume on-circline H (stereographic z)
thus on-sphere-circle (inv-stereographic-circline H) z
proof transfer
fix H M
fix M H
obtain A B C D where HH: Rep-circline-mat H = (A, B, C, D)
  by (cases Rep-circline-mat H) auto
have *: is-real A is-real D C = cnj B
  using Rep-circline-mat[of H] HH hermitean-elems[of A B C D]
  by auto
obtain x y z where MM: Rep-riemann-sphere M = (x, y, z)
  by (cases Rep-riemann-sphere M) auto
assume **: on-circline-rep H (stereographic-coords M)
show on-sphere-circle-rep (inv-stereographic-circline-rep H) M
proof (cases z=1)
  case True
  hence x = 0 y = 0
    using MM Rep-riemann-sphere[of M]
    by auto
  thus ?thesis
    using HH MM ** `z = 1`

```

```

by (simp add: on-sphere-circle-rep-def on-circline-rep-def Let-def vec-cnj-def stereographic-coords-rep)
next
case False
hence  $(x * \operatorname{Re} A + (1 - z) * \operatorname{Re} B) * x - (-(\operatorname{y} * \operatorname{Re} A) + -((1 - z) * \operatorname{Im} B)) * y + (x * \operatorname{Re} B + y * \operatorname{Im} B + (1 - z) * \operatorname{Re} D) * (1 - z) = 0$ 
using HH MM ***
by (simp add: on-circline-rep-def Let-def vec-cnj-def stereographic-coords-rep complex-eq-iff)
hence  $\operatorname{Re} A * (x * x + y * y) + 2 * \operatorname{Re} B * x * (1 - z) + 2 * \operatorname{Im} B * y * (1 - z) + \operatorname{Re} D * (1 - z) * (1 - z) = 0$ 
by (simp add: field-simps)
moreover
have  $x * x + y * y = (1 + z) * (1 - z)$ 
using MM Rep-riemann-sphere[of M]
by (simp add: field-simps)
ultimately
have  $\operatorname{Re} A * (1 + z) * (1 - z) + 2 * \operatorname{Re} B * x * (1 - z) + 2 * \operatorname{Im} B * y * (1 - z) + \operatorname{Re} D * (1 - z) * (1 - z) = 0$ 
by simp
hence  $(\operatorname{Re} A * (1 + z) + 2 * \operatorname{Re} B * x + 2 * \operatorname{Im} B * y + \operatorname{Re} D * (1 - z)) * (1 - z) = 0$ 
by (simp add: field-simps)
hence  $\operatorname{Re} A * (1 + z) + 2 * \operatorname{Re} B * x + 2 * \operatorname{Im} B * y + \operatorname{Re} D * (1 - z) = 0$ 
using  $(z \neq 1)$ 
by simp
thus ?thesis
using MM HH *
by (simp add: on-sphere-circle-rep-def field-simps)
qed
qed
qed

lemma stereographic-sphere-circle-set':
stereographic `sphere-circle-set (inv-stereographic-circline H) = circline-set H
unfolding sphere-circle-set-def circline-set-def
apply safe
proof-
fix x
assume on-sphere-circle (inv-stereographic-circline H) x
thus on-circline H (stereographic x)
using stereographic-sphere-circle-set''
by simp
next
fix x
assume on-circline H x
show x ∈ stereographic ` {z. on-sphere-circle (inv-stereographic-circline H) z}
proof
show x = stereographic (inv-stereographic x)
by (simp add: stereographic-inv-stereographic)

```

```

next
  show inv-stereographic  $x \in \{z. \text{on-sphere-circle} (\text{inv-stereographic-circline } H)$ 
 $\}$ 
    using stereographic-sphere-circle-set'[of  $H$  inv-stereographic  $x$ ] (on-circline  $H$ 
 $x$ )
      by (simp add: stereographic-inv-stereographic)
qed
qed

lemma stereographic-sphere-circle-set:
  shows stereographic `sphere-circle-set  $H = \text{circline-set} (\text{stereographic-circline } H)$ 
using stereographic-sphere-circle-set'[of stereographic-circline  $H$ ]
using inv-stereographic-circline-stereographic-circline
unfolding comp-def
by (metis id-apply)

lemma bij stereographic-circline
using stereographic-circline-inv-stereographic-circline inv-stereographic-circline-stereographic-circline
by (metis bij-def image-compose inj-iff inj-imp-surj-inv inj-on-imageI2 inv-id surj-id
surj-iff)

lemma bij inv-stereographic-circline
using stereographic-circline-inv-stereographic-circline inv-stereographic-circline-stereographic-circline
by (metis bij-def image-compose inj-iff inj-imp-surj-inv inj-on-imageI2 inv-id surj-id
surj-iff)

```

11.3 Some special circlines

Unit circle

definition unit-circle-rep **where**
[*simp*]: unit-circle-rep = Abs-circline-mat (1, 0, 0, -1)

lemma [*simp*]: Rep-circline-mat (Abs-circline-mat (1, 0, 0, -1)) = (1, 0, 0, -1)
by (auto *simp add: Abs-circline-mat-inverse hermitean-def mat-adj-def mat-cnj-def*)

lemma [*simp*]: Rep-circline-mat unit-circle-rep = (1, 0, 0, -1)
by *simp*

lift-definition unit-circle :: circline **is** unit-circle-rep
done

lemma one-on-unit-circle: $1_h \in \text{circline-set unit-circle}$
unfolding circline-set-def
by (simp, transfer, *simp add: on-circline-rep-def Let-def vec-cnj-def*)

x-axis

definition x-axis-rep **where** x-axis-rep = Abs-circline-mat (0, ii, -ii, 0)
lift-definition x-axis :: circline **is** x-axis-rep
done

lemma [simp]: $\text{Rep-circline-mat}(\text{Abs-circline-mat}(0, ii, -ii, 0)) = (0, ii, -ii, 0)$
using $\text{Abs-circline-mat-inverse}[\text{of } (0, ii, -ii, 0)]$
by (simp add: hermitean-def mat-adj-def mat-cnj-def complex-cnj)

lemma [simp]: $\text{Rep-circline-mat} \text{x-axis-rep} = (0, ii, -ii, 0)$
unfolding x-axis-rep-def
by simp

lemma [simp]: $0_h \in \text{circline-set} \text{x-axis}$ $1_h \in \text{circline-set} \text{x-axis}$ $\infty_h \in \text{circline-set} \text{x-axis}$
unfolding circline-set-def
by auto (transfer, simp add: on-circline-rep-def Let-def vec-cnj-def) +

Point 0_h as a circline

definition $\text{circline-point-0h-rep}$ **where** $\text{circline-point-0h-rep} = \text{Abs-circline-mat}(1, 0, 0, 0)$

lift-definition $\text{circline-point-0h} :: \text{circline}$ **is** $\text{circline-point-0h-rep}$
done

lemma [simp]: $\text{Rep-circline-mat}(\text{Abs-circline-mat}(1, 0, 0, 0)) = (1, 0, 0, 0)$
using $\text{Abs-circline-mat-inverse}$
by (simp add: hermitean-def mat-adj-def mat-cnj-def)

lemma [simp]: $\text{Rep-circline-mat} \text{circline-point-0h-rep} = (1, 0, 0, 0)$
unfolding $\text{circline-point-0h-rep-def}$
by simp

imaginary unit circle

definition $\text{imag-unit-circle-rep}$ **where**
[**simp**]: $\text{imag-unit-circle-rep} = \text{Abs-circline-mat}(1, 0, 0, 1)$

lemma [simp]: $\text{Rep-circline-mat}(\text{Abs-circline-mat}(1, 0, 0, 1)) = (1, 0, 0, 1)$
by (auto simp add: Abs-circline-mat-inverse hermitean-def mat-adj-def mat-cnj-def)

lemma [simp]: $\text{Rep-circline-mat} \text{imag-unit-circle-rep} = (1, 0, 0, 1)$
by simp

lift-definition $\text{imag-unit-circle} :: \text{circline}$ **is** $\text{imag-unit-circle-rep}$
done

11.4 Moebius action on circlines

definition $\text{moebius-circline-rep} :: \text{moebius-mat} \Rightarrow \text{circline-mat} \Rightarrow \text{circline-mat}$ **where**

$\text{moebius-circline-rep} M H =$
(let $M = \text{Rep-moebius-mat} M$;

$H = \text{Rep-circline-mat } H$
 in $\text{Abs-circline-mat} (\text{congruence} (\text{mat-inv } M) H))$

lemma [simp]: $\text{Rep-circline-mat} (\text{Abs-circline-mat} (\text{congruence} (\text{mat-inv} (\text{Rep-moebius-mat } M)) (\text{Rep-circline-mat } H))) = \text{congruence} (\text{mat-inv} (\text{Rep-moebius-mat } M)) (\text{Rep-circline-mat } H)$
proof (rule *Abs-circline-mat-inverse, safe*)
show *hermitean* ($\text{congruence} (\text{mat-inv} (\text{Rep-moebius-mat } M)) (\text{Rep-circline-mat } H))$
using *Rep-circline-mat*[of H]
using *hermitean-congruence*
by *simp*
next
assume $\text{congruence} (\text{mat-inv} (\text{Rep-moebius-mat } M)) (\text{Rep-circline-mat } H) = \text{mat-zero}$
thus *False*
using *Rep-circline-mat*[of H] *Rep-moebius-mat*[of M] *mat-det-inv*
using *congruence-nonzero*
by *auto*
qed

lemma *moebius-circline-rep-Rep* [simp]: $\text{Rep-circline-mat} (\text{moebius-circline-rep } M H) = \text{congruence} (\text{mat-inv} (\text{Rep-moebius-mat } M)) (\text{Rep-circline-mat } H)$
by (simp add: *moebius-circline-rep-def Let-def*)

lift-definition *moebius-circline* :: *moebius* \Rightarrow *circline* \Rightarrow *circline* **is** *moebius-circline-rep*
proof–
fix $M M' H H'$
assume *moebius-mat-eq* $M M' \text{ circline-mat-eq } H H'$
thus *circline-mat-eq* (*moebius-circline-rep* $M H$) (*moebius-circline-rep* $M' H')$
by (auto simp add: *mat-inv-mult-sm complex-cnj*) (rule-tac $x=ka / Re (k * cnj k)$ in *exI*, auto simp add: *complex-mult-cnj-cmod power2-eq-square*)
qed

lemma *moebius-preserve-circline-type*:
shows *circline-type* (*moebius-circline* $M H$) = *circline-type* H
proof (*transfer*)
fix $M H$
show *circline-type-rep* (*moebius-circline-rep* $M H$) = *circline-type-rep* H
unfolding *circline-type-rep-def Let-def*
apply *simp*
using *Re-det-sgn-congruence*[of *Rep-circline-mat* H *mat-inv* (*Rep-moebius-mat* M)]
using *Rep-circline-mat*[of H] *Rep-moebius-mat*[of M] *mat-det-inv*[of *Rep-moebius-mat* M]
by *simp*
qed

lemma *moebius-circline-rep*:

```

shows moebius-pt-rep M ` {z. on-circline-rep H z} = {z. on-circline-rep (moebius-circline-rep
M H) z}
proof (safe)
fix z
let ?M = Rep-moebius-mat M
let ?H = Rep-circline-mat H
let ?z = Rep-homo-coords z
let ?H' = Rep-circline-mat H'
let ?z' = Rep-moebius-mat M *mv Rep-homo-coords z
let ?H'' = mat-adj (mat-inv ?M) *mm ?H *mm (mat-inv ?M)
assume on-circline-rep H z
hence quad-form ?z ?H = 0
by (simp add: on-circline-rep-def Let-def)
hence quad-form ?z' ?H'' = 0
using quad-form-congruence[of ?M ?z ?H] Rep-moebius-mat[of M]
by simp
thus on-circline-rep (moebius-circline-rep M H) (moebius-pt-rep M z)
by (auto simp add: moebius-circline-rep-def on-circline-rep-def moebius-pt-rep-def
Let-def)
next
fix z
let ?z = Rep-homo-coords z
let ?M = Rep-moebius-mat M
let ?H = Rep-circline-mat H
let ?iM = mat-inv ?M
let ?z' = mat-inv ?M *mv ?z

assume on-circline-rep (moebius-circline-rep M H) z
hence quad-form ?z (congruence (mat-inv ?M) ?H) = 0
unfolding on-circline-rep-def Let-def
by simp

have ?z' ≠ (0, 0)
using Rep-homo-coords[of z] mult-mv-nonzero[of ?z ?iM] Rep-moebius-mat[of
M] mat-det-inv[of ?M]
by simp
hence *: Rep-homo-coords (Abs-homo-coords ?z') = ?z'
by (simp add: Abs-homo-coords-inverse)

show z ∈ moebius-pt-rep M ` {z. on-circline-rep H z}
proof
show z = moebius-pt-rep M (Abs-homo-coords ?z')
using * Rep-moebius-mat[of M] eye-mv-l[of ?z]
unfolding moebius-pt-rep-def Let-def
by (simp add: mat-inv-r Rep-homo-coords-inverse)
next
have Rep-moebius-mat M *mm mat-inv (Rep-moebius-mat M) *mv Rep-homo-coords
z = ?z
using Rep-moebius-mat[of M]

```

```

by (subst mat-inv-r) (auto simp add: simp del: eye-def)
thus Abs-homo-coords ?z' ∈ {z. on-circline-rep H z}
  using *
  using <quad-form ?z (congruence (mat-inv ?M) ?H) = 0> Rep-moebius-mat[of
M]
  by (auto simp add: on-circline-rep-def Let-def simp del: quad-form-def) (subst
quad-form-congruence[of ?M ?iM *mv ?z ?H, symmetric], auto)
  qed
qed

lemma moebius-circline-set:
  shows moebius-pt M ` circline-set H = circline-set (moebius-circline M H) (is
?lhs = ?rhs)
proof
  show ?lhs ⊆ ?rhs
  proof (safe)
    fix z::complex-homo
    assume z ∈ circline-set H
    thus moebius-pt M z ∈ circline-set (moebius-circline M H)
      unfolding circline-set-def
      using moebius-circline-rep
      by simp (transfer, auto)
  qed
next
  show ?rhs ⊆ ?lhs
  proof
    fix z
    assume z ∈ circline-set (moebius-circline M H)
    thus z ∈ moebius-pt M ` circline-set H
      using assms
      unfolding circline-set-def
      apply (simp add: image-def)
    proof (transfer)
      fix M H z
      assume on-circline-rep (moebius-circline-rep M H) z
      then obtain z' where on-circline-rep H z' z = moebius-pt-rep M z'
        using moebius-circline-rep[of M H]
        by auto
      thus ∃ z'. on-circline-rep H z' ∧ z ≈ moebius-pt-rep M z'
        by (rule-tac x=z' in exI, simp, rule-tac x=1 in exI, simp)
    qed
  qed
qed

lemma
  inj-moebius-circline: inj (moebius-circline M)
unfolding inj-on-def
proof (safe)
  fix H H'

```

```

assume moebius-circline M H = moebius-circline M H'
thus H = H'
proof (transfer)
  fix M H H'
  let ?M = Rep-moebius-mat M
  let ?iM = mat-inv ?M
  let ?H = Rep-circline-mat H and ?H' = Rep-circline-mat H'
  assume circline-mat-eq (moebius-circline-rep M H) (moebius-circline-rep M H')
  then obtain k where congruence ?iM ?H' = congruence ?iM (cor k *sm ?H)
  k ≠ 0
  by auto
  thus circline-mat-eq H H'
  using Rep-moebius-mat[of M] inj-congruence[of ?iM ?H' cor k *sm ?H]
  mat-det-inv[of ?M]
  by auto
  qed
qed

lemma [simp]:
  moebius-circline id-moebius H = H
proof transfer
  fix H
  show circline-mat-eq (moebius-circline-rep id-moebius-rep H) H
  by (cases Rep-circline-mat H, simp) (rule-tac x=1 in exI, simp add: mat-adj-def
  mat-cnj-def)
qed

lemma moebius-circline-comp:
  moebius-circline M1 (moebius-circline M2 H) = moebius-circline (moebius-comp
  M1 M2) H
proof (transfer)
  fix M1 M2 H
  show circline-mat-eq (moebius-circline-rep M1 (moebius-circline-rep M2 H))
  (moebius-circline-rep (moebius-comp-rep M1 M2) H)
  using congruence-congruence Rep-moebius-mat[of M1] Rep-moebius-mat[of M2]
  by (simp add: mat-inv-mult-mm, rule-tac x=1 in exI, simp)
qed

lemma moebius-circline-comp-inv [simp]:
  moebius-circline (moebius-inv M) (moebius-circline M H) = H
  by (subst moebius-circline-comp) simp

lemma moebius-circline-comp-inv' [simp]:
  moebius-circline M (moebius-circline (moebius-inv M) H) = H
  by (subst moebius-circline-comp) simp

lemma
  moebius-circline-set-mem:

```

```

moebius-pt M z ∈ circline-set (moebius-circline M H) ←→ z ∈ circline-set H
using moebius-circline-set[of M H, symmetric] bij-moebius-pt[of M]
by (auto simp add: bij-def inj-on-def)

```

11.5 Conjugation, reciprocation and inversion of circlines

Conjugation of circlines

definition circline-cnj-rep **where**

```
circline-cnj-rep H = Abs-circline-mat (mat-cnj (Rep-circline-mat H))
```

```

lemma [simp]: Rep-circline-mat (Abs-circline-mat (mat-cnj (Rep-circline-mat H)))  

= mat-cnj (Rep-circline-mat H)  

using Rep-circline-mat[of H] hermitean-mat-cnj nonzero-mat-cnj  

by (auto simp add: Abs-circline-mat-inverse)

```

```

lemma [simp]: Rep-circline-mat (circline-cnj-rep H) = mat-cnj (Rep-circline-mat H)  

by (simp add: circline-cnj-rep-def)

```

lift-definition circline-cnj :: circline ⇒ circline **is** circline-cnj-rep

proof –

```
fix H H'  

assume circline-mat-eq H H'  

thus circline-mat-eq (circline-cnj-rep H) (circline-cnj-rep H')  

using Rep-circline-mat[of H] Rep-circline-mat[of H']  

by auto
qed
```

lemma cnj-homo-circline-set':

```
shows cnj-homo ` circline-set H ⊆ circline-set (circline-cnj H)
```

proof (safe)

```
fix z  

assume z ∈ circline-set H
```

```
thus cnj-homo z ∈ circline-set (circline-cnj H)
```

unfolding circline-set-def

apply simp

proof (transfer)

fix z H

assume on-circline-rep H z

obtain z1 z2 **where** zz: Rep-homo-coords z = (z1, z2)

by (rule obtain-homo-coords)

have (cnj z1, cnj z2) *_{vm} Rep-circline-mat H *_{vv} (z1, z2) = 0

using <on-circline-rep H z> zz

unfolding on-circline-rep-def Let-def

by (simp add: vec-cnj-def)

hence cnj ((cnj z1, cnj z2) *_{vm} Rep-circline-mat H *_{vv} (z1, z2)) = 0

by simp

hence (z1, z2) *_{vm} mat-cnj (Rep-circline-mat H) *_{vv} (cnj z1, cnj z2) = 0

```

by (subst (asm) cnj-mult-vv) (cases Rep-circline-mat H, simp add: vec-cnj-def
mat-cnj-def complex-cnj)
  thus on-circline-rep (circline-cnj-rep H) (cnj-homo-coords z)
    unfolding on-circline-rep-def Let-def
    using zz
    by (simp add: vec-cnj-def)
  qed
qed

lemma [simp]: circline-cnj (circline-cnj H) = H
by (transfer) (auto simp add: circline-cnj-rep-def Rep-circline-mat-inverse, rule-tac
x=1 in exI, simp)

lemma cnj-homo-circline-set:
  shows cnj-homo ` circline-set H = circline-set (circline-cnj H) (is ?lhs = ?rhs)
proof (safe)
  fix z
  assume z ∈ circline-set (circline-cnj H)
  show z ∈ cnj-homo ` circline-set H
  proof
    show z = cnj-homo (cnj-homo z)
    by simp
  next
    show cnj-homo z ∈ circline-set H
    using ⟨z ∈ circline-set (circline-cnj H)⟩
    using cnj-homo-circline-set'[of circline-cnj H]
    by auto
  qed
  next
    fix z
    assume z ∈ circline-set H
    thus cnj-homo z ∈ circline-set (circline-cnj H)
      using cnj-homo-circline-set'[of H]
      by auto
  qed

```

Reciprocal and inversion of circlines

```

definition circline-swap-AD-rep where
  circline-swap-AD-rep H =
    (let (A, B, C, D) = Rep-circline-mat H
     in Abs-circline-mat (D, B, C, A))

```

```

lemma
  shows [simp]: Rep-circline-mat (circline-swap-AD-rep H) = (let (A, B, C, D)
  = Rep-circline-mat H in (D, B, C, A))
proof-
  obtain A B C D where hh: Rep-circline-mat H = (A, B, C, D)
  by (cases Rep-circline-mat H) auto
  have hermitean (D, B, C, A) ∧ (D, B, C, A) ≠ mat-zero

```

```

using Rep-circline-mat[of H] hh
by (auto simp add: hermitean-def mat-adj-def mat-cnj-def)
thus ?thesis
  using hh
  unfolding circline-swap-AD-rep-def Let-def
  by (cases Rep-circline-mat H) (simp add: Abs-circline-mat-inverse)
qed

lift-definition circline-swap-AD :: circline ⇒ circline is circline-swap-AD-rep
proof-
  fix H H' :: circline-mat
  assume circline-mat-eq H H'
  thus circline-mat-eq (circline-swap-AD-rep H) (circline-swap-AD-rep H')
    by (cases Rep-circline-mat H, cases Rep-circline-mat H') auto
qed

lemma reciprocal-circline-set:
  shows reciprocal-homo ` circline-set H = circline-set ((circline-cnj ∘ circline-swap-AD)
H)
  proof (subst reciprocal-moebius, subst moebius-circline-set)
    have moebius-circline reciprocal-moebius H = (circline-cnj ∘ circline-swap-AD)
H
      unfolding reciprocal-moebius-def
    proof (transfer)
      fix H :: circline-mat
      obtain A B C D where H: Rep-circline-mat H = (A, B, C, D)
        by (cases Rep-circline-mat H) blast
      thus circline-mat-eq (moebius-circline-rep (mk-moebius-rep 0 1 1 0) H) ((circline-cnj-rep
        ∘ circline-swap-AD-rep) H)
        using Rep-circline-mat[of H]
        by (simp add: mat-adj-def mat-cnj-def hermitean-def) (rule-tac x=1 in exI,
simp)
      qed
      thus circline-set (moebius-circline reciprocal-moebius H) = circline-set ((circline-cnj
        ∘ circline-swap-AD) H)
        by simp
    qed
  qed

lemma inversion-circline-set:
  shows inversion-homo ` circline-set H = circline-set (circline-swap-AD H)
  unfolding inversion-homo-def image-comp
  by (subst reciprocal-circline-set, subst cnj-homo-circline-set, rule arg-cong[where
f=circline-set]) simp

```

11.6 Circline uniqueness

11.6.1 Zero type circline uniqueness

```

lemma unique-circline-type-zero-0h':
  shows (circline-type circline-point-0h = 0 ∧ 0h ∈ circline-set circline-point-0h)

```

```

 $\wedge$ 
   $(\forall H. \text{circline-type } H = 0 \wedge 0_h \in \text{circline-set } H \longrightarrow H = \text{circline-point-0h})$ 
unfolding circline-set-def
proof (safe)
  show circline-type circline-point-0h = 0
    by (transfer) (simp add: circline-type-rep-def circline-point-0h-rep-def)
next
  show on-circline circline-point-0h 0_h
    by (transfer) (simp add: on-circline-rep-def Let-def vec-cnj-def)
next
  fix H
  assume circline-type H = 0 on-circline H 0_h
  thus H = circline-point-0h
  proof (transfer)
    fix H
    obtain A B C D where HH: Rep-circline-mat H = (A, B, C, D)
      by (cases Rep-circline-mat H) auto
    hence *: C = cnj B is-real A
      using Rep-circline-mat[of H] hermitean-elems[of A B C D]
      by auto
    assume circline-type-rep H = 0 on-circline-rep H zero-homo-rep
    thus circline-mat-eq H circline-point-0h-rep
      using * Rep-circline-mat[of H] HH
      by (simp add: on-circline-rep-def Let-def Abs-circline-mat-inverse vec-cnj-def
circline-type-rep-def sgn-minus sgn-mult sgn-zero-iff)
        (rule-tac x=1/Re A in exI, cases A, cases B, simp add: complex-of-real-Re
sgn-zero-iff)
    qed
  qed

lemma unique-circline-type-zero-0h:
  shows  $\exists! H. \text{circline-type } H = 0 \wedge 0_h \in \text{circline-set } H$ 
  using unique-circline-type-zero-0h'
  by auto

lemma unique-circline-type-zero:
  shows  $\exists! H. \text{circline-type } H = 0 \wedge z \in \text{circline-set } H$ 
proof-
  obtain M where ++: moebius-pt M z = 0_h
    using ex-moebius-1[of z]
    by auto
  have  $+++: z = \text{moebius-pt}(\text{moebius-inv } M) 0_h$ 
    by (subst ++[symmetric]) simp
  then obtain H0 where *: circline-type H0 = 0  $\wedge 0_h \in \text{circline-set } H0$  and
   $**: \forall H'. \text{circline-type } H' = 0 \wedge 0_h \in \text{circline-set } H' \longrightarrow H' = H0$ 
    using unique-circline-type-zero-0h
    by auto
  let ?H' = moebius-circline(moebius-inv M) H0
  show ?thesis

```

```

unfolding Ex1-def
using * +++
proof (rule-tac x=?H' in exI, simp add: moebius-preserve-circline-type moebius-circline-set[symmetric], safe)
  fix H'
  assume circline-type H' = 0 moebius-pt (moebius-inv M) 0_h ∈ circline-set H'
  hence 0_h ∈ circline-set (moebius-circline M H')
    by (metis + + + + imageI moebius-circline-set)
  hence moebius-circline M H' = H0
    using **[rule-format, of moebius-circline M H']
    using moebius-preserve-circline-type[of M H'] ⟨circline-type H' = 0⟩
      by simp
    thus H' = moebius-circline (moebius-inv M) H0
      by auto
  qed
qed

```

11.6.2 Negative type circline uniqueness

```

lemma unique-circline-01inf':
  0_h ∈ circline-set x-axis ∧ 1_h ∈ circline-set x-axis ∧ ∞_h ∈ circline-set x-axis ∧
  (∀ H. 0_h ∈ circline-set H ∧ 1_h ∈ circline-set H ∧ ∞_h ∈ circline-set H → H
  = x-axis)
proof safe
  fix H
  assume 0_h ∈ circline-set H 1_h ∈ circline-set H ∞_h ∈ circline-set H
  thus H = x-axis
    unfolding circline-set-def
    apply simp
    proof (transfer)
      fix H
      obtain A B C D where HH: Rep-circline-mat H = (A, B, C, D)
        by (cases Rep-circline-mat H) auto
      have *: C = cnj B A = 0 ∧ D = 0 → B ≠ 0
        using hermitean-elems[of A B C D] Rep-circline-mat[of H] HH
        by auto
      obtain Bx By where B = Complex Bx By
        by (cases B) auto
      assume on-circline-rep H zero-homo-rep on-circline-rep H one-homo-rep on-circline-rep
      H inf-homo-rep
      thus circline-mat-eq H x-axis-rep
        using * HH ⟨C = cnj B⟩ ⟨B = Complex Bx By⟩
        by (simp add: on-circline-rep-def Let-def mk-circline-rep-def Abs-circline-mat-inverse
        vec-cnj-def complex-of-real-def i-def, rule-tac x=1/By in exI, auto )
      qed
qed simp-all

lemma unique-circline-set:
  assumes A ≠ B A ≠ C B ≠ C

```

shows $\exists! H. A \in \text{circline-set } H \wedge B \in \text{circline-set } H \wedge C \in \text{circline-set } H$
proof—
let $?P = \lambda A B C. A \neq B \wedge A \neq C \wedge B \neq C \longrightarrow (\exists! H. A \in \text{circline-set } H \wedge B \in \text{circline-set } H \wedge C \in \text{circline-set } H)$
have $?P A B C$
proof (rule *wlog-moebius-01inf*[of $?P$])
fix $M a b c$
let $?M = \text{moebius-pt } M$
assume $?P a b c$
show $?P (?M a) (?M b) (?M c)$
proof
assume $?M a \neq ?M b \wedge ?M a \neq ?M c \wedge ?M b \neq ?M c$
hence $a \neq b b \neq c a \neq c$
using *bij-moebius-pt*[of M]
by (auto simp add: *bij-def inj-on-def*)
hence $\exists! H. a \in \text{circline-set } H \wedge b \in \text{circline-set } H \wedge c \in \text{circline-set } H$
using $\langle ?P a b c \rangle$
by *simp*
then obtain H **where**
 $*: a \in \text{circline-set } H \wedge b \in \text{circline-set } H \wedge c \in \text{circline-set } H$ **and**
 $**: \forall H'. a \in \text{circline-set } H' \wedge b \in \text{circline-set } H' \wedge c \in \text{circline-set } H' \longrightarrow H' = H$
unfolding *Ex1-def*
by *auto*
let $?H' = \text{moebius-circline } M H$
show $\exists! H. ?M a \in \text{circline-set } H \wedge \text{moebius-pt } M b \in \text{circline-set } H \wedge \text{moebius-pt } M c \in \text{circline-set } H$
unfolding *Ex1-def*
proof (rule-tac $x=?H'$ in *exI*, rule)
show $?M a \in \text{circline-set } ?H' \wedge ?M b \in \text{circline-set } ?H' \wedge ?M c \in \text{circline-set } ?H'$
using * *moebius-circline-set-mem*[of $M - H$]
by *blast*
next
show $\forall H'. ?M a \in \text{circline-set } H' \wedge ?M b \in \text{circline-set } H' \wedge ?M c \in \text{circline-set } H' \longrightarrow H' = ?H'$
proof (safe)
fix H'
let $?iH' = \text{moebius-circline } (\text{moebius-inv } M) H'$
assume $?M a \in \text{circline-set } H' ?M b \in \text{circline-set } H' ?M c \in \text{circline-set } H'$
hence $a \in \text{circline-set } ?iH' \wedge b \in \text{circline-set } ?iH' \wedge c \in \text{circline-set } ?iH'$
using *moebius-circline-set-mem*[of $M - ?iH'$, simplified]
by *simp*
hence $H = ?iH'$
using **
by *simp*
thus $H' = \text{moebius-circline } M H$
by *simp*

```

qed
qed
qed
next
show ?P 0h 1h ∞h
  using unique-circline-01inf'
  unfolding Ex1-def
  by (safe, rule-tac x=x-axis in exI) auto
qed fact+
thus ?thesis
  using assms
  by simp
qed

```

11.7 Circline set cardinality

11.7.1 Diagonal circlines

definition circline-diag-rep **where**

circline-diag-rep $H \longleftrightarrow$ mat-diagonal (Rep-circline-mat H)

lemma [simp]: mat-diagonal $H \longleftrightarrow (\text{let } (A, B, C, D) = H \text{ in } B = 0 \wedge C = 0)$
by (cases H) simp

lift-definition circline-diag :: circline \Rightarrow bool **is** circline-diag-rep
by (auto simp add: circline-diag-rep-def)

lemma det-zero-trace-zero:

assumes mat-det $A = 0$ mat-trace $A = (0::complex)$ hermitean A

shows $A = \text{mat-zero}$

using assms

proof-

{

fix $a d c$

assume $a * d = \text{cnj } c * c a + d = 0$ $\text{cnj } a = a$

from $\langle a + d = 0 \rangle$ have $d = -a$

by (metis add-eq-0-iff)

hence $-(\text{cor}(\text{Re } a))^2 = (\text{cor}(\text{cmod } c))^2$

using ⟨cnj a = a⟩ eq-cnj-iff-real[of a]

using ⟨a*d = cnj c * c⟩

using complex-mult-cnj-cmod[of cnj c]

by (simp add: complex-of-real-Re power2-eq-square)

hence $-(\text{Re } a)^2 \geq 0$

using zero-le-power2[of cmod c]

by (metis Re-complex-of-real cor-squared of-real-minus)

hence $a = 0$

using zero-le-power2[of Re a]

using ⟨cnj a = a⟩ eq-cnj-iff-real[of a]

by (cases a) simp

}

note * = this

```

obtain a b c d where A = (a, b, c, d)
  by (cases A) auto
thus ?thesis
  using *[of a d c] *[of d a c]
  using assms ⟨A = (a, b, c, d)⟩
  by (auto simp add: hermitean-def mat-adj-def mat-cnj-def)
qed

lemma circline-diagonalize:
  shows ∃ M H'. moebius-circline M H = H' ∧ circline-diag H'
using assms
proof transfer
fix H
obtain A B C D where HH: Rep-circline-mat H = (A, B, C, D)
  by (cases Rep-circline-mat H) auto
hence HH-elems: is-real A is-real D C = cnj B
  using hermitean-elems[of A B C D] Rep-circline-mat[of H]
  by auto
obtain M k1 k2 where *: mat-det M ≠ 0 unitary M congruence M (Rep-circline-mat
H) = (k1, 0, 0, k2) is-real k1 is-real k2
  using hermitean-diagonizable[of Rep-circline-mat H] Rep-circline-mat[of H]
  by auto
have k1 ≠ 0 ∨ k2 ≠ 0
  using ⟨congruence M (Rep-circline-mat H) = (k1, 0, 0, k2)⟩ Rep-circline-mat[of
H] congruence-nonzero[of Rep-circline-mat H M] ⟨mat-det M ≠ 0⟩
  by auto

have **: Rep-circline-mat (Abs-circline-mat (k1, 0, 0, k2)) = (k1, 0, 0, k2)
  apply (rule Abs-circline-mat-inverse)
  using ⟨is-real k1⟩ ⟨is-real k2⟩ ⟨k1 ≠ 0 ∨ k2 ≠ 0⟩
  by (auto simp add: hermitean-def mat-adj-def mat-cnj-def eq-cnj-iff-real[symmetric])

thus ∃ M H'. circline-mat-eq (moebius-circline-rep M H) H' ∧ circline-diag-rep
H'
  using * mat-det-inv[of M]
  by (rule-tac x=Abs-moebius-mat (mat-inv M) in exI, rule-tac x=Abs-circline-mat
(k1, 0, 0, k2) in exI)
    (simp add: Abs-moebius-mat-inverse circline-diag-rep-def, rule-tac x=1 in
exI, simp)
qed

lemma wlog-circline-diag:
assumes ⋀ H. circline-diag H ⟹ P H
  ⋀ M H. P H ⟹ P (moebius-circline M H)
shows P H
proof-
obtain M H' where moebius-circline M H = H' circline-diag H'
  using circline-diagonalize[of H]

```

```

    by auto
  hence  $P$  (moebius-circline  $M$   $H$ )
    using assms(1)
    by simp
  thus ?thesis
    using assms(2)[of moebius-circline  $M$   $H$  moebius-inv  $M$ ]
    by simp
qed

```

11.7.2 Zero type circline set cardinality

```

lemma circline-type-zero-card-eq1-0h:
  assumes circline-type  $H = 0$   $0_h \in \text{circline-set } H$ 
  shows circline-set  $H = \{0_h\}$ 
using assms
unfolding circline-set-def
proof(safe)
  fix  $z$ 
  assume on-circline  $H z$  circline-type  $H = 0$  on-circline  $H 0_h$ 
  hence  $H = \text{circline-point-}0h$ 
    using unique-circline-type-zero-0h'
    unfolding circline-set-def
    by simp
  thus  $z = 0_h$ 
    using (on-circline  $H z)
  proof transfer
    fix  $H z$ 
    assume circline-mat-eq  $H$  circline-point-0h-rep on-circline-rep  $H z$ 
    thus  $z \approx \text{zero-homo-rep}$ 
      using Rep-homo-coords[of z]
      by (cases Rep-homo-coords  $z$ , cases Rep-circline-mat  $H$ ) (simp add: circline-point-0h-rep-def
on-circline-rep-def Let-def vec-cnj-def, rule-tac  $x=1/b$  in exI, auto)
    qed
  qed$ 
```

lemma *bij-image-singleton*:

$\llbracket f : A = \{b\}; f a = b; \text{bij } f \rrbracket \implies A = \{a\}$

by (metis (mono-tags) *bij-betw-imp-inj-on* *image-empty* *image-insert* *inj-vimage-image-eq*)

```

lemma circline-type-zero-card-eq1:
  assumes circline-type  $H = 0$ 
  shows  $\exists z. \text{circline-set } H = \{z\}$ 
proof-
  have  $\exists z. \text{on-circline } H z$ 
  using assms
  proof transfer
    fix  $H$ 
    obtain  $A B C D$  where HH: Rep-circline-mat  $H = (A, B, C, D)$ 
    by (cases Rep-circline-mat  $H$ ) auto

```

```

hence  $C = \text{cnj } B$  is-real  $A$  is-real  $D$ 
  using Rep-circline-mat[of  $H$ ] hermitean-elems[of  $A$   $B$   $C$   $D$ ]
  by auto
assume circline-type-rep  $H = 0$ 
hence mat-det (Rep-circline-mat  $H$ ) = 0
  using Rep-circline-mat[of  $H$ ] mat-det-hermitean-real[of Rep-circline-mat  $H$ ]
  by (auto simp add: circline-type-rep-def sgn-zero-iff) (metis complex-surj
complex-zero-def)
hence  $A*D = B*C$ 
  using HH
  by simp
show Ex (on-circline-rep  $H$ )
proof (cases  $A \neq 0 \vee B \neq 0$ )
  case True
  thus ?thesis
    using HH (A*D = B*C)
    by (rule-tac x=Abs-homo-coords (-B, A) in exI) (auto simp add: on-circline-rep-def
Let-def Abs-homo-coords-inverse vec-cnj-def complex-cnj field-simps)
next
  case False
  thus ?thesis
    using HH (C = cnj B)
    by (rule-tac x=Abs-homo-coords (1, 0) in exI) (simp add: Abs-homo-coords-inverse
on-circline-rep-def Let-def vec-cnj-def)
qed
qed
then obtain  $z$  where on-circline  $H z$ 
  by auto
obtain  $M$  where moebius-pt  $M z = 0_h$ 
  using ex-moebius-1[of  $z$ ]
  by auto
hence  $0_h \in \text{circline-set} (\text{moebius-circline } M H)$ 
  using (on-circline  $H z$ )
  by (subst moebius-circline-set[of  $M H$ , symmetric]) (force simp add: circline-set-def)
hence circline-set (moebius-circline  $M H$ ) = { $0_h$ }
  using circline-type-zero-card-eq1-0h[of moebius-circline  $M H$ ] (circline-type  $H$ 
= 0)
  by (auto simp add: moebius-preserve-circline-type)
hence circline-set  $H = \{z\}$ 
  using (moebius-pt  $M z = 0_h$ )
  using bij-moebius-pt[of  $M$ ] bij-image-singleton[of moebius-pt  $M$  circline-set  $H$  -
 $z$ ]
  by (subst (asm) moebius-circline-set[symmetric]) simp
thus ?thesis
  by auto
qed

```

11.7.3 Negative type circline set cardinality

```

lemma quad-form-diagonal-iff:
  assumes k1 ≠ 0 is-real k1 is-real k2 Re k1 * Re k2 < 0
  shows quad-form (z1, 1) (k1, 0, 0, k2) = 0 ↔ (exists φ. z1 = rcis (sqrt (Re (-k2 / k1))) φ)
proof-
  have Re (-k2/k1) ≥ 0
  using ⟨Re k1 * Re k2 < 0⟩ ⟨is-real k1⟩ ⟨is-real k2⟩ ⟨k1 ≠ 0⟩
  by (auto simp add: Re-divide-real) (metis less-asym mult-neg-neg mult-pos-pos not-less zero-less-divide-iff)

  have quad-form (z1, 1) (k1, 0, 0, k2) = 0 ↔ (cor (cmod z1))² = -k2 / k1
  using assms add-eq-0-iff[of k2 k1*(cor (cmod z1))²]
  using eq-divide-imp[of k1 (cor (cmod z1))² -k2]
  by (auto simp add: vec-cnj-def field-simps complex-mult-cnj-cmod)
  also have ... ↔ (cmod z1)² = Re (-k2 / k1)
  using assms
  apply (subst complex-eq-if-Re-eq)
  using Re-complex-of-real[of (cmod z1)²]
  by auto (metis is-real-complex-of-real of-real-power, metis div-reals)
  also have ... ↔ cmod z1 = sqrt (Re (-k2 / k1))
  by (metis norm-ge-zero real-sqrt-ge-0-iff real-sqrt-pow2 real-sqrt-power)
  also have ... ↔ (exists φ. z1 = rcis (sqrt (Re (-k2 / k1))) φ)
  using rcis-cmod-arg[of z1, symmetric] assms abs-of-nonneg[of sqrt (Re (-k2 / k1))]
  using ⟨Re (-k2/k1) ≥ 0⟩
  by auto
  finally show ?thesis
  .
qed

```



```

lemma circline-type-neg-card-gt3-diag:
  assumes circline-type H < 0 circline-diag H
  shows ∃ A B C. A ≠ B ∧ A ≠ C ∧ B ≠ C ∧ {A, B, C} ⊆ circline-set H
using assms
unfolding circline-set-def
apply (simp del: HOL.ex-simps)
proof (transfer)
  fix H
  obtain A B C D where HH: Rep-circline-mat H = (A, B, C, D)
  by (cases Rep-circline-mat H) auto
  hence HH-elems: is-real A is-real D C = cnj B
  using hermitean-elems[of A B C D] Rep-circline-mat[of H]
  by auto
  assume circline-diag-rep H circline-type-rep H < 0
  hence B = 0 C = 0 Re A * Re D < 0 A ≠ 0
  using HH ⟨is-real A⟩ ⟨is-real D⟩
  unfolding circline-diag-rep-def circline-type-rep-def
  by auto

```

```

let ?x = sqrt (Re (- D / A))
let ?A = (rcis ?x 0, 1)
let ?B = (rcis ?x (pi/2), 1)
let ?C = (rcis ?x pi, 1)
from quad-form-diagonal-iff[OF ‹A ≠ 0› ‹is-real A› ‹is-real D› ‹Re A * Re D < 0›]
have quad-form ?A (A, 0, 0, D) = 0 quad-form ?B (A, 0, 0, D) = 0 quad-form
?C (A, 0, 0, D) = 0
by (auto simp del: rcis-zero-arg)
moreover
have Re (D / A) < 0
using ‹Re A * Re D < 0› ‹A ≠ 0› ‹is-real A› ‹is-real D›
by (subst Re-divide-real) (auto, metis divide-less-0-iff mult-eq-0-iff mult-neg-neg
mult-pos-pos not-less-iff-gr-or-eq)
hence ¬ Abs-homo-coords ?A ≈ Abs-homo-coords ?B ∧ ¬ Abs-homo-coords ?A
≈ Abs-homo-coords ?C ∧ ¬ Abs-homo-coords ?B ≈ Abs-homo-coords ?C
unfolding rcis-def
by (auto simp add: Abs-homo-coords-inverse cis-def)
ultimately
show ∃ A B C. ¬ A ≈ B ∧ ¬ A ≈ C ∧ ¬ B ≈ C ∧ (on-circline-rep H A ∧
on-circline-rep H B ∧ on-circline-rep H C)
using HH ‹B = 0› ‹C = 0›
by (rule-tac x=Abs-homo-coords ?A in exI, rule-tac x=Abs-homo-coords ?B in
exI, rule-tac x=Abs-homo-coords ?C in exI)
(simp add: on-circline-rep-def Abs-homo-coords-inverse Let-def)
qed

lemma circline-type-neg-card-gt3:
assumes circline-type H < 0
shows ∃ A B C. A ≠ B ∧ A ≠ C ∧ B ≠ C ∧ {A, B, C} ⊆ circline-set H
proof-
obtain M H' where moebius-circline M H = H' circline-diag H'
using circline-diagonalize[of H] assms
by auto
moreover
hence circline-type H' < 0
using assms moebius-preserve-circline-type
by auto
ultimately
obtain A B C where A ≠ B A ≠ C B ≠ C {A, B, C} ⊆ circline-set H'
using circline-type-neg-card-gt3-diag[of H']
by auto
let ?iM = moebius-inv M
have moebius-circline ?iM H' = H
using ‹moebius-circline M H = H'›[symmetric]
by simp
let ?A = moebius-pt ?iM A and ?B = moebius-pt ?iM B and ?C = moebius-pt
?iM C
have ?A ∈ circline-set H ?B ∈ circline-set H ?C ∈ circline-set H

```

```

using <moebius-circline ?iM H' = H>[symmetric] <\{A, B, C\} ⊆ circline-set H'
  by (simp-all add: moebius-circline-set[symmetric])
moreover
have ?A ≠ ?B ?A ≠ ?C ?B ≠ ?C
  using bij-moebius-pt[of moebius-inv M] <A ≠ B> <A ≠ C> <B ≠ C>
  unfolding bij-def inj-on-def
  by blast+
ultimately
show ?thesis
  by auto
qed

```

11.7.4 Positive type circline set cardinality

```

lemma circline-type-pos-card-eq0-diag:
assumes circline-diag H circline-type H > 0
shows circline-set H = {}
using assms
unfolding circline-set-def
apply simp
proof transfer
fix H
obtain A B C D where HH: Rep-circline-mat H = (A, B, C, D)
  by (cases Rep-circline-mat H) auto
hence HH-elems: is-real A is-real D C = cnj B
  using hermitean-elems[of A B C D] Rep-circline-mat[of H]
  by auto
assume circline-diag-rep H 0 < circline-type-rep H
hence B = 0 C = 0 Re A * Re D > 0 A ≠ 0
  using <circline-diag-rep H> HH <is-real A> <is-real D>
  unfolding circline-diag-rep-def circline-type-rep-def
  by auto
show ∀ x. ¬ on-circline-rep H x
proof
fix x
obtain x1 x2 where xx:Rep-homo-coords x = (x1, x2)
  by (rule obtain-homo-coords)
have (Re A > 0 ∧ Re D > 0) ∨ (Re A < 0 ∧ Re D < 0)
  using <Re A * Re D > 0
  by (metis linorder-neqE-linordered-idom mult-eq-0-iff zero-less-mult-pos zero-less-mult-pos2)
moreover
have (Re (x1 * cnj x1) ≥ 0 ∧ Re (x2 * cnj x2) > 0) ∨ (Re (x1 * cnj x1) >
0 ∧ Re (x2 * cnj x2) ≥ 0)
  using Rep-homo-coords[of x] xx
  by auto (metis complex-surj complex-zero-def sum-squares-gt-zero-iff) +
ultimately
have Re A * Re (x1 * cnj x1) + Re D * Re (x2 * cnj x2) ≠ 0
  apply auto
apply (metis add-less-cancel-left add-pos-pos mult-eq-0-iff mult-pos-pos sum-squares-eq-zero-iff)

```

```

sum-squares-gt-zero-iff)
  apply (metis (lifting, no-types) add-pos-pos comm-semiring-1-class.normalize-semiring-rules(6)
mult-eq-0-iff mult-pos-pos sum-squares-gt-zero-iff)
    apply (metis add-less-cancel-left add-neg-neg mult-eq-0-iff mult-pos-neg2
sum-squares-eq-zero-iff sum-squares-gt-zero-iff)
      apply (metis (lifting, no-types) add-neg-neg comm-semiring-1-class.normalize-semiring-rules(6)
mult-eq-0-iff mult-pos-neg2 sum-squares-gt-zero-iff)
        done
      hence A * (x1 * cnj x1) + D * (x2 * cnj x2) ≠ 0
        using ‹is-real A› ‹is-real D›
        by (metis Re-mult-real complex-Re-add complex-Re-zero)
      thus ¬ on-circline-rep H x
        using HH ‹B = 0› ‹C = 0› xx
        unfolding on-circline-rep-def Let-def
        by (simp add: vec-cnj-def field-simps)
      qed
qed

lemma circline-type-pos-card-eq0:
  assumes circline-type H > 0
  shows circline-set H = {}
proof-
  obtain M H' where moebius-circline M H = H' circline-diag H'
    using circline-diagonalize[of H] assms
    by auto
  moreover
  hence circline-type H' > 0
    using assms moebius-preserve-circline-type
    by auto
  ultimately
  have circline-set H' = {}
    using circline-type-pos-card-eq0-diag[of H']
    by auto
  let ?iM = moebius-inv M
  have moebius-circline ?iM H' = H
    using ‹moebius-circline M H = H'›[symmetric]
    by simp
  thus ?thesis
    using ‹circline-set H' = {}›
    by (auto simp add: moebius-circline-set[symmetric])
qed

```

11.7.5 Cardinality determines type

```

lemma card-eq1-circline-type-zero:
  assumes ∃ z. circline-set H = {z}
  shows circline-type H = 0
proof (cases circline-type H < 0)
  case True

```

```

thus ?thesis
  using circline-type-neg-card-gt3[of H] assms
  by auto
next
  case False
  show ?thesis
  proof (cases circline-type H > 0)
    case True
    thus ?thesis
      using circline-type-pos-card-eq0[of H] assms
      by auto
  next
    case False
    thus ?thesis
      using linorder.cases[of "circline-type H"] lessThan0
      by simp
  qed
qed

```

11.7.6 Circline set is injective

```

lemma inj-circline-set:
  assumes circline-set H = circline-set H' circline-set H ≠ {}
  shows H = H'
proof (cases circline-type H < 0)
  case True
  then obtain A B C where A ≠ B A ≠ C B ≠ C {A, B, C} ⊆ circline-set H
    using circline-type-neg-card-gt3[of H]
    by auto
  hence ∃!H. A ∈ circline-set H ∧ B ∈ circline-set H ∧ C ∈ circline-set H
    using unique-circline-set[of A B C]
    by simp
  thus ?thesis
    using linorder.cases[of "circline-set H = circline-set H'"]
    by auto
next
  case False
  show ?thesis
  proof (cases circline-type H = 0)
    case True
    moreover
    then obtain A where {A} = circline-set H
      using circline-type-zero-card-eq1[of H]
      by auto
    moreover
    hence circline-type H' = 0
      using linorder.cases[of "circline-set H = circline-set H'"]
      by auto
    ultimately

```

```

show ?thesis
  using unique-circline-type-zero[of A] ⟨circline-set H = circline-set H'⟩
  by auto
next
  case False
  hence circline-type H > 0
    using ⟨¬ (circline-type H < 0)⟩
    by auto
  thus ?thesis
    using ⟨circline-set H ≠ {}⟩ circline-type-pos-card-eq0[of H]
    by auto
qed
qed

```

11.8 Symmetric points wrt. circline

definition circline-symmetric-rep **where**

```

circline-symmetric-rep z1 z2 H ⟷
  (let z1 = Rep-homo-coords z1;
   z2 = Rep-homo-coords z2;
   H = Rep-circline-mat H in
   bilinear-form z1 z2 H = 0)

```

lift-definition circline-symmetric :: complex-homo ⇒ complex-homo ⇒ circline
 \Rightarrow bool **is** circline-symmetric-rep
by (auto simp add: circline-symmetric-rep-def bilinear-form-scale-m bilinear-form-scale-v1
bilinear-form-scale-v2 simp del: vec-cnj-sv quad-form-def bilinear-form-def)

lemma symmetry-principle:

```

assumes circline-symmetric z1 z2 H
shows circline-symmetric (moebius-pt M z1) (moebius-pt M z2) (moebius-circline
M H)
using assms
proof (transfer)
  fix z1 z2 H M
  assume circline-symmetric-rep z1 z2 H
  thus circline-symmetric-rep (moebius-pt-rep M z1) (moebius-pt-rep M z2) (moebius-circline-rep
M H)
    apply (auto simp add: circline-symmetric-rep-def simp del: bilinear-form-def)
    using Rep-moebius-mat[of M]
    by (subst bilinear-form-congruence[symmetric]) simp-all
qed

```

Symmetry wrt. unit-circle

lemma circline-symmetric-0inf-disc: circline-symmetric 0_h ∞_h unit-circle
by (transfer) (simp add: circline-symmetric-rep-def vec-cnj-def)

lemma circline-symmetric-inv-homo-disc: circline-symmetric a (inversion-homo
a) unit-circle

```

unfolding inversion-homo-def
by (transfer) (case-tac Rep-homo-coords a, auto simp add: circline-symmetric-rep-def
vec-cnj-def split-def Let-def)

lemma circline-symmetric-inv-homo-disc':
assumes circline-symmetric a a' unit-circle
shows a' = inversion-homo a
unfolding inversion-homo-def
using assms
proof (transfer)
fix a a'
obtain a1 a2 where aa: Rep-homo-coords a = (a1, a2)
by (rule obtain-homo-coords)
obtain a1' a2' where aa': Rep-homo-coords a' = (a1', a2')
by (rule obtain-homo-coords)
assume *: circline-symmetric-rep a a' unit-circle-rep
show a' ≈ (cnj-homo-coords ∘ reciprocal-homo-coords) a
proof (cases a1' = 0)
case True
thus ?thesis
using aa aa' * Rep-homo-coords[of a'] Rep-homo-coords[of a]
by (auto simp add: circline-symmetric-rep-def vec-cnj-def field-simps)
next
case False
show ?thesis
proof (cases a2 = 0)
case True
thus ?thesis
using (a1' ≠ 0)
using aa aa' * Rep-homo-coords[of a]
by (simp add: circline-symmetric-rep-def vec-cnj-def field-simps)
next
case False
thus ?thesis
using (a1' ≠ 0) aa aa' *
by (simp add: circline-symmetric-rep-def vec-cnj-def field-simps) (rule-tac
x=cnj a2 / a1' in exI, simp add: field-simps)
qed
qed
qed

```

11.9 Oriented circlines; discs

```

definition ocircline-mat-eq where
[simp]: ocircline-mat-eq A B ↔ (exists k::real. k > 0 ∧ Rep-circline-mat B =
complex-of-real k *sm (Rep-circline-mat A))

lemma [simp]: ocircline-mat-eq H H
by (simp, rule-tac x=1 in exI, simp)

```

```

quotient-type ocircline = circline-mat / ocircline-mat-eq
proof (rule equivpI)
  show reflp ocircline-mat-eq
    unfolding reflp-def
    by (auto, rule-tac x=1 in exI, simp)
next
  show symp ocircline-mat-eq
    unfolding symp-def
    by (auto, rule-tac x=1/k in exI, simp)
next
  show transp ocircline-mat-eq
    unfolding transp-def
    by (auto, rule-tac x=ka*k in exI, simp add: mult-pos-pos)
qed

lift-definition on-ocircline :: ocircline  $\Rightarrow$  complex-homo  $\Rightarrow$  bool is on-circline-rep
by (auto simp add: on-circline-rep-def quad-form-scale-m quad-form-scale-v Let-def
simp del: vec-cnj-sv quad-form-def)

definition ocircline-set :: ocircline  $\Rightarrow$  complex-homo set where
  ocircline-set H = {z. on-ocircline H z}

disc and disc complement

definition in-ocircline-rep where
  in-ocircline-rep H z  $\longleftrightarrow$ 
    (let z = Rep-homo-coords z;
H = Rep-circline-mat H
in Re (quad-form z H) < 0)

lift-definition in-ocircline :: ocircline  $\Rightarrow$  complex-homo  $\Rightarrow$  bool is in-ocircline-rep
proof-
  fix H H' z z'
  assume ocircline-mat-eq H H' z ≈ z'
  then obtain k k' where
    *: 0 < k Rep-circline-mat H' = cor k *sm Rep-circline-mat H k' ≠ 0 Rep-homo-coords
z' = k' *sv Rep-homo-coords z
    by auto
    hence quad-form (Rep-homo-coords z') (Rep-circline-mat H') = cor k * cor
((cmod k')^2) * quad-form (Rep-homo-coords z) (Rep-circline-mat H)
    by (simp add: quad-form-scale-v quad-form-scale-m del: vec-cnj-sv quad-form-def)
    hence Re (quad-form (Rep-homo-coords z') (Rep-circline-mat H')) =
k * (cmod k')^2 * Re (quad-form (Rep-homo-coords z) (Rep-circline-mat H))
    using Rep-circline-mat[of H] quad-form-hermitean-real[of Rep-circline-mat H]
    by (simp add: complex-of-real-Re power2-eq-square)
  thus in-ocircline-rep H z = in-ocircline-rep H' z'
  unfolding in-ocircline-rep-def Let-def
  using (k > 0) (k' ≠ 0)
  apply auto

```

```

apply (metis mult-pos-neg mult-pos-pos norm-eq-zero zero-less-power2)
apply (metis comm-semiring-1-class.normalize-semiring-rules(10) mult-less-cancel-left-pos
zero-less-norm-iff zero-less-power)
done
qed

definition disc where
disc H = {z. in-ocircline H z}

definition out-ocircline-rep where
out-ocircline-rep H z  $\longleftrightarrow$ 
(let z = Rep-homo-coords z;
H = Rep-circline-mat H
in Re (quad-form z H) > 0)

lift-definition out-ocircline :: ocircline  $\Rightarrow$  complex-homo  $\Rightarrow$  bool is out-ocircline-rep
proof-
fix H H' z z'
assume ocircline-mat-eq H H' z  $\approx$  z'
then obtain k k' where
*: 0 < k Rep-circline-mat H' = cor k *sm Rep-circline-mat H k'  $\neq$  0 Rep-homo-coords
z' = k' *sv Rep-homo-coords z
by auto
hence quad-form (Rep-homo-coords z') (Rep-circline-mat H') = cor k * cor
((cmod k')2) * quad-form (Rep-homo-coords z) (Rep-circline-mat H)
by (simp add: quad-form-scale-v quad-form-scale-m del: vec-cnj-sv quad-form-def)
hence Re (quad-form (Rep-homo-coords z') (Rep-circline-mat H')) =
k * (cmod k')2 * Re (quad-form (Rep-homo-coords z) (Rep-circline-mat H))
using Rep-circline-mat[of H] quad-form-hermitean-real[of Rep-circline-mat H]
by (simp add: complex-of-real-Re power2-eq-square)
thus out-ocircline-rep H z = out-ocircline-rep H' z'
unfolding out-ocircline-rep-def Let-def
using {k > 0} {k'  $\neq$  0}
apply auto
apply (metis mult-pos-pos norm-eq-zero zero-less-power2)
apply (metis comm-semiring-1-class.normalize-semiring-rules(10) mult-less-cancel-left-pos
zero-less-norm-iff zero-less-power)
done
qed

definition disc-compl where
disc-compl H = {z. out-ocircline H z}

lemma in-on-out: in-ocircline H z  $\vee$  on-ocircline H z  $\vee$  out-ocircline H z
proof transfer
fix z H
show in-ocircline-rep H z  $\vee$  on-circline-rep H z  $\vee$  out-ocircline-rep H z
using Rep-circline-mat[of H] quad-form-hermitean-real[of Rep-circline-mat H]
Rep-homo-coords z]

```

by (*simp add: in-ocircline-rep-def on-circline-rep-def out-ocircline-rep-def Let-def*)
(metis complex-Im-zero complex-Re-zero complex-equality linorder-cases)
qed

lemma *disc H ∪ disc-compl H ∪ ocircline-set H = UNIV*
unfolding *disc-def disc-compl-def ocircline-set-def*
using *in-on-out[of H]*
by *auto*

lemma
disc-inter-disc-compl: disc H ∩ disc-compl H = {}
unfolding *disc-def disc-compl-def*
by *auto (transfer, auto simp add: in-ocircline-rep-def out-ocircline-rep-def Let-def)*

lemma
disc-inter-ocircline-set: disc H ∩ ocircline-set H = {}
unfolding *disc-def ocircline-set-def*
by *auto (transfer, simp add: in-ocircline-rep-def on-circline-rep-def Let-def)*

lemma
disc-compl-inter-ocircline-set: disc-compl H ∩ ocircline-set H = {}
unfolding *disc-compl-def ocircline-set-def*
by *auto (transfer, simp add: out-ocircline-rep-def on-circline-rep-def Let-def)*

Opposite orientation

definition *opposite-ocircline-rep where*
opposite-ocircline-rep H =
(let H = Rep-circline-mat H in
*Abs-circline-mat (-1 *_{sm} H))*

lemma *circline-mat-mult-m1 [simp]: Rep-circline-mat (Abs-circline-mat (-1 *_{sm} Rep-circline-mat H)) = (-1 *_{sm} Rep-circline-mat H)*
proof-

have $-1 = \text{cor}(-1)$
by (*simp add: complex-of-real-def*)
thus *?thesis*
using *circline-mat-mult-sm-Rep[of -1 H]*
by *auto*

qed

lemma [*simp*]: *Rep-circline-mat (opposite-ocircline-rep H) = (-1 *_{sm} Rep-circline-mat H)*
unfolding *opposite-ocircline-rep-def*
by *auto*

lift-definition *opposite-ocircline :: ocircline ⇒ ocircline* **is** *opposite-ocircline-rep*
by *auto*

lemma *opposite-ocircline-rep-opposite-ocircline-rep*

```

[simp]: opposite-ocircline-rep (opposite-ocircline-rep H) = H
by (simp add: opposite-ocircline-rep-def Rep-circline-mat-inverse)

lemma opposite-ocircline-opposite-ocircline
[simp]: opposite-ocircline (opposite-ocircline H) = H
by (transfer) (auto, rule-tac x=1 in exI, simp)

lemma ocircline-set-opposite-ocircline
[simp]: ocircline-set (opposite-ocircline H) = ocircline-set H
unfolding ocircline-set-def
by auto (transfer, auto simp add: on-circline-rep-def quad-form-scale-m simp del:
quad-form-def)+

lemma disc-compl-opposite: disc-compl (opposite-ocircline H) = disc H
unfolding disc-def disc-compl-def
apply auto
apply (transfer)
apply (auto simp add: in-ocircline-rep-def out-ocircline-rep-def quad-form-scale-m
simp del: quad-form-def)
apply (transfer)
apply (auto simp add: in-ocircline-rep-def out-ocircline-rep-def quad-form-scale-m
simp del: quad-form-def)
done

lemma disc-opposite:
disc (opposite-ocircline H) = disc-compl H
using disc-compl-opposite[of opposite-ocircline H]
by simp

of-ocircline, pos-oriented, of-circline

lift-definition of-ocircline :: ocircline ⇒ circline is id::circline-mat ⇒ circline-mat
by auto (rule-tac x=k in exI, simp)

lemma of-ocircline-opposite-ocircline [simp]:
of-ocircline (opposite-ocircline H) = of-ocircline H
by (transfer) (auto, rule-tac x=-1 in exI, simp)

lemma circline-set-ocircline-set [simp]:
circline-set (of-ocircline H) = ocircline-set H
unfolding ocircline-set-def circline-set-def
by (safe) (transfer, simp)+

lemma inj-of-ocircline:
assumes of-ocircline H = of-ocircline H'
shows H = H' ∨ H = opposite-ocircline H'
using assms
by (transfer) (auto, metis linorder-neqE-linordered-idom neg-0-less-iff-less of-real-minus)

```

```

lemma inj-ocircline-set:
  assumes ocircline-set  $H = \text{ocircline-set } H'$ 
  shows  $H = H' \vee H = \text{opposite-ocircline } H'$ 
proof-
  from assms have circline-set (of-ocircline  $H) = \text{circline-set (of-ocircline } H')$ 
  circline-set (of-ocircline  $H') \neq \{\}$ 
  using circline-set-ocircline-set[symmetric, of  $H)]$  circline-set-ocircline-set[symmetric,
  of  $H']$ 
  by blast+
  hence of-ocircline  $H = \text{of-ocircline } H'$ 
  by (simp add: inj-circline-set)
  thus ?thesis
  by (rule inj-of-ocircline)
qed

```

```

definition pos-oriented-rep where
  pos-oriented-rep  $H \longleftrightarrow$ 
    (let  $(A, B, C, D) = \text{Rep-circline-mat } H$ 
     in  $(\text{Re } A > 0 \vee (\text{Re } A = 0 \wedge ((B \neq 0 \wedge \arg B > 0) \vee (B = 0 \wedge \text{Re } D > 0))))$ )
lemma pos-oriented-rep: pos-oriented-rep  $H \vee \text{pos-oriented-rep } (\text{opposite-ocircline-rep } H)$ 
proof-
  obtain  $A B C D$  where  $\text{HH: Rep-circline-mat } H = (A, B, C, D)$ 
  by (cases Rep-circline-mat  $H$ ) auto
  moreover
  hence  $\text{Re } A = 0 \wedge \text{Re } D = 0 \longrightarrow B \neq 0$ 
  using Rep-circline-mat[of  $H)]$  hermitean-elems[of  $A B C D]$ 
  by (cases  $A$ , cases  $D$ ) auto
  moreover
  have  $B \neq 0 \wedge \neg 0 < \arg B \longrightarrow 0 < \arg (-B)$ 
  using MoreComplex.canon-ang-plus-pi2[of  $\arg B)]$  arg-bounded[of  $B]$ 
  by (auto simp add: arg-uminus)
  ultimately
  show ?thesis
  by (auto simp add: pos-oriented-rep-def)
qed

```

```

lift-definition pos-oriented :: ocircline  $\Rightarrow$  bool is pos-oriented-rep
apply (auto simp add: pos-oriented-rep-def mult-pos-pos)
apply (metis arg-mult-real-positive)
apply (metis arg-mult-real-positive)
apply (metis zero-less-mult-pos)+
apply (metis arg-mult-real-positive)
apply (metis arg-mult-real-positive)

```

```

apply (metis zero-less-mult-pos)+  

done

lemma pos-oriented: pos-oriented  $H \vee$  pos-oriented (opposite-ocircline  $H$ )  

by (transfer) (rule pos-oriented-rep)

lemma pos-oriented-opposite-ocircline:  

  pos-oriented (opposite-ocircline  $H$ )  $\longleftrightarrow \neg$  pos-oriented  $H$   

proof transfer  

  fix  $H$   

  obtain  $A B C D$  where  $HH: Rep\text{-circline-mat } H = (A, B, C, D)$   

    by (cases Rep-circline-mat  $H$ ) auto  

  moreover  

  hence  $Re A = 0 \wedge Re D = 0 \longrightarrow B \neq 0$   

    using Rep-circline-mat[of  $H$ ] hermitean-elems[of  $A B C D$ ]  

    by (cases  $A$ , cases  $D$ ) auto  

  moreover  

  have  $B \neq 0 \wedge \neg 0 < arg B \longrightarrow 0 < arg (-B)$   

    using MoreComplex.canon-ang-plus-pi2[of arg  $B$ ] arg-bounded[of  $B$ ]  

    by (auto simp add: arg-uminus)  

  moreover  

  have  $B \neq 0 \wedge 0 < arg B \longrightarrow \neg 0 < arg (-B)$   

    using MoreComplex.canon-ang-plus-pi1[of arg  $B$ ] arg-bounded[of  $B$ ]  

    by (auto simp add: arg-uminus)  

  ultimately  

  show pos-oriented-rep (opposite-ocircline-rep  $H$ ) = ( $\neg$  pos-oriented-rep  $H$ )  

    unfolding pos-oriented-rep-def  

    by simp (metis not-less-iff-gr-or-eq)
  qed

lemma pos-oriented-circle-inf:  

  assumes  $\infty_h \notin$  circline-set  $H$   

  shows pos-oriented  $H \longleftrightarrow \infty_h \notin disc H$   

using assms  

unfolding circline-set-def disc-def  

apply simp  

proof transfer  

  fix  $H$   

  obtain  $A B C D$  where  $HH: Rep\text{-circline-mat } H = (A, B, C, D)$   

    by (cases Rep-circline-mat  $H$ ) auto  

  hence is-real  $A$   

    using Rep-circline-mat[of  $H$ ] hermitean-elems  

    by auto  

  assume  $\neg$  on-circline-rep  $H$  inf-homo-rep  

  thus pos-oriented-rep  $H$  = ( $\neg$  in-ocircline-rep  $H$  inf-homo-rep)  

    using HH (is-real  $A$ )  

    by (cases  $A$ ) (auto simp add: on-circline-rep-def in-ocircline-rep-def Let-def  

      pos-oriented-rep-def vec-cnj-def)
  qed

```

```

lemma
  assumes is-circle (of-ocircline H) (a, r) = euclidean-circle (of-ocircline H)
  circline-type (of-ocircline H) < 0
  shows pos-oriented H  $\longleftrightarrow$  of-complex a ∈ disc H
  using assms
  unfolding disc-def
  apply simp
  proof transfer
    fix H a r
    obtain A B C D where HH: Rep-circline-mat H = (A, B, C, D)
      by (cases Rep-circline-mat H) auto
    hence is-real A is-real D C = cnj B
      using Rep-circline-mat[of H] hermitean-elems
      by auto

    assume  $*: \neg \text{circline-A0-rep}(\text{id } H) (a, r) = \text{euclidean-circle-rep}(\text{id } H) \text{ circline-type-rep}$ 
     $(\text{id } H) < 0$ 
    hence  $A \neq 0 \text{ Re } A \neq 0$ 
      using HH (is-real A)
      by (case-tac[] A) (auto simp add: circline-A0-rep-def)
    have  $\text{Re}(A*D - B*C) < 0$ 
      using <circline-type-rep (id H) < 0> HH
      by (simp add: circline-type-rep-def)
    have  $(A * (D * \text{cnj } A) - B * (C * \text{cnj } A)) / (A * \text{cnj } A) = (A*D - B*C) / A$ 
      using <A ≠ 0>
      by (simp add: field-simps)
    hence  $0 < \text{Re } A \longleftrightarrow \text{Re}((A * (D * \text{cnj } A) - B * (C * \text{cnj } A)) / (A * \text{cnj } A)) < 0$ 
      using (is-real A) (A ≠ 0) (Re (A*D - B*C) < 0)
      by (auto simp add: Re-divide-real, metis divide-less-0-iff less-iff-diff-less-0, metis
      divide-less-0-iff less-iff-diff-less-0 mult-neg-neg zero-less-mult-pos)
      thus pos-oriented-rep H = in-ocircline-rep H (of-complex-coords a)
      using HH (Re A ≠ 0) * (is-real A)
      by (simp add: circline-A0-rep-def euclidean-circle-rep-def pos-oriented-rep-def
      in-ocircline-rep-def Let-def vec-cnj-def complex-cnj field-simps)
    qed

definition of-circline-rep :: circline-mat ⇒ circline-mat where
  of-circline-rep H = (if pos-oriented-rep H then H else opposite-ocircline-rep H)

lift-definition of-circline :: circline ⇒ ocircline is of-circline-rep
proof –
  fix H H'
  assume circline-mat-eq H H'
  then obtain k where  $*: k \neq 0 \text{ Rep-circline-mat } H' = \text{cor } k *_{sm} \text{Rep-circline-mat } H$ 

```

```

by auto
obtain A B C D where HH: Rep-circline-mat H = (A, B, C, D)
  by (cases Rep-circline-mat H) auto
obtain A' B' C' D' where HH': Rep-circline-mat H' = (A', B', C', D')
  by (cases Rep-circline-mat H') auto

show ocircline-mat-eq (of-circline-rep H) (of-circline-rep H')
proof (cases Re A > 0)
  case True
    show ?thesis
    proof (cases k > 0)
      case True
        hence Re A' > 0
        using ⟨Re A > 0⟩ * HH HH'
        by (auto simp add: mult-pos-pos)
      thus ?thesis
        using ⟨Re A > 0⟩
        using * ⟨k > 0⟩ HH HH'
        by (auto simp add: pos-oriented-rep-def of-circline-rep-def Let-def split-def)
    next
      case False
      hence k < 0
      using ⟨k ≠ 0⟩
      by auto
      hence Re A' < 0
      using ⟨Re A > 0⟩ * HH HH'
      by (auto simp add: mult-neg-pos)
      thus ?thesis
        using ⟨Re A > 0⟩
        using * ⟨k < 0⟩ HH HH'
        using circline-mat-mult-sm-Rep[of -k H]
        by (auto simp add: pos-oriented-rep-def of-circline-rep-def Let-def split-def)
    (rule-tac x=-k in exI, simp)
  qed
next
case False
show ?thesis
proof (cases Re A < 0)
  case True
  show ?thesis
  proof (cases k > 0)
    case True
    hence Re A' < 0
    using ⟨Re A < 0⟩
    using * HH HH'
    by (auto simp add: mult-pos-neg)
  moreover
  have -1 = cor (-1)
    by (simp add: complex-of-real-def)

```

```

ultimately
show ?thesis
  using ⟨Re A < 0⟩
  using * ⟨k > 0⟩ HH HH'
  using circline-mat-mult-sm-Rep[of -k H]
  using circline-mat-mult-sm-Rep[of -1 H]
  by (auto simp add: pos-oriented-rep-def of-circline-rep-def Let-def split-def)
next
  case False
  hence k < 0
    using ⟨k ≠ 0⟩
    by simp
  hence Re A' > 0
    using ⟨Re A < 0⟩
    using * HH HH'
    by (auto simp add: mult-neg-neg)
moreover
have -1 = cor (-1)
  by (simp add: complex-of-real-def)
ultimately
show ?thesis
  using ⟨Re A < 0⟩
  using * ⟨k < 0⟩ HH HH'
  using circline-mat-mult-sm-Rep[of -k H]
  using circline-mat-mult-sm-Rep[of -1 H]
  by (auto simp add: pos-oriented-rep-def of-circline-rep-def Let-def split-def)
(rule-tac x=-k in exI, simp)
qed
next
  case False
  hence Re A = 0
    using ⟨¬ Re A > 0⟩
    by auto
  hence Re A' = 0
    using * HH HH'
    by auto

  show ?thesis
  proof (cases B ≠ 0)
    case True
    show ?thesis
    proof (cases arg B > 0)
      case True
      show ?thesis
      proof (cases arg B' > 0)
        case True
        hence k > 0
          using ⟨arg B > 0⟩
          using * HH HH' arg-mult[of cor k B] ⟨B ≠ 0⟩ ⟨k ≠ 0⟩
      qed
    qed
  qed

```

```

using arg-complex-of-real-negative[of k] arg-complex-of-real-positive[of
k]
using MoreComplex.canon-ang-plus-pi1[of arg B] arg-bounded[of B]
by (cases k > 0) (auto simp add: arg-mult field-simps)
thus ?thesis
  using ⟨arg B > 0⟩ ⟨arg B' > 0⟩ ⟨B ≠ 0⟩ ⟨Re A = 0⟩ ⟨Re A' = 0⟩ HH
HH' *
  by (auto simp add: pos-oriented-rep-def of-circline-rep-def)
next
  case False
  hence k < 0
    using ⟨arg B > 0⟩
    using * HH HH' arg-mult[of cor k B] ⟨B ≠ 0⟩ ⟨k ≠ 0⟩
    using arg-complex-of-real-negative[of k] arg-complex-of-real-positive[of
k]
    by (cases k > 0) (auto simp add: arg-mult field-simps canon-ang-arg)
    thus ?thesis
      using ⟨arg B > 0⟩ ⟨¬ arg B' > 0⟩ ⟨Re A = 0⟩ ⟨Re A' = 0⟩ ⟨B ≠ 0⟩ HH
HH' *
      using circline-mat-mult-sm-Rep[of -k H]
      by (auto simp add: pos-oriented-rep-def of-circline-rep-def) (rule-tac
x=-k in exI, simp)+
    qed
  next
  case True
  show ?thesis
  proof (cases arg B' > 0)
    case True
    hence k < 0
      using ⟨¬ arg B > 0⟩
      using * HH HH' arg-mult[of cor k B] ⟨B ≠ 0⟩ ⟨k ≠ 0⟩
      using arg-complex-of-real-negative[of k] arg-complex-of-real-positive[of
k]
      by (cases k > 0) (auto simp add: arg-mult field-simps canon-ang-arg)
      thus ?thesis
        using ⟨¬ arg B > 0⟩ ⟨arg B' > 0⟩ ⟨B ≠ 0⟩ ⟨Re A = 0⟩ ⟨Re A' = 0⟩ HH
HH' *
        by (auto simp add: pos-oriented-rep-def of-circline-rep-def) (rule-tac
x=-k in exI, simp)+
    next
    case False
    hence k > 0
      using ⟨¬ arg B > 0⟩
      using * HH HH' arg-mult[of cor k B] ⟨B ≠ 0⟩ ⟨k ≠ 0⟩
      using arg-complex-of-real-negative[of k] arg-complex-of-real-positive[of
k]
      using MoreComplex.canon-ang-plus-pi2[of arg B] arg-bounded[of B]
      by (cases k > 0) (auto simp add: arg-mult field-simps canon-ang-arg)
      thus ?thesis

```

```

using  $\neg \arg B > 0 \wedge \neg \arg B' > 0 \wedge \langle \operatorname{Re} A = 0 \rangle \wedge \langle \operatorname{Re} A' = 0 \rangle \wedge \langle B \neq 0 \rangle$ 
HH HH' *
  using circline-mat-mult-sm-Rep[of -k H]
  by (auto simp add: pos-oriented-rep-def of-circline-rep-def)
qed
qed
next
case False
hence  $B' = 0$ 
  using * HH HH'
  by simp
have  $\operatorname{Re} D \neq 0$ 
  using  $\langle \operatorname{Re} A = 0 \rangle \wedge \neg B \neq 0$ 
  using Rep-circline-mat[of H] HH hermitean-elems[of A B C D]
  by (cases A, cases D) auto
show ?thesis
  using  $\neg B \neq 0 \wedge B' = 0 \wedge \langle \operatorname{Re} A = 0 \rangle \wedge \langle \operatorname{Re} A' = 0 \rangle \wedge \langle \operatorname{Re} D \neq 0 \rangle$  HH HH' *
  apply (auto simp add: of-circline-rep-def pos-oriented-rep-def)
  apply (metis zero-less-mult-pos2)
  apply (rule-tac x=-k in exI, simp, metis linorder-cases mult-pos-pos)
  apply (rule-tac x=-k in exI, simp, metis linorder-cases zero-less-mult-pos)
  apply (rule-tac x=k in exI, simp, metis mult-neg-neg neqE)
done
qed
qed
qed
qed
qed

lemma pos-oriented-of-circline: pos-oriented (of-circline H)
proof (transfer)
fix H
show pos-oriented-rep (of-circline-rep H)
  using pos-oriented-rep[of H]
  unfolding of-circline-rep-def
  by auto
qed

lemma of-ocircline-of-circline [simp]: of-ocircline (of-circline H) = H
apply (transfer)
apply (auto simp add: of-circline-rep-def)
by (rule-tac x=1 in exI, simp) (rule-tac x=-1 in exI, auto simp add: pos-oriented-rep-def
complex-of-real-def)

lemma of-circline-of-ocircline-pos-oriented [simp]:
assumes pos-oriented H
shows of-circline (of-ocircline H) = H
using assms
by (transfer) (simp add: of-circline-rep-def, rule-tac x=1 in exI, simp)

```

```

lemma ocircline-set-circline-set[simp]: ocircline-set (of-circline H) = circline-set H
  unfolding ocircline-set-def circline-set-def
  proof (safe)
    fix z
    assume on-ocircline (of-circline H) z
    thus on-circline H z
      by (transfer) (auto simp add: on-circline-rep-def of-circline-rep-def Let-def
quad-form-scale-m simp del: quad-form-def split: split-if-asm)
  next
    fix z
    assume on-circline H z
    thus on-ocircline (of-circline H) z
      by (transfer) (auto simp add: on-circline-rep-def of-circline-rep-def Let-def
quad-form-scale-m simp del: quad-form-def)
  qed

lemma inj-of-circline:
  assumes of-circline H = of-circline H'
  shows H = H'
  using assms
  proof (transfer)
    fix H H'
    assume ocircline-mat-eq (of-circline-rep H) (of-circline-rep H')
    then obtain k where k > 0 Rep-circline-mat (of-circline-rep H') = cor k *sm
Rep-circline-mat (of-circline-rep H)
      by auto
    thus circline-mat-eq H H'
      using mult-sm-inv-l[of -1 Rep-circline-mat H' cor k *sm Rep-circline-mat H]
      using mult-sm-inv-l[of -1 Rep-circline-mat H'(-(cor k)) *sm Rep-circline-mat
H]
      apply (auto simp add: of-circline-rep-def split: split-if-asm)
      apply (rule-tac x=k in exI, simp)
      apply (rule-tac x=-k in exI, simp)
      apply (rule-tac x=-k in exI, simp)
      apply (rule-tac x=k in exI, simp)
      done
  qed

lemma of-circline-of-ocircline:
  shows of-circline (of-ocircline H') = H' ∨ of-circline (of-ocircline H') = opposite-ocircline H'
  proof (cases pos-oriented H')
    case True
    thus ?thesis
      by auto
  next
    case False
    hence pos-oriented (opposite-ocircline H')

```

```

using pos-oriented
by auto
thus ?thesis
  using of-ocircline-opposite-ocircline[of H']
  using of-circline-of-ocircline-pos-oriented [of opposite-ocircline H']
  by auto
qed

```

11.10 Some special oriented circlines and discs

```

lift-definition mk-ocircline :: complex  $\Rightarrow$  complex  $\Rightarrow$  complex  $\Rightarrow$  complex  $\Rightarrow$  ocircline is mk-circline-rep
by (simp add: mk-circline-rep-def, rule-tac x=1 in exI, simp)

```

oriented unit circle and unit disc

```

lift-definition ounit-circle :: ocircline is unit-circle-rep
done

```

```

definition unit-disc = disc ounit-circle

```

```

lemma zero-in-unit-disc:  $0_h \in$  unit-disc
unfolding unit-disc-def disc-def
by (simp, transfer) (simp add: in-ocircline-rep-def Let-def vec-cnj-def)

```

```

lemma inf-notin-unit-disc:  $\infty_h \notin$  unit-disc
unfolding unit-disc-def disc-def
by (simp, transfer) (simp add: in-ocircline-rep-def Let-def vec-cnj-def)

```

```

lemma of-ocircline-ounit-circle [simp]: of-ocircline ounit-circle = unit-circle
by (transfer) (auto, rule-tac x=1 in exI, simp)

```

```

lemma of-circline-unit-circline [simp]: of-circline (unit-circle) = ounit-circle
by (transfer) (auto simp add: pos-oriented-rep-def of-circline-rep-def, rule-tac x=1
in exI, simp)

```

Oriented x axis and lower half plane

```

lift-definition o-x-axis :: ocircline is x-axis-rep
done

```

```

lemma o-x-axis-pos-oriented: pos-oriented o-x-axis
by transfer (simp add: pos-oriented-rep-def)

```

```

lemma of-ocircline-o-x-axis [simp]: of-ocircline o-x-axis = x-axis
by transfer (simp del: circline-mat-eq-def)

```

```

lemma of-circline-x-axis [simp]: of-circline x-axis = o-x-axis
using of-circline-of-ocircline-pos-oriented[of o-x-axis]
using o-x-axis-pos-oriented
by simp

```

lemma *ocircline-set-circline-set-x-axis*: *ocircline-set o-x-axis = circline-set x-axis*
by (*subst of-circline-x-axis[symmetric]*, *subst ocircline-set-circline-set, simp*)

lemma [*simp*]: $ii_h \notin disc\ o\text{-}x\text{-axis}$
unfolding *disc-def*
by *simp (transfer, simp add: in-ocircline-rep-def Let-def vec-cnj-def)*

lemma [*simp*]: $ii_h \in disc\ (\text{opposite-ocircline}\ o\text{-}x\text{-axis})$
unfolding *disc-def*
by *simp (transfer, simp add: in-ocircline-rep-def Let-def vec-cnj-def)*

11.11 Moebius action on oriented circlines and discs

lift-definition *moebius-ocircline* :: *moebius* \Rightarrow *ocircline* \Rightarrow *ocircline* **is** *moebius-circline-rep*
proof –

fix $M M' H H'$
assume *moebius-mat-eq M M' ocircline-mat-eq H H'*
thus *ocircline-mat-eq (moebius-circline-rep M H) (moebius-circline-rep M' H')*
by (*auto simp add: mat-inv-mult-sm complex-cnj*) (*rule-tac x=ka / Re (k * cnj k) in exI, auto simp add: complex-mult-cnj-cmod power2-eq-square, metis divide-pos-pos mult-eq-0-iff norm-mult zero-less-norm-iff*)
qed

lemma *moebius-circline-ocircline*:
moebius-circline M H = of-ocircline (moebius-ocircline M (of-circline H))
apply (*transfer*)
apply (*auto simp add: of-circline-rep-def*)
apply (*rule-tac x=1 in exI, simp*)
apply (*rule-tac x=-1 in exI, simp add: of-real-neg-numeral*)
done

lemma *moebius-ocircline-circline*:
moebius-ocircline M H = of-circline (moebius-circline M (of-ocircline H)) \vee
moebius-ocircline M H = opposite-ocircline (of-circline (moebius-circline M (of-ocircline H)))
apply (*transfer*)
apply (*auto simp add: of-circline-rep-def*)
apply (*rule-tac x=1 in exI, simp*)
apply (*erule-tac x=1 in allE, simp*)
done

lemma
inj-moebius-ocircline: inj (moebius-ocircline M)
unfolding *inj-on-def*
proof (*safe*)
fix $H H'$
assume *moebius-ocircline M H = moebius-ocircline M H'*
thus $H = H'$

```

proof (transfer)
fix M H H'
let ?M = Rep-moebius-mat M
let ?iM = mat-inv ?M
let ?H = Rep-circline-mat H and ?H' = Rep-circline-mat H'
assume ocircline-mat-eq (moebius-circline-rep M H) (moebius-circline-rep M H')
then obtain k where congruence ?iM ?H' = congruence ?iM (cor k *sm ?H)
k > 0
by auto
thus ocircline-mat-eq H H'
using Rep-moebius-mat[of M] inj-congruence[of ?iM ?H' cor k *sm ?H]
mat-det-inv[of ?M]
by auto
qed
qed

lemma moebius-ocircline-comp:
moebius-ocircline M1 (moebius-ocircline M2 H) = moebius-ocircline (moebius-comp M1 M2) H
proof (transfer)
fix M1 M2 H
show ocircline-mat-eq (moebius-circline-rep M1 (moebius-circline-rep M2 H))
(moebius-circline-rep (moebius-comp-rep M1 M2) H)
using congruence-congruence Rep-moebius-mat[of M1] Rep-moebius-mat[of M2]
by (simp add: mat-inv-mult-mm, rule-tac x=1 in exI, simp)
qed

lemma [simp]:
moebius-ocircline id-moebius H = H
proof transfer
fix H
show ocircline-mat-eq (moebius-circline-rep id-moebius-rep H) H
by (cases Rep-circline-mat H, simp) (rule-tac x=1 in exI, simp add: mat-adj-def mat-cnj-def)
qed

lemma moebius-ocircline-comp-inv[simp]:
moebius-ocircline (moebius-inv M) (moebius-ocircline M H) = H
by (subst moebius-ocircline-comp) simp

lemma moebius-circline-opposite-ocircline [simp]:
moebius-ocircline M (opposite-ocircline H) = opposite-ocircline (moebius-ocircline M H)
by transfer (auto, rule-tac x=1 in exI, simp)

lemma moebius-ocircline-set:
shows moebius-pt M ` ocircline-set H = ocircline-set (moebius-ocircline M H)

```

```

(is ?lhs = ?rhs)
proof-
  have moebius-pt M ` ocircline-set H = circline-set (moebius-circline M (of-ocircline
H))
    by (subst moebius-circline-set[symmetric]) simp
  thus ?thesis
    using moebius-ocircline-circline[of M H]
    by auto
qed

lemma moebius-disc:
  moebius-pt M ` (disc H) = disc (moebius-ocircline M H)
proof (safe)
  fix z
  assume z ∈ disc H
  thus moebius-pt M z ∈ disc (moebius-ocircline M H)
    unfolding disc-def
  proof (safe)
    assume in-ocircline H z
    thus in-ocircline (moebius-ocircline M H) (moebius-pt M z)
      unfolding in-ocircline-def
    proof (transfer)
      fix H z M
      assume in-ocircline-rep H z
      thus in-ocircline-rep (moebius-circline-rep M H) (moebius-pt-rep M z)
        using Rep-moebius-mat[of M] quad-form-congruence[of Rep-moebius-mat M
Rep-homo-coords z]
        by (simp add: in-ocircline-rep-def moebius-circline-rep-def Let-def)
    qed
  qed
next
  fix z
  assume z ∈ disc (moebius-ocircline M H)
  thus z ∈ moebius-pt M ` disc H
    unfolding disc-def
  proof(safe)
    assume in-ocircline (moebius-ocircline M H) z
    show z ∈ moebius-pt M ` Collect (in-ocircline H)
    proof
      show z = moebius-pt M (moebius-pt (moebius-inv M) z)
        using moebius-inv[of M] bij-moebius-pt[of M]
        by (simp add: bij-def) (metis surj-f-inv-f)
    next
      show moebius-pt (moebius-inv M) z ∈ Collect (in-ocircline H)
        using in-ocircline (moebius-ocircline M H) z
      proof (safe, transfer)
        fix M H z
        have congruence (mat-inv (mat-inv (Rep-moebius-mat M))) (congruence
(mat-inv (Rep-moebius-mat M)) (Rep-circline-mat H)) =
          Rep-circline-mat H

```

```

using Rep-moebius-mat[of M]
by (simp add: congruence-congruence-inv)
hence quad-form (Rep-homo-coords z) (congruence (mat-inv (Rep-moebius-mat
M)) (Rep-circline-mat H)) =
quad-form (mat-inv (Rep-moebius-mat M) *mv Rep-homo-coords z)
(Rep-circline-mat H)
using quad-form-congruence[of mat-inv (Rep-moebius-mat M) Rep-homo-coords
z congruence (mat-inv (Rep-moebius-mat M)) (Rep-circline-mat H)]
using Rep-moebius-mat[of M] mat-det-inv[of Rep-moebius-mat M]
by simp
moreover
assume in-ocircline-rep (moebius-circline-rep M H) z
ultimately
show in-ocircline-rep H (moebius-pt-rep (moebius-inv-rep M) z)
by (auto simp add: in-ocircline-rep-def Let-def)
qed
qed
qed
qed

lemma moebius-disc-compl:
moebius-pt M ` (disc-compl H) = disc-compl (moebius-ocircline M H)
proof (safe)
fix z
assume z ∈ disc-compl H
thus moebius-pt M z ∈ disc-compl (moebius-ocircline M H)
unfolding disc-compl-def
proof (safe)
assume out-ocircline H z
thus out-ocircline (moebius-ocircline M H) (moebius-pt M z)
proof (transfer)
fix H z M
assume out-ocircline-rep H z
thus out-ocircline-rep (moebius-circline-rep M H) (moebius-pt-rep M z)
using Rep-moebius-mat[of M] quad-form-congruence[of Rep-moebius-mat M
Rep-homo-coords z]
by (simp add: out-ocircline-rep-def moebius-circline-rep-def Let-def)
qed
qed
next
fix z
assume z ∈ disc-compl (moebius-ocircline M H)
thus z ∈ moebius-pt M ` disc-compl H
unfolding disc-compl-def
proof (safe)
assume out-ocircline (moebius-ocircline M H) z
show z ∈ moebius-pt M ` Collect (out-ocircline H)
proof
show z = moebius-pt M (moebius-pt (moebius-inv M) z)

```

```

using moebius-inv[of M] bij-moebius-pt[of M]
by (simp add: bij-def) (metis surj-f-inv-f)
next
show moebius-pt (moebius-inv M) z ∈ Collect (out-ocircline H)
  using <out-ocircline (moebius-ocircline M H) z>
proof (safe, transfer)
fix M H z
have congruence (mat-inv (mat-inv (Rep-moebius-mat M))) (congruence
(mat-inv (Rep-moebius-mat M)) (Rep-circline-mat H)) =
  Rep-circline-mat H
  using Rep-moebius-mat[of M]
  by (simp add: congruence-congruence-inv)
hence quad-form (Rep-homo-coords z) (congruence (mat-inv (Rep-moebius-mat
M)) (Rep-circline-mat H)) =
  quad-form (mat-inv (Rep-moebius-mat M) *mv Rep-homo-coords z)
(Rep-circline-mat H)
  using quad-form-congruence[of mat-inv (Rep-moebius-mat M) Rep-homo-coords
z congruence (mat-inv (Rep-moebius-mat M)) (Rep-circline-mat H)]
    using Rep-moebius-mat[of M] mat-det-inv[of Rep-moebius-mat M]
    by simp
moreover
assume out-ocircline-rep (moebius-circline-rep M H) z
ultimately
show out-ocircline-rep H (moebius-pt-rep (moebius-inv-rep M) z)
  by (auto simp add: out-ocircline-rep-def Let-def)
qed
qed
qed
qed

```

```

lemma similarity-preserves-lines:
assumes a ≠ 0
shows ∞h ∈ ocircline-set H ←→ ∞h ∈ ocircline-set (moebius-ocircline (similarity-moebius
a b) H) (is ?lhs = ?rhs)
proof
assume ?lhs
thus ?rhs
  using similarity-inf-fixed[OF ⟨a ≠ 0⟩, of b]
  by (subst moebius-ocircline-set[symmetric]) force
next
assume ?rhs
thus ?lhs
  using similarity-only-inf-to-inf[OF ⟨a ≠ 0⟩, of b]
  by (subst (asm) moebius-ocircline-set[symmetric]) (auto, metis)
qed

```

lemma similarity-preserve-orientation':

assumes $a \neq 0 M = \text{similarity-moebius } a b H' = \text{moebius-ocircline } M H \infty_h \notin \text{ocircline-set } H$
shows pos-oriented $H \rightarrow \text{pos-oriented } H'$
proof
have $\infty_h \notin \text{ocircline-set } H'$
using assms similarity-preserves-lines
by auto
assume pos-oriented H
hence $\infty_h \in \text{disc-compl } H$
using $\langle \infty_h \notin \text{ocircline-set } H \rangle \text{ pos-oriented-circle-inf}[\text{of } H] \text{ in-on-out}$
unfolding disc-def disc-compl-def ocircline-set-def
by auto
hence $\infty_h \in \text{disc-compl } H'$
using $\langle M = \text{similarity-moebius } a b \rangle \langle H' = \text{moebius-ocircline } M H \rangle$
using similarity-inf-fixed[$OF \langle a \neq 0 \rangle, of b$]
by (simp, subst moebius-disc-compl[symmetric], force)
thus pos-oriented H'
using pos-oriented-circle-inf[of $H'] disc-inter-disc-compl[of $H'] \langle \infty_h \notin \text{ocircline-set } H' \rangle$
by auto
qed$

lemma similarity-preserve-orientation:
assumes $a \neq 0 M = \text{similarity-moebius } a b H' = \text{moebius-ocircline } M H \infty_h \notin \text{ocircline-set } H$
shows pos-oriented $H \longleftrightarrow \text{pos-oriented } H'$
proof–
have $\infty_h \notin \text{ocircline-set } H'$
using assms similarity-preserves-lines
by auto

have $*: H = \text{moebius-ocircline } (-\text{similarity-moebius } a b) H'$
using $\langle H' = \text{moebius-ocircline } M H \rangle \langle M = \text{similarity-moebius } a b \rangle$
by simp
thus ?thesis
using $\langle a \neq 0 \rangle$
using similarity-preserve-orientation'[$OF \langle a \neq 0 \rangle \langle M = \text{similarity-moebius } a b \rangle \langle H' = \text{moebius-ocircline } M H \rangle \langle \infty_h \notin \text{ocircline-set } H \rangle$]
using similarity-preserve-orientation'[$OF -\text{similarity-moebius-inv}[of a b, OF \langle a \neq 0 \rangle] * \langle \infty_h \notin \text{ocircline-set } H' \rangle$]
by auto
qed

lemma $0_h \in \text{disc-compl } (\text{mk-ocircline } -1 (2*ii) (-2*ii) 1)$
unfolding disc-compl-def
by simp (transfer, simp add: out-ocircline-rep-def mk-circline-rep-def Abs-homo-coords-inverse Let-def Abs-circline-mat-inverse hermitean-def mat-adj-def mat-cnj-def vec-cnj-def complex-cnj)
lemma $\neg \text{pos-oriented } (\text{mk-ocircline } -1 (2*ii) (-2*ii) 1)$

```

by transfer (simp add: mk-circline-rep-def Abs-circline-mat-inverse hermitean-def
mat-adj-def mat-cnj-def complex-cnj pos-oriented-rep-def)
lemma circline-type (mk-circline -1 (2*ii) (-2*ii) 1) = -1
by transfer (simp add: mk-circline-rep-def Abs-circline-mat-inverse hermitean-def
mat-adj-def mat-cnj-def complex-cnj circline-type-rep-def)
lemma  $\theta_h \in disc\text{-compl}$  (mk-ocircline 1 (2*ii) (-2*ii) 1)
unfolding disc-compl-def
by simp (transfer, simp add: out-ocircline-rep-def mk-circline-rep-def Abs-homo-coords-inverse
Let-def Abs-circline-mat-inverse hermitean-def mat-adj-def mat-cnj-def vec-cnj-def
complex-cnj)
lemma pos-oriented (mk-ocircline 1 (2*ii) (-2*ii) 1)
by transfer (simp add: mk-circline-rep-def Abs-circline-mat-inverse hermitean-def
mat-adj-def mat-cnj-def complex-cnj pos-oriented-rep-def)
lemma circline-type (mk-circline 1 (2*ii) (-2*ii) 1) = -1
by transfer (simp add: mk-circline-rep-def Abs-circline-mat-inverse hermitean-def
mat-adj-def mat-cnj-def complex-cnj circline-type-rep-def)

lemma reciprocal-preserve-orientation:
assumes  $\theta_h \in disc\text{-compl } H M = reciprocal\text{-moebius } H' = moebius\text{-ocircline } M$ 
H
shows pos-oriented  $H'$ 
proof-
have  $\infty_h \in disc\text{-compl } H'$ 
using assms
by simp (subst moebius-disc-compl[symmetric], subst reciprocal-moebius[symmetric],
force)
thus pos-oriented  $H'$ 
using pos-oriented-circle-inf[of  $H'$ ] disc-inter-disc-compl[of  $H'$ ] disc-compl-inter-ocircline-set[of
 $H'$ ]
by auto
qed

lemma reciprocal-not-preserve-orientation:
assumes  $\theta_h \in disc H M = reciprocal\text{-moebius } H' = moebius\text{-ocircline } M H$ 
shows  $\neg$  pos-oriented  $H'$ 
proof-
have  $\infty_h \in disc H'$ 
using assms
by simp (subst moebius-disc[symmetric], subst reciprocal-moebius[symmetric],
force)
thus  $\neg$  pos-oriented  $H'$ 
using pos-oriented-circle-inf[of  $H'$ ] disc-inter-ocircline-set[of  $H'$ ]
by auto
qed

lemma pole-in-disc:
assumes  $M = mk\text{-moebius } a b c d c \neq 0 a*d - b*c \neq 0$ 
assumes is-pole  $M z z \in disc H H' = moebius\text{-ocircline } M H$ 
shows  $\neg$  pos-oriented  $H'$ 

```

proof–

```
let ?t1 = translation-moebius (a / c)
let ?rd = rotation-dilatation-moebius ((b * c - a * d) / (c * c))
let ?r = reciprocal-moebius
let ?t2 = translation-moebius (d / c)

have 0h = moebius-pt (translation-moebius (d/c)) z
  using pole-mk-moebius[of a b c d z] assms
  by simp

have z ∉ ocircline-set H
  using ⟨z ∈ disc H⟩ disc-inter-ocircline-set[of H]
  by auto
hence 0h ∉ ocircline-set (moebius-ocircline ?t2 H)
  using ⟨0h = moebius-pt ?t2 z⟩
  using inj-image-mem-iff[of moebius-pt ?t2 z ocircline-set H] bij-moebius-pt
  by (subst moebius-ocircline-set[symmetric]) (simp add: bij-def)
hence ∞h ∉ ocircline-set (moebius-ocircline (?r + ?t2) H)
  using reciprocal-homo-only-0-to-inf
  by (simp add: moebius-ocircline-comp[symmetric]) (subst moebius-ocircline-set[symmetric],
  subst reciprocal-moebius[symmetric], auto, metis)
hence ∞h ∉ ocircline-set (moebius-ocircline (?rd + ?r + ?t2) H)
  using ⟨a*d - b*c ≠ 0⟩ ⟨c ≠ 0⟩
  using similarity-preserves-lines
  unfolding rotation-dilatation-moebius-def
  by (simp add: moebius-ocircline-comp[symmetric])

have ¬ pos-oriented (moebius-ocircline (?r + ?t2) H)
  using pole-mk-moebius[of a b c d z] assms
  apply (simp add: moebius-ocircline-comp[symmetric])
  apply (subst reciprocal-not-preserve-orientation, simp-all)
  apply (subst moebius-disc[symmetric])
  apply force
  done
hence ¬ pos-oriented (moebius-ocircline (?rd + ?r + ?t2) H)
  using *
  using ⟨a*d - b*c ≠ 0⟩ ⟨c ≠ 0⟩
  unfolding rotation-dilatation-moebius-def
  by (simp add: moebius-ocircline-comp[symmetric]) (subst similarity-preserve-orientation[symmetric],
  simp-all)
hence ¬ pos-oriented (moebius-ocircline (?t1 + ?rd + ?r + ?t2) H)
  using **
  unfolding translation-moebius-def
  by (simp add: moebius-ocircline-comp[symmetric]) (subst similarity-preserve-orientation[symmetric],
  simp-all)

thus ?thesis
  using assms
  by simp (subst moebius-decomposition, auto simp add: moebius-ocircline-comp[symmetric])
```

qed

lemma pole-in-disc-compl:
assumes $M = \text{mk-moebius } a \ b \ c \ d \ c \neq 0 \ a*d - b*c \neq 0$
assumes is-pole $M z \ z \in \text{disc-compl } H \ H' = \text{moebius-ocircline } M \ H$
shows pos-oriented H'
proof–
let $?t1 = \text{translation-moebius } (a / c)$
let $?rd = \text{rotation-dilatation-moebius } ((b * c - a * d) / (c * c))$
let $?r = \text{reciprocal-moebius}$
let $?t2 = \text{translation-moebius } (d / c)$

have $\theta_h = \text{moebius-pt } (\text{translation-moebius } (d/c)) z$
using pole-mk-moebius[of a b c d z] assms
by simp

have $z \notin \text{ocircline-set } H$
using $\langle z \in \text{disc-compl } H \rangle \ \text{disc-compl-inter-ocircline-set}[of \ H]$
by auto
hence $\theta_h \notin \text{ocircline-set } (\text{moebius-ocircline } ?t2 \ H)$
using $\langle \theta_h = \text{moebius-pt } ?t2 \ z \rangle$
using inj-image-mem-iff[of moebius-pt ?t2 z ocircline-set H] bij-moebius-pt
by (subst moebius-ocircline-set[symmetric]) (simp add: bij-def)
hence $*: \infty_h \notin \text{ocircline-set } (\text{moebius-ocircline } (?r + ?t2) \ H)$
using reciprocal-homo-only-0-to-inf
by (simp add: moebius-ocircline-comp[symmetric]) (subst moebius-ocircline-set[symmetric],
subst reciprocal-moebius[symmetric], auto, metis)
hence $**: \infty_h \notin \text{ocircline-set } (\text{moebius-ocircline } (?rd + ?r + ?t2) \ H)$
using $\langle a*d - b*c \neq 0 \rangle \ \langle c \neq 0 \rangle$
using similarity-preserves-lines
unfolding rotation-dilatation-moebius-def
by (simp add: moebius-ocircline-comp[symmetric])

have pos-oriented ($\text{moebius-ocircline } (?r + ?t2) \ H$)
using pole-mk-moebius[of a b c d z] assms
apply (simp add: moebius-ocircline-comp[symmetric])
apply (subst reciprocal-preserve-orientation, simp-all)
apply (subst moebius-disc-compl[symmetric])
apply force
done
hence pos-oriented ($\text{moebius-ocircline } (?rd + ?r + ?t2) \ H$)
using *
using $\langle a*d - b*c \neq 0 \rangle \ \langle c \neq 0 \rangle$
unfolding rotation-dilatation-moebius-def
by (simp add: moebius-ocircline-comp[symmetric]) (subst similarity-preserve-orientation[symmetric],
simp-all)
hence pos-oriented ($\text{moebius-ocircline } (?t1 + ?rd + ?r + ?t2) \ H$)
using **
unfolding translation-moebius-def

by (*simp add: moebius-ocircline-comp[symmetric]*) (*subst similarity-preserve-orientation[symmetric]*, *simp-all*)

thus *?thesis*
using *assms*
by simp (*subst moebius-decomposition, auto simp add: moebius-ocircline-comp[symmetric]*)
qed

11.12 Oriented circlines uniqueness

lemma *ocircline-01inf*:

assumes $0_h \in \text{ocircline-set } H \wedge 1_h \in \text{ocircline-set } H \wedge \infty_h \in \text{ocircline-set } H$
shows $H = o\text{-}x\text{-axis} \vee H = \text{opposite-ocircline } o\text{-}x\text{-axis}$

proof –

have $0_h \in \text{circline-set } (\text{of-ocircline } H) \wedge 1_h \in \text{circline-set } (\text{of-ocircline } H) \wedge \infty_h \in \text{circline-set } (\text{of-ocircline } H)$
using *assms*
by simp
hence $\text{of-ocircline } H = x\text{-axis}$
using *unique-circline-01inf'*
by auto
thus $H = o\text{-}x\text{-axis} \vee H = \text{opposite-ocircline } o\text{-}x\text{-axis}$
by (*metis inj-of-ocircline of-ocircline-o-x-axis*)
qed

lemma *unique-ocircline-01inf*: $\exists! H. 0_h \in \text{ocircline-set } H \wedge 1_h \in \text{ocircline-set } H \wedge \infty_h \in \text{ocircline-set } H \wedge ii_h \notin \text{disc } H$

proof
show $0_h \in \text{ocircline-set } o\text{-}x\text{-axis} \wedge 1_h \in \text{ocircline-set } o\text{-}x\text{-axis} \wedge \infty_h \in \text{ocircline-set } o\text{-}x\text{-axis} \wedge ii_h \notin \text{disc } o\text{-}x\text{-axis}$
by (*simp add: ocircline-set-circline-set-x-axis*)
next
fix *H*
assume $0_h \in \text{ocircline-set } H \wedge 1_h \in \text{ocircline-set } H \wedge \infty_h \in \text{ocircline-set } H \wedge ii_h \notin \text{disc } H$
hence $0_h \in \text{ocircline-set } H \wedge 1_h \in \text{ocircline-set } H \wedge \infty_h \in \text{ocircline-set } H \wedge ii_h \notin \text{disc } H$
by auto
hence $H = o\text{-}x\text{-axis} \vee H = \text{opposite-ocircline } o\text{-}x\text{-axis}$
using *ocircline-01inf*
by simp
thus $H = o\text{-}x\text{-axis}$
using *(ii_h ∉ disc H)*
by auto
qed

lemma *unique-ocircline-set*:

assumes $A \neq B A \neq C B \neq C$
shows $\exists! H. \text{pos-oriented } H \wedge (A \in \text{ocircline-set } H \wedge B \in \text{ocircline-set } H \wedge C$

$\in \text{ocircline-set } H)$

proof –

obtain M **where** $*: \text{moebius-pt } M A = 0_h \text{ moebius-pt } M B = 1_h \text{ moebius-pt } M$

$C = \infty_h$

using $\text{ex-moebius-01inf}[\text{OF assms}]$

by auto

let $?iM = \text{moebius-pt}(\text{moebius-inv } M)$

have $**: ?iM 0_h = A \quad ?iM 1_h = B \quad ?iM \infty_h = C$

using $\text{bij-moebius-pt}[\text{of moebius-inv } M] *$

by $(\text{auto simp add: moebius-inv}) (\text{metis bij-def bij-moebius-pt inv-f-eq}) +$

let $?H = \text{moebius-ocircline}(\text{moebius-inv } M) \text{ o-x-axis}$

have $1: A \in \text{ocircline-set } ?H \quad B \in \text{ocircline-set } ?H \quad C \in \text{ocircline-set } ?H$

by $-(\text{subst moebius-ocircline-set[symmetric]}, \text{subst**[symmetric]}, \text{simp add: ocircline-set-circline-set-x-axis}) +$

have $2: \bigwedge H'. A \in \text{ocircline-set } H' \wedge B \in \text{ocircline-set } H' \wedge C \in \text{ocircline-set } H' \implies H' = ?H \vee H' = \text{opposite-ocircline } ?H$

proof –

fix H'

let $?H' = \text{ocircline-set } H' \text{ and } ?H'' = \text{ocircline-set}(\text{moebius-ocircline } M H')$

assume $A \in \text{ocircline-set } H' \wedge B \in \text{ocircline-set } H' \wedge C \in \text{ocircline-set } H'$

hence $\text{moebius-pt } M A \in ?H'' \text{ moebius-pt } M B \in ?H'' \text{ moebius-pt } M C \in ?H''$

using $\text{moebius-ocircline-set}[\text{of } M H']$

by auto

hence $0_h \in ?H'' \quad 1_h \in ?H'' \quad \infty_h \in ?H''$

using *

by auto

hence $\text{moebius-ocircline } M H' = \text{o-x-axis} \vee \text{moebius-ocircline } M H' = \text{opposite-ocircline}$
 o-x-axis

using ocircline-01inf

by auto

hence $\text{o-x-axis} = \text{moebius-ocircline } M H' \vee \text{o-x-axis} = \text{opposite-ocircline}$
 $(\text{moebius-ocircline } M H')$

by auto

thus $H' = ?H \vee H' = \text{opposite-ocircline } ?H$

proof

assume $*: \text{o-x-axis} = \text{moebius-ocircline } M H'$

show $H' = \text{moebius-ocircline}(\text{moebius-inv } M) \text{ o-x-axis} \vee H' = \text{opposite-ocircline}$
 $(\text{moebius-ocircline}(\text{moebius-inv } M) \text{ o-x-axis})$

by $(\text{rule disjI1}) (\text{subst } *, \text{simp})$

next

assume $*: \text{o-x-axis} = \text{opposite-ocircline}(\text{moebius-ocircline } M H')$

show $H' = \text{moebius-ocircline}(\text{moebius-inv } M) \text{ o-x-axis} \vee H' = \text{opposite-ocircline}$
 $(\text{moebius-ocircline}(\text{moebius-inv } M) \text{ o-x-axis})$

by $(\text{rule disjI2}) (\text{subst } *, \text{simp})$

qed

qed

show $?thesis$

```

proof (cases pos-oriented ?H)
  case True
  thus ?thesis
    unfolding Ex1-def
    proof (rule-tac x=moebius-ocircline (moebius-inv M) o-x-axis in exI, simp
add: 1, safe)
      fix y
      assume pos-oriented y A ∈ ocircline-set y B ∈ ocircline-set y C ∈ ocircline-set
y
      thus y = moebius-ocircline (moebius-inv M) o-x-axis
        using 2[of y] True
        by (auto simp add: pos-oriented-opposite-ocircline)
    qed
  next
  case False
  thus ?thesis
    unfolding Ex1-def
    proof (rule-tac x=opposite-ocircline (moebius-ocircline (moebius-inv M) o-x-axis)
in exI, simp add: 1 pos-oriented-opposite-ocircline, safe)
      fix y
      assume pos-oriented y A ∈ ocircline-set y B ∈ ocircline-set y C ∈ ocircline-set
y
      thus y = opposite-ocircline (moebius-ocircline (moebius-inv M) o-x-axis)
        using 2[of y] False
        by (auto simp add: pos-oriented-opposite-ocircline)
    qed
  qed
qed

```

```

definition chordal-circle-rep where
  chordal-circle-rep a r =
    (let (a1, a2) = Rep-homo-coords a in
     mk-circline-rep (4*a2*cnj a2 - (cor r)^2*(a1*cnj a1 + a2*cnj a2)) (-4*a1*cnj
a2) (-4*cnj a1*a2) (4*a1*cnj a1 - (cor r)^2*(a1*cnj a1 + a2*cnj a2)))
lemma [simp]: Rep-circline-mat (chordal-circle-rep a r) = (let (a1, a2) = Rep-homo-coords
a in
  (4*a2*cnj a2 - (cor r)^2*(a1*cnj a1 + a2*cnj a2), -4*a1*cnj a2, -4*cnj
a1*a2, 4*a1*cnj a1 - (cor r)^2*(a1*cnj a1 + a2*cnj a2)))
proof-
  obtain a1 a2 where aa: Rep-homo-coords a = (a1, a2)
    by (rule obtain-homo-coords)
  hence (4 * a2 * cnj a2 - (cor r)^2 * (a1 * cnj a1 + a2 * cnj a2), -4 * a1 *
cnj a2, -4 * cnj a1 * a2, 4 * a1 * cnj a1 - (cor r)^2 * (a1 * cnj a1 + a2 * cnj
a2)) ∈ {H. hermitean H ∧ H ≠ mat-zero}
    using Rep-homo-coords[of a]
    by (auto simp add: hermitean-def mat-adj-def mat-cnj-def complex-cnj power2-eq-square)
  thus ?thesis

```

```

using aa
by (simp add: chordal-circle-rep-def split-def Let-def mk-circline-rep-def Abs-circline-mat-inverse)
qed

lift-definition chordal-circle :: complex-homo  $\Rightarrow$  real  $\Rightarrow$  circline is chordal-circle-rep
proof-
  fix a b r
  obtain a1 a2 where aa: Rep-homo-coords a = (a1, a2)
    by (rule obtain-homo-coords)
  obtain b1 b2 where bb: Rep-homo-coords b = (b1, b2)
    by (rule obtain-homo-coords)
  assume a  $\approx$  b
  then obtain k where Rep-homo-coords b = (k * a1, k * a2) k  $\neq$  0
    using aa bb
    by auto
  moreover
  have cor (Re (k * cnj k)) = k * cnj k
    by (metis complex-In-mult-cnj-zero complex-of-real-Re)
  ultimately
  show circline-mat-eq (chordal-circle-rep a r) (chordal-circle-rep b r)
    using aa bb
    by (auto simp add: complex-cnj) (rule-tac x=Re (k*cnj k) in exI, auto simp
add: field-simps)
qed

lemma sqrt-1-plus-square:  $\sqrt{1 + a^2} \neq 0$ 
  by (smt real-sqrt-less-mono real-sqrt-zero realpow-square-minus-le)

lemma
  assumes dist-homo z a = r
  shows z  $\in$  circline-set (chordal-circle a r)
  proof (cases a  $\neq$   $\infty_h$ )
    case True
    then obtain a' where a = of-complex a'
      using inf-homo-or-complex-homo[of a]
      by auto
    let ?A = 4 - (cor r)2 * (1 + (a'*cnj a')) and ?B = -4*a' and ?C=-4*cnj
    a' and ?D = 4*a'*cnj a' - (cor r)2 * (1 + (a'*cnj a'))
    have hh: (?A, ?B, ?C, ?D)  $\in$  {H. hermitean H  $\wedge$  H  $\neq$  mat-zero}
      by (auto simp add: hermitean-def mat-adj-def mat-cnj-def complex-cnj power2-eq-square)
    hence *: chordal-circle (of-complex a') r = mk-circline ?A ?B ?C ?D
      by (transfer, simp add: mk-circline-rep-def Abs-circline-mat-inverse) (rule-tac
x=1 in exI, simp)

    show ?thesis
    proof (cases z  $\neq$   $\infty_h$ )
      case True
      then obtain z' where z = of-complex z'
        using inf-homo-or-complex-homo[of z] inf-homo-or-complex-homo[of a]

```

```

    by auto
have 2 * cmod (z' - a') / (sqrt (1 + (cmod z')^2) * sqrt (1 + (cmod a')^2)) =
r
  using dist-homo-finite[of z' a'] assms <z = of-complex z'> <a = of-complex a'>
  by auto
hence 4 * (cmod (z' - a'))^2 / ((1 + (cmod z')^2) * (1 + (cmod a')^2)) = r^2
  by (auto simp add: power-mult-distrib power-divide field-simps)
moreover
have sqrt (1 + (cmod z')^2) ≠ 0 sqrt (1 + (cmod a')^2) ≠ 0
  using sqrt-1-plus-square
  by simp+
hence (1 + (cmod z')^2) * (1 + (cmod a')^2) ≠ 0
  by simp
ultimately
have 4 * (cmod (z' - a'))^2 = r^2 * ((1 + (cmod z')^2) * (1 + (cmod a')^2))
  by (simp add: field-simps)
hence 4 * Re ((z' - a')*cnj (z' - a')) = r^2 * (1 + Re (z'*cnj z')) * (1 +
Re (a'*cnj a'))
  by ((subst cmod-square[symmetric])+, simp)
hence 4 * (Re(z'*cnj z') - Re(a'*cnj z') - Re(cnj a'*z') + Re(a'*cnj a')) =
r^2 * (1 + Re (z'*cnj z')) * (1 + Re (a'*cnj a'))
  by (simp add: complex-cnj field-simps)
hence Re (?A * z' * cnj z' + ?B * cnj z' + ?C * z' + ?D) = 0
  by (simp add: power2-eq-square field-simps)
hence ?A * z' * cnj z' + ?B * cnj z' + ?C * z' + ?D = 0
  by (subst complex-eq-if-Re-eq) (auto simp add: power2-eq-square)
hence (cnj z' * ?A + ?C) * z' + (cnj z' * ?B + ?D) = 0
  by algebra
hence z ∈ circline-set (mk-circline ?A ?B ?C ?D)
  using <z = of-complex z'> hh
  unfolding circline-set-def
  by simp (transfer, simp add: of-complex-coords-def Abs-homo-coords-inverse
on-circline-rep-def Let-def Abs-circline-mat-inverse mk-circline-rep-def vec-cnj-def)
thus ?thesis
  using *
  by (subst <a = of-complex a'>) simp
next
case False
hence 2 / sqrt (1 + (cmod a')^2) = r
  using assms <a = of-complex a'>
  using dist-homo-infinite2[of a']
  by simp
moreover
have sqrt (1 + (cmod a')^2) ≠ 0
  using sqrt-1-plus-square
  by simp
ultimately
have 2 = r * sqrt (1 + (cmod a')^2)
  by (simp add: field-simps)

```

```

hence  $4 = (r * \sqrt{1 + (\text{cmod } a')^2})^2$ 
  by simp
hence  $4 = r^2 * (1 + (\text{cmod } a')^2)$ 
  by (simp add: power-mult-distrib)
hence  $\text{Re}(4 - (\text{cor } r)^2 * (1 + (a' * \text{cnj } a'))) = 0$ 
  by (subst (asm) cmod-square) (simp add: field-simps power2-eq-square)
hence  $4 - (\text{cor } r)^2 * (1 + (a' * \text{cnj } a')) = 0$ 
  by (subst complex-eq-if-Re-eq) (auto simp add: power2-eq-square)
hence circline-A0 (mk-circline ?A ?B ?C ?D)
  using hh
by simp (transfer, simp add: circline-A0-rep-def mk-circline-rep-def Abs-circline-mat-inverse)
hence  $z \in \text{circline-set}(\text{mk-circline } ?A ?B ?C ?D)$ 
  using inf-in-circline-set[of mk-circline ?A ?B ?C ?D]
  using  $\neg z \neq \infty_h$ 
  by simp
thus ?thesis
  using *
  by (subst (a = of-complex a')) simp
qed
next
case False
let ?A =  $-(\text{cor } r)^2$  and ?B = 0 and ?C = 0 and ?D =  $4 - (\text{cor } r)^2$ 
have hh:  $(?A, ?B, ?C, ?D) \in \{H. \text{hermitean } H \wedge H \neq \text{mat-zero}\}$ 
  by (auto simp add: hermitean-def mat-adj-def mat-cnj-def complex-cnj power2-eq-square)
hence *: chordal-circle a r = mk-circline ?A ?B ?C ?D
  using  $\neg a \neq \infty_h$ 
  by simp (transfer, simp add: mk-circline-rep-def Abs-circline-mat-inverse,
rule-tac x=1 in exI, simp)

show ?thesis
proof (cases z =  $\infty_h$ )
  case True
  show ?thesis
    using assms (z =  $\infty_h$ )  $\neg a \neq \infty_h$ 
    using * hh
    by (simp, subst inf-in-circline-set, transfer, simp add: circline-A0-rep-def
mk-circline-rep-def Abs-circline-mat-inverse)
  next
  case False
  then obtain z' where z = of-complex z'
    using inf-homo-or-complex-homo[of z]
    by auto
  have  $2 / \sqrt{1 + (\text{cmod } z')^2} = r$ 
    using assms (z = of-complex z')  $\neg a \neq \infty_h$ 
    using dist-homo-infinite2[of z']
    by simp
  moreover
  have  $\sqrt{1 + (\text{cmod } z')^2} \neq 0$ 
    using sqrt-1-plus-square

```

```

by simp
ultimately
have  $\sqrt{2} = r * \sqrt{1 + (\text{cmod } z')^2}$ 
  by (simp add: field-simps)
hence  $4 = (r * \sqrt{1 + (\text{cmod } z')^2})^2$ 
  by simp
hence  $4 = r^2 * (1 + (\text{cmod } z')^2)$ 
  by (simp add: power-mult-distrib)
hence  $\text{Re}(4 - (\text{cor } r)^2 * (1 + (z' * \text{cnj } z')))) = 0$ 
  by (subst (asm) cmod-square) (simp add: field-simps power2-eq-square)
hence  $-(\text{cor } r)^2 * z' * \text{cnj } z' + 4 - (\text{cor } r)^2 = 0$ 
  by (subst complex-eq-if-Re-eq) (auto simp add: power2-eq-square field-simps)
hence  $z \in \text{circline-set}(\text{mk-circline } ?A ?B ?C ?D)$ 
  using hh
  unfolding circline-set-def
  by (subst (z = of-complex z'), simp) (transfer, auto simp add: on-circline-rep-def
Let-def mk-circline-rep-def Abs-circline-mat-inverse vec-cnj-def field-simps)
thus ?thesis
  using *
  by simp
qed
qed

lemma [simp]:  $\sqrt{4} = 2$ 
proof-
  have  $\sqrt{2^2} = 2$ 
    by (metis abs-numeral real-sqrt-abs)
  thus ?thesis
    by simp
qed

lemma
assumes  $z \in \text{circline-set}(\text{chordal-circle } a r) \quad r \geq 0$ 
shows  $\text{dist-homo } z a = r$ 
proof (cases a =  $\infty_h$ )
  case False
  then obtain a' where  $a = \text{of-complex } a'$ 
    using inf-homo-or-complex-homo[of a]
    by auto

  let ?A =  $4 - (\text{cor } r)^2 * (1 + (a' * \text{cnj } a'))$  and ?B =  $-4 * a'$  and ?C =  $-4 * \text{cnj } a'$  and ?D =  $4 * a' * \text{cnj } a' - (\text{cor } r)^2 * (1 + (a' * \text{cnj } a'))$ 
  have hh:  $(?A, ?B, ?C, ?D) \in \{H. \text{hermitean } H \wedge H \neq \text{mat-zero}\}$ 
    by (auto simp add: hermitean-def mat-adj-def mat-cnj-def complex-cnj power2-eq-square)
  hence  $*: \text{chordal-circle}(\text{of-complex } a') r = \text{mk-circline } ?A ?B ?C ?D$ 
    by (transfer, simp add: mk-circline-rep-def Abs-circline-mat-inverse) (rule-tac
      x=1 in exI, simp)

  show ?thesis

```

```

proof (cases  $z = \infty_h$ )
  case False
    then obtain  $z'$  where  $z = \text{of-complex } z'$ 
      using inf-homo-or-complex-homo[of  $z$ ] inf-homo-or-complex-homo[of  $a$ ]
      by auto
    hence  $z \in \text{circline-set}(\text{mk-circline } ?A ?B ?C ?D)$ 
      using assms ‹ $a = \text{of-complex } a'$ › *
      by simp
    hence  $(\text{cnj } z' * ?A + ?C) * z' + (\text{cnj } z' * ?B + ?D) = 0$ 
      using hh
      unfolding circline-set-def
      by (subst (asm) ‹ $z = \text{of-complex } z'$ ›, simp) (transfer, simp add: on-circline-rep-def)
    Let-def mk-circline-rep-def Abs-circline-mat-inverse vec-cnj-def)
    hence  $?A * z' * \text{cnj } z' + ?B * \text{cnj } z' + ?C * z' + ?D = 0$ 
      by algebra
    hence  $\text{Re} (?A * z' * \text{cnj } z' + ?B * \text{cnj } z' + ?C * z' + ?D) = 0$ 
      by (simp add: power2-eq-square field-simps)
    hence  $4 * \text{Re} ((z' - a') * \text{cnj } (z' - a')) = r^2 * (1 + \text{Re}(z' * \text{cnj } z')) * (1 + \text{Re}(a' * \text{cnj } a'))$ 
      by (simp add: complex-cnj field-simps power2-eq-square)
    hence  $4 * (\text{cmod}(z' - a'))^2 = r^2 * ((1 + (\text{cmod } z')^2) * (1 + (\text{cmod } a')^2))$ 
      by (subst cmod-square)+ simp
    moreover
      have  $\sqrt{1 + (\text{cmod } z')^2} \neq 0$   $\sqrt{1 + (\text{cmod } a')^2} \neq 0$ 
        using sqrt-1-plus-square
        by simp+
      hence  $(1 + (\text{cmod } z')^2) * (1 + (\text{cmod } a')^2) \neq 0$ 
        by simp
    ultimately
      have  $4 * (\text{cmod}(z' - a'))^2 / ((1 + (\text{cmod } z')^2) * (1 + (\text{cmod } a')^2)) = r^2$ 
        by (simp add: field-simps)
      hence  $2 * \text{cmod}(z' - a') / (\sqrt{1 + (\text{cmod } z')^2} * \sqrt{1 + (\text{cmod } a')^2})$ 
       $= r$ 
        using ‹ $r \geq 0$ ›
        by (subst (asm) real-sqrt-eq-iff[symmetric]) (simp add: real-sqrt-mult real-sqrt-divide)
    thus ?thesis
      using ‹ $z = \text{of-complex } z'$ › ‹ $a = \text{of-complex } a'$ ›
      using dist-homo-finite[of  $z' a'$ ]
      by simp
  next
    case True
    have  $z \in \text{circline-set}(\text{mk-circline } ?A ?B ?C ?D)$ 
      using assms ‹ $a = \text{of-complex } a'$ › *
      by simp
    hence circline-A0 (mk-circline ?A ?B ?C ?D)
      using inf-in-circline-set[mk-circline ?A ?B ?C ?D]
      using ‹ $z = \infty_h$ ›
      by simp
    hence  $4 - (\text{cor } r)^2 * (1 + (a' * \text{cnj } a')) = 0$ 

```

```

using hh
by transfer (simp add: circline-A0-rep-def mk-circline-rep-def Abs-circline-mat-inverse)
hence Re (4 - (cor r)2 * (1 + (a' * cnj a'))) = 0
  by simp
hence 4 = r2 * (1 + (cmod a')2)
  by (subst cmod-square) (simp add: power2-eq-square)
hence 2 = r * sqrt (1 + (cmod a')2)
  using ‹r ≥ 0›
  by (subst (asm) real-sqrt-eq-iff[symmetric]) (simp add: real-sqrt-mult)
moreover
have sqrt (1 + (cmod a')2) ≠ 0
  using sqrt-1-plus-square
  by simp
ultimately
have 2 / sqrt (1 + (cmod a')2) = r
  by (simp add: field-simps)
thus ?thesis
  using ‹a = of-complex a'› ‹z = ∞h›
  using dist-homo-infinite2[of a']
  by simp
qed
next
case True
let ?A = -(cor r)2 and ?B = 0 and ?C = 0 and ?D = 4 - (cor r)2
have hh: (?A, ?B, ?C, ?D) ∈ {H. hermitean H ∧ H ≠ mat-zero}
  by (auto simp add: hermitean-def mat-adj-def mat-cnj-def complex-cnj power2-eq-square)
hence *: chordal-circle a r = mk-circline ?A ?B ?C ?D
  using ‹a = ∞h›
  by simp (transfer, simp add: mk-circline-rep-def Abs-circline-mat-inverse,
rule-tac x=1 in exI, simp)

show ?thesis
proof (cases z = ∞h)
  case True
  thus ?thesis
    using ‹a = ∞h› assms * hh
    by simp (subst (asm) inf-in-circline-set, transfer, simp add: circline-A0-rep-def
mk-circline-rep-def Abs-circline-mat-inverse)
  next
  case False
  then obtain z' where z = of-complex z'
    using inf-homo-or-complex-homo[of z] inf-homo-or-complex-homo[of a]
    by auto
  hence z ∈ circline-set (mk-circline ?A ?B ?C ?D)
    using assms *
    by simp
  hence - (cor r)2 * z'*cnj z' + 4 - (cor r)2 = 0
    using hh
    unfolding circline-set-def

```

```

apply (subst (asm) ⟨z = of-complex z'⟩)
  by (simp, transfer) (simp add: on-circline-rep-def mk-circline-rep-def Let-def
vec-cnj-def Abs-circline-mat-inverse, algebra)
  hence 4 - (cor r)2 * (1 + (z'*cnj z')) = 0
    by (simp add: field-simps)
  hence Re (4 - (cor r)2 * (1 + (z' * cnj z'))) = 0
    by simp
  hence 4 = r2 * (1 + (cmod z')2)
    by (subst cmod-square) (simp add: power2-eq-square)
  hence 2 = r * sqrt (1 + (cmod z')2)
    using ⟨r ≥ 0⟩
    by (subst (asm) real-sqrt-eq-iff[symmetric]) (simp add: real-sqrt-mult)
moreover
have sqrt (1 + (cmod z')2) ≠ 0
  using sqrt-1-plus-square
  by simp
ultimately
have 2 / sqrt (1 + (cmod z')2) = r
  by (simp add: field-simps)
thus ?thesis
  using ⟨z = of-complex z'⟩ ⟨a = ∞b⟩
  using dist-homo-infinite2[of z']
  by simp
qed
qed

```

lemma chordal-circle-radius-positive:

```

assumes hermitean (A, B, C, D) Re (mat-det (A, B, C, D)) ≤ 0 B ≠ 0
dsc = sqrt(Re ((D-A)2 + 4 * (B*cnj B))) a1 = (A - D + cor dsc) / (2 * C)
a2 = (A - D - cor dsc) / (2 * C)
shows Re (A*a1/B) ≥ -1 ∧ Re (A*a2/B) ≥ -1
proof-
  from assms have is-real A is-real D C = cnj B
    using hermitean-elems
    by auto
  have *: A*a1/B = ((A - D + cor dsc) / (2 * (B * cnj B))) * A
    using ⟨B ≠ 0⟩ ⟨C = cnj B⟩ ⟨a1 = (A - D + cor dsc) / (2 * C)⟩
    by (simp add: field-simps) algebra
  have **: A*a2/B = ((A - D - cor dsc) / (2 * (B * cnj B))) * A
    using ⟨B ≠ 0⟩ ⟨C = cnj B⟩ ⟨a2 = (A - D - cor dsc) / (2 * C)⟩
    by (simp add: field-simps) algebra
  have dsc ≥ 0
  proof-
    have 0 ≤ Re ((D - A)2) + 4 * Re ((cor (cmod B))2)
      using ⟨is-real A⟩ ⟨is-real D⟩
      by (subst cor-squared, subst Re-complex-of-real) (simp add: power2-eq-square)
    thus ?thesis
      using ⟨dsc = sqrt(Re ((D-A)2 + 4*(B*cnj B)))⟩
      by (subst (asm) complex-mult-cnj-cmod) simp
  qed

```

```

qed
hence  $\operatorname{Re}(A - D - \operatorname{cor dsc}) \leq \operatorname{Re}(A - D + \operatorname{cor dsc})$ 
  by simp
moreover
have  $\operatorname{Re}(2 * (B * \operatorname{cnj} B)) > 0$ 
  using  $\langle B \neq 0 \rangle$ 
  by (subst complex-mult-cnj-cmod, simp add: power2-eq-square) (metis norm-eq-zero
not-real-square-gt-zero)
ultimately
have  $\operatorname{xxx}: \operatorname{Re}(A - D + \operatorname{cor dsc}) / \operatorname{Re}(2 * (B * \operatorname{cnj} B)) \geq \operatorname{Re}(A - D - \operatorname{cor dsc}) / \operatorname{Re}(2 * (B * \operatorname{cnj} B))$  (is ?lhs  $\geq$  ?rhs)
  by (metis divide-right-mono less-eq-real-def)

have  $\operatorname{Re} A * \operatorname{Re} D \leq \operatorname{Re}(B * \operatorname{cnj} B)$ 
  using  $\langle \operatorname{Re}(\operatorname{mat-det}(A, B, C, D)) \leq 0 \rangle \langle C = \operatorname{cnj} B \rangle \langle \operatorname{is-real} A \rangle \langle \operatorname{is-real} D \rangle$ 
  by simp

show ?thesis
proof (cases  $\operatorname{Re} A > 0$ )
  case True
  hence  $\operatorname{Re}(A * a1/B) \geq \operatorname{Re}(A * a2/B)$ 
    using * **  $\langle \operatorname{Re}(2 * (B * \operatorname{cnj} B)) > 0 \rangle \langle B \neq 0 \rangle \langle \operatorname{is-real} A \rangle \langle \operatorname{is-real} D \rangle$  xxx
    using mult-right-mono[of ?rhs ?lhs  $\operatorname{Re} A$ ]
    apply simp
    apply (subst Re-divide-real, simp, simp)
    apply (subst Re-divide-real, simp, simp)
    apply (subst Re-mult-real, simp) +
    apply simp
    done
  moreover
  have  $\operatorname{Re}(A * a2/B) \geq -1$ 
  proof -
    from  $\operatorname{Re} A * \operatorname{Re} D \leq \operatorname{Re}(B * \operatorname{cnj} B)$ 
    have  $\operatorname{Re}(A^2) \leq \operatorname{Re}(B * \operatorname{cnj} B) + \operatorname{Re}((A - D) * A)$ 
      using  $\langle \operatorname{Re} A > 0 \rangle \langle \operatorname{is-real} A \rangle \langle \operatorname{is-real} D \rangle$ 
      by (simp add: power2-eq-square field-simps)
    have  $1 \leq \operatorname{Re}(B * \operatorname{cnj} B) / \operatorname{Re}(A^2) + \operatorname{Re}(A - D) / \operatorname{Re} A$ 
      using  $\langle \operatorname{Re} A > 0 \rangle \langle \operatorname{is-real} A \rangle \langle \operatorname{is-real} D \rangle$ 
      using divide-right-mono[OF  $\operatorname{Re}(A^2) \leq \operatorname{Re}(B * \operatorname{cnj} B) + \operatorname{Re}((A - D) * A)$ ],
of  $\operatorname{Re}(A^2)$ ]
      by (simp add: power2-eq-square add-divide-distrib)
    have  $4 * \operatorname{Re}(B * \operatorname{cnj} B) \leq 4 * (\operatorname{Re}(B * \operatorname{cnj} B))^2 / \operatorname{Re}(A^2) + 2 * \operatorname{Re}(A - D) / \operatorname{Re} A * 2 * \operatorname{Re}(B * \operatorname{cnj} B)$ 
      using mult-right-mono[OF  $1 \leq \operatorname{Re}(B * \operatorname{cnj} B) / \operatorname{Re}(A^2) + \operatorname{Re}(A - D) / \operatorname{Re} A$ , of  $4 * \operatorname{Re}(B * \operatorname{cnj} B)$ ]
      by (simp add: distrib-right) (simp add: power2-eq-square field-simps)
  moreover
  have  $A \neq 0$ 
    using  $\langle \operatorname{Re} A > 0 \rangle$ 

```

```

by auto
hence  $4 * (Re(B * cnj B))^2 / Re(A^2) = Re(4 * (B * cnj B)^2 / A^2)$ 
  using Re-divide-real[of  $A^2$   $4 * (B * cnj B)^2$ ] ⟨Re  $A > 0$ ⟩ ⟨is-real  $A$ ⟩
    by (auto simp add: power2-eq-square)
moreover
have  $2 * Re(A - D) / Re A * 2 * Re(B * cnj B) = Re(2 * (A - D) / A * 2 * B * cnj B)$ 
  using ⟨is-real  $A$ ⟩ ⟨is-real  $D$ ⟩ ⟨ $A \neq 0$ ⟩
  using Re-divide-real[of  $A$   $(4 * A - 4 * D) * B * cnj B$ ]
    by (simp add: field-simps)
ultimately
have  $Re((D - A)^2 + 4 * B * cnj B) \leq Re((A - D)^2 + 4 * (B * cnj B)^2 / A^2 + 2 * (A - D) / A * 2 * B * cnj B)$ 
  by (simp add: field-simps power2-eq-square)
hence  $Re((D - A)^2 + 4 * B * cnj B) \leq Re((A - D) + 2 * B * cnj B / A)^2$ 
  using ⟨ $A \neq 0$ ⟩
  by (subst power2-sum) (simp add: power2-eq-square field-simps)
hence  $dsc \leq \sqrt{Re((A - D) + 2 * B * cnj B / A)^2})$ 
  using ⟨ $dsc = \sqrt{Re((D - A)^2 + 4 * (B * cnj B))}$ ⟩
    by simp
moreover
have  $Re((A - D) + 2 * B * cnj B / A)^2 = (Re((A - D) + 2 * B * cnj B / A))^2$ 
  using ⟨is-real  $A$ ⟩ ⟨is-real  $D$ ⟩ div-reals
  by (simp add: power2-eq-square)
ultimately
have  $dsc \leq |Re(A - D + 2 * B * cnj B / A)|$ 
  by simp
moreover
have  $Re(A - D + 2 * B * cnj B / A) \geq 0$ 
proof-
  have  $Re(A^2 + B * cnj B) \geq 0$ 
    using ⟨is-real  $A$ ⟩
    by (simp add: power2-eq-square)
  hence  $Re(A^2 + 2 * B * cnj B - A * D) \geq 0$ 
    using ⟨ $Re A * Re D \leq Re(B * cnj B)$ ⟩
    using ⟨is-real  $A$ ⟩ ⟨is-real  $D$ ⟩
    by simp
  show ?thesis
    using divide-right-mono[OF ⟨Re  $(A^2 + 2 * B * cnj B - A * D) \geq 0$ ⟩, of  $Re A$ ] ⟨Re  $A > 0$ ⟩ ⟨is-real  $A$ ⟩ ⟨ $A \neq 0$ ⟩
      by (simp add: add-divide-distrib diff-divide-distrib del: complex-Re-mult)
      (subst Re-divide-real, auto simp add: power2-eq-square field-simps)
qed
ultimately
have  $dsc \leq Re(A - D + 2 * B * cnj B / A)$ 
  by simp
hence  $-Re(2 * (B * cnj B) / A) \leq Re((A - D - cor dsc))$ 

```

```

by (simp add: field-simps)
hence  $- (Re (2 * (B * cnj B)) / Re A) \leq Re (A - D - cor dsc)$ 
  using (is-real A) (A ≠ 0)
  by (subst (asm) Re-divide-real, auto)
from divide-right-mono[OF this, of Re (2 * B * cnj B)]
have  $- 1 / Re A \leq Re (A - D - cor dsc) / Re (2 * B * cnj B)$ 
  using (Re A > 0) (B ≠ 0) (A ≠ 0) (0 < Re (2 * (B * cnj B)))
  by (simp add: field-simps del: complex-Re-mult)
from mult-right-mono[OF this, of Re A]
show ?thesis
  using (is-real A) (is-real D) (B ≠ 0) (Re A > 0) (A ≠ 0)
  apply (subst **)
  apply (subst Re-mult-real, simp add: div-reals)
  apply (subst Re-divide-real, simp, simp)
  apply (simp add: field-simps)
  done
qed
ultimately
show ?thesis
  by simp
next
case False
show ?thesis
proof (cases Re A < 0)
  case True
  hence  $Re (A*a1/B) \leq Re (A*a2/B)$ 
    using * ** (Re (2 * (B * cnj B)) > 0) (B ≠ 0) (is-real A) (is-real D) xxx
    using mult-right-mono-neg[of ?rhs ?lhs Re A]
    apply simp
    apply (subst Re-divide-real, simp, simp)
    apply (subst Re-divide-real, simp, simp)
    apply (subst Re-mult-real, simp) +
    apply simp
    done
  moreover
  have  $Re (A*a1/B) \geq -1$ 
  proof-
    from (Re A * Re D ≤ Re (B*cnj B))
    have  $Re (A^2) \leq Re (B*cnj B) - Re ((D - A)*A)$ 
      using (Re A < 0) (is-real A) (is-real D)
      by (simp add: power2-eq-square field-simps)
    hence  $1 \leq Re (B*cnj B) / Re (A^2) - Re (D - A) / Re A$ 
      using (Re A < 0) (is-real A) (is-real D)
      by (simp add: power2-eq-square diff-divide-distrib)
    have  $4 * Re(B*cnj B) \leq 4 * (Re (B*cnj B))^2 / Re (A^2) - 2 * Re (D - A) / Re A * 2 * Re(B*cnj B)$ 
      using mult-right-mono[OF (1 ≤ Re (B*cnj B) / Re (A^2) - Re (D - A))]

```

```

/  $\operatorname{Re} A$ , of  $4 * \operatorname{Re}(B * \operatorname{cnj} B)$ ]
  by (simp add: left-diff-distrib) (simp add: power2-eq-square field-simps)
  moreover
    have  $A \neq 0$ 
      using ⟨ $\operatorname{Re} A < 0$ ⟩
      by auto
    hence  $4 * (\operatorname{Re}(B * \operatorname{cnj} B))^2 / \operatorname{Re}(A^2) = \operatorname{Re}(4 * (B * \operatorname{cnj} B)^2 / A^2)$ 
      using Re-divide-real[of  $A^2 4 * (B * \operatorname{cnj} B)^2$ ] ⟨ $\operatorname{Re} A < 0$ ⟩ ⟨is-real A⟩
      by (auto simp add: power2-eq-square)
  moreover
    have  $2 * \operatorname{Re}(D - A) / \operatorname{Re} A * 2 * \operatorname{Re}(B * \operatorname{cnj} B) = \operatorname{Re}(2 * (D - A) / A * 2 * B * \operatorname{cnj} B)$ 
      using ⟨is-real A⟩ ⟨is-real D⟩ ⟨ $A \neq 0$ ⟩
      using Re-divide-real[of A  $(4 * D - 4 * A) * B * \operatorname{cnj} B$ ]
      by (simp add: field-simps)
  ultimately
    have  $\operatorname{Re}((D - A)^2 + 4 * B * \operatorname{cnj} B) \leq \operatorname{Re}((D - A)^2 + 4 * (B * \operatorname{cnj} B)^2 / A^2 - 2 * (D - A) / A * 2 * B * \operatorname{cnj} B)$ 
      by (simp add: field-simps power2-eq-square)
    hence  $\operatorname{Re}((D - A)^2 + 4 * B * \operatorname{cnj} B) \leq \operatorname{Re}(((D - A) - 2 * B * \operatorname{cnj} B / A)^2)$ 
      using ⟨ $A \neq 0$ ⟩
      by (subst power2-diff) (simp add: power2-eq-square field-simps)
    hence  $dsc \leq \sqrt{\operatorname{Re}(((D - A) - 2 * B * \operatorname{cnj} B / A)^2)}$ 
      using ⟨ $dsc = \sqrt{\operatorname{Re}((D - A)^2 + 4 * (B * \operatorname{cnj} B))}$ ⟩
      by simp
  moreover
    have  $\operatorname{Re}(((D - A) - 2 * B * \operatorname{cnj} B / A)^2) = (\operatorname{Re}((D - A) - 2 * B * \operatorname{cnj} B / A))^2$ 
      using ⟨is-real A⟩ ⟨is-real D⟩ div-reals
      by (simp add: power2-eq-square)
  ultimately
    have  $dsc \leq |\operatorname{Re}(D - A - 2 * B * \operatorname{cnj} B / A)|$ 
      by simp
  moreover
    have  $\operatorname{Re}(D - A - 2 * B * \operatorname{cnj} B / A) \geq 0$ 
  proof-
    have  $\operatorname{Re}(A^2 + B * \operatorname{cnj} B) \geq 0$ 
      using ⟨is-real A⟩
      by (simp add: power2-eq-square)
    hence  $\operatorname{Re}(A^2 + 2 * B * \operatorname{cnj} B - A * D) \geq 0$ 
      using ⟨ $\operatorname{Re} A * \operatorname{Re} D \leq \operatorname{Re}(B * \operatorname{cnj} B)$ ⟩
      using ⟨is-real A⟩ ⟨is-real D⟩
      by simp
    show ?thesis
      using divide-right-mono-neg[OF ⟨ $\operatorname{Re}(A^2 + 2 * B * \operatorname{cnj} B - A * D) \geq 0$ ⟩,
      of Re A] ⟨ $\operatorname{Re} A < 0$ ⟩ ⟨is-real A⟩ ⟨ $A \neq 0$ ⟩
      by (simp add: add-divide-distrib diff-divide-distrib del: complex-Re-mult)
      (subst Re-divide-real, auto simp add: power2-eq-square field-simps)

```

```

qed
ultimately
have  $dsc \leq Re(D - A - 2 * B * cnj B / A)$ 
  by simp
hence  $-Re(2 * (B * cnj B) / A) \geq Re((A - D + cor dsc))$ 
  by (simp add: field-simps)
hence  $(Re(2 * (B * cnj B)) / Re A) \geq Re(A - D + cor dsc)$ 
  using ⟨is-real A⟩ ⟨A ≠ 0⟩
  by (subst (asm) Re-divide-real, auto)
from divide-right-mono[OF this, of Re(2 * B * cnj B)]
have  $-1 / Re A \geq Re(A - D + cor dsc) / Re(2 * B * cnj B)$ 
  using ⟨Re A < 0⟩ ⟨B ≠ 0⟩ ⟨A ≠ 0⟩ ⟨0 < Re(2 * (B * cnj B))⟩
  by (simp add: field-simps del: complex-Re-mult)
from mult-right-mono-neg[OF this, of Re A]
show ?thesis
  using ⟨is-real A⟩ ⟨is-real D⟩ ⟨B ≠ 0⟩ ⟨Re A < 0⟩ ⟨A ≠ 0⟩
  apply (subst *)
  apply (subst Re-mult-real, simp add: div-reals)
  apply (subst Re-divide-real, simp, simp)
  apply (simp add: field-simps)
  done
qed
ultimately
show ?thesis
  by simp
next
  case False
  hence  $A = 0$ 
    using ⟨¬Re A > 0⟩ ⟨is-real A⟩
    by (cases A) simp
  thus ?thesis
    by simp
qed
qed
qed

definition chordal-circles-rep where
chordal-circles-rep H =
  (let (A, B, C, D) = Rep-circline-mat H;
   dsc = sqrt(Re((D-A)2 + 4 * (B*cnj B)));
   a1 = (A - D + cor dsc) / (2 * C);
   r1 = sqrt((4 - Re((-4 * a1/B) * A)) / (1 + Re(a1*cnj a1)));
   a2 = (A - D - cor dsc) / (2 * C);
   r2 = sqrt((4 - Re((-4 * a2/B) * A)) / (1 + Re(a2*cnj a2)));
   in ((a1, r1), (a2, r2)))
lift-definition chordal-circles :: ocircline ⇒ (complex × real) × (complex × real)
is chordal-circles-rep
proof –

```

```

fix H1 H2
obtain A1 B1 C1 D1 where hh1: (A1, B1, C1, D1) = Rep-circline-mat H1
  by (cases Rep-circline-mat H1) auto
obtain A2 B2 C2 D2 where hh2: (A2, B2, C2, D2) = Rep-circline-mat H2
  by (cases Rep-circline-mat H2) auto

assume ocircline-mat-eq H1 H2
then obtain k where *:  $k > 0$   $A2 = \operatorname{cor} k * A1 B2 = \operatorname{cor} k * B1 C2 = \operatorname{cor} k * C1 D2 = \operatorname{cor} k * D1$ 
  using hh1[symmetric] hh2[symmetric]
  by auto
let ?dsc1 =  $\sqrt{\operatorname{Re}((D1 - A1)^2 + 4 * (B1 * \operatorname{cnj} B1)))}$  and ?dsc2 =  $\sqrt{\operatorname{Re}((D2 - A2)^2 + 4 * (B2 * \operatorname{cnj} B2)))}$ 
let ?a11 =  $(A1 - D1 + \operatorname{cor} ?dsc1) / (2 * C1)$  and ?a12 =  $(A2 - D2 + \operatorname{cor} ?dsc2) / (2 * C2)$ 
let ?a21 =  $(A1 - D1 - \operatorname{cor} ?dsc1) / (2 * C1)$  and ?a22 =  $(A2 - D2 - \operatorname{cor} ?dsc2) / (2 * C2)$ 
let ?r11 =  $\sqrt{(4 - \operatorname{Re}((-4 * ?a11/B1) * A1)) / (1 + \operatorname{Re} (?a11 * \operatorname{cnj} ?a11)))}$ 
let ?r12 =  $\sqrt{(4 - \operatorname{Re}((-4 * ?a12/B2) * A2)) / (1 + \operatorname{Re} (?a12 * \operatorname{cnj} ?a12)))}$ 
let ?r21 =  $\sqrt{(4 - \operatorname{Re}((-4 * ?a21/B1) * A1)) / (1 + \operatorname{Re} (?a21 * \operatorname{cnj} ?a21)))}$ 
let ?r22 =  $\sqrt{(4 - \operatorname{Re}((-4 * ?a22/B2) * A2)) / (1 + \operatorname{Re} (?a22 * \operatorname{cnj} ?a22)))}$ 

have  $\operatorname{Re}((D2 - A2)^2 + 4 * (B2 * \operatorname{cnj} B2)) = k^2 * \operatorname{Re}((D1 - A1)^2 + 4 * (B1 * \operatorname{cnj} B1))$ 
  using *
  by (simp add: power2-eq-square field-simps)
hence ?dsc2 =  $k * ?dsc1$ 
  using  $\langle k > 0 \rangle$ 
  by (simp add: real-sqrt-mult)
hence  $A2 - D2 + \operatorname{cor} ?dsc2 = \operatorname{cor} k * (A1 - D1 + \operatorname{cor} ?dsc1)$   $A2 - D2 - \operatorname{cor} ?dsc2 = \operatorname{cor} k * (A1 - D1 - \operatorname{cor} ?dsc1)$   $2 * C2 = \operatorname{cor} k * (2 * C1)$ 
  using *
  by (auto simp add: field-simps)
hence ?a12 = ?a11 ?a22 = ?a21
  using  $\langle k > 0 \rangle$ 
  by simp-all
moreover
have  $\operatorname{Re}((-4 * ?a12/B2) * A2) = \operatorname{Re}((-4 * ?a11/B1) * A1)$ 
  using *
  by (subst (?a12 = ?a11)) (simp, simp add: field-simps)
have ?r12 = ?r11
  by (subst (?r12 = ?r11)) (simp, simp add: field-simps)
have ?r12 = ?r11
  by (subst (?r12 = ?r11)) (simp, simp add: field-simps)
moreover
have  $\operatorname{Re}((-4 * ?a22/B2) * A2) = \operatorname{Re}((-4 * ?a21/B1) * A1)$ 
  using *
  by (subst (?a22 = ?a21)) (simp, simp add: field-simps)
have ?r22 = ?r21
  by (subst (?r22 = ?r21)) (simp, simp add: field-simps)

```

```

(?a22 = ?a21) +) simp
moreover
have chordal-circles-rep H1 = ((?a11, ?r11), (?a21, ?r21))
  using hh1[symmetric] hh2[symmetric]
  unfolding chordal-circles-rep-def Let-def
  by simp
moreover
have chordal-circles-rep H1 = ((?a11, ?r11), (?a21, ?r21))
  using hh1[symmetric]
  unfolding chordal-circles-rep-def Let-def
  by simp
moreover
have chordal-circles-rep H2 = ((?a12, ?r12), (?a22, ?r22))
  using hh2[symmetric]
  unfolding chordal-circles-rep-def Let-def
  by simp
ultimately
show chordal-circles-rep H1 = chordal-circles-rep H2
  by metis
qed

lemma chordal-circle-det-positive:
fixes x y :: real
assumes x * y < 0
shows x / (x - y) > 0
proof (cases x > 0)
  case True
  hence y < 0
    using ‹x * y < 0›
    by (metis mult-less-cancel-left-pos mult-zero-right)
  have x - y > 0
    using ‹x > 0› ‹y < 0›
    by auto
  thus ?thesis
    using ‹x > 0›
    by (metis zero-less-divide-iff)
next
  case False
  hence y > 0 x < 0
    using ‹x * y < 0›
    by – (metis mult-less-cancel-left-disj mult-zero-right, metis less-linear mult-zero-left)
  have x - y < 0
    using ‹x < 0› ‹y > 0›
    by auto
  thus ?thesis
    using ‹x < 0›
    by (metis zero-less-divide-iff)
qed

```

lemma *chordal-circle1*:

assumes *is-real A is-real D Re (A * D) < 0 r = sqrt(Re ((4*A)/(A-D)))*

shows *mk-circeline A 0 0 D = chordal-circle ∞_h r*

using *assms*

proof *transfer*

fix *A D r*

assume *: *is-real A is-real D Re (A * D) < 0 r = sqrt (Re ((4*A)/(A-D)))*

hence *A ≠ 0 ∨ D ≠ 0*

by *auto*

hence *(A, 0, 0, D) ∈ {H. hermitean H ∧ H ≠ mat-zero}*

using *eq-cnj-iff-real[of A] eq-cnj-iff-real[of D]* *

unfolding *hermitean-def*

by *(simp add: mat-adj-def mat-cnj-def)*

moreover

have *(- (cor r)², 0, 0, 4 - (cor r)²) ∈ {H. hermitean H ∧ H ≠ mat-zero}*

by *(simp add: hermitean-def mat-adj-def mat-cnj-def complex-cnj power2-eq-square)*

moreover

have *A ≠ D*

using *⟨Re (A * D) < 0⟩ ⟨is-real A⟩ ⟨is-real D⟩*

by *auto*

have *Re ((4*A)/(A-D)) ≥ 0*

proof-

have *Re A / Re (A - D) ≥ 0*

using *⟨Re (A * D) < 0⟩ ⟨is-real A⟩ ⟨is-real D⟩*

using *chordal-circle-det-positive[of Re A Re D]*

by *simp*

thus *?thesis*

using *⟨is-real A⟩ ⟨is-real D⟩ ⟨A ≠ D⟩*

by *(subst Re-divide-real, auto) (metis mult-nonneg-nonpos zero-le-divide-iff zero-le-mult-iff zero-le-numeral)*

qed

moreover

have *- (cor (sqrt (Re (4 * A / (A - D)))))² = cor (Re (4 / (D - A))) * A*

using *⟨Re ((4*A)/(A-D)) ≥ 0⟩ ⟨is-real A⟩ ⟨is-real D⟩ ⟨A ≠ D⟩*

by *(subst cor-squared, subst real-sqrt-power[symmetric], simp) (simp add: Re-divide-real Re-multiply-real complex-of-real-Re of-real-numeral minus-divide-right)*

moreover

have *4 * (A - D) - 4 * A = 4 * -D*

by *(simp add: field-simps)*

hence *4 - 4 * A / (A - D) = -4 * D / (A - D)*

using *⟨A ≠ D⟩*

by *(smt ab-semigroup-mult-class.mult-ac(1) diff-divide-eq-iff eq-iff-diff-eq-0 mult-minus1 mult-minus1-right mult-numeral-1-right neg-numeral-def right-diff-distrib-numeral times-divide-eq-right)*

hence *4 - 4 * A / (A - D) = 4 * D / (D - A)*

by *(metis (hide-lams, no-types) minus-diff-eq minus-divide-left minus-divide-right minus-mult-left neg-numeral-def)*

hence *4 - (cor (sqrt (Re (4 * A / (A - D)))))² = cor (Re (4 / (D - A))) * D*

```

using ⟨Re ((4*A)/(A-D)) ≥ 0⟩ ⟨is-real A⟩ ⟨is-real D⟩ ⟨A ≠ D⟩
by (subst cor-squared, subst real-sqrt-power[symmetric], simp) (simp add:
  Re-divide-real Re-multiply-real complex-of-real-Re of-real-numeral)
ultimately
show circline-mat-eq (mk-circline-rep A 0 0 D) (chordal-circle-rep inf-homo-rep
r)
using ⟨is-real A⟩ ⟨is-real D⟩ ⟨A ≠ D⟩ ⟨r = sqrt(Re ((4*A)/(A-D)))⟩
by (simp add: chordal-circle-rep-def mk-circline-rep-def Abs-circline-mat-inverse)
(rule-tac x=Re(4/(D-A)) in exI, auto)
qed

lemma chordal-circle2:
assumes is-real A is-real D Re (A * D) < 0 r = sqrt(Re ((4*D)/(D-A)))
shows mk-circline A 0 0 D = chordal-circle 0_h r
using assms
proof transfer
fix A D r
assume *: is-real A is-real D Re (A * D) < 0 r = sqrt (Re ((4*D)/(D-A)))
hence A ≠ 0 ∨ D ≠ 0
by auto
hence (A, 0, 0, D) ∈ {H. hermitean H ∧ H ≠ mat-zero}
using eq-cnj-iff-real[of A] eq-cnj-iff-real[of D] *
unfolding hermitean-def
by (simp add: mat-adj-def mat-cnj-def)
moreover
have (4 - (cor r)2, 0, 0, - (cor r)2) ∈ {H. hermitean H ∧ H ≠ mat-zero}
by (simp add: hermitean-def mat-adj-def mat-cnj-def complex-cnj power2-eq-square)
(metis mult-zero-right of-real-0 zero-neq-numeral)
moreover
have A ≠ D
using ⟨Re (A * D) < 0⟩ ⟨is-real A⟩ ⟨is-real D⟩
by auto
have Re((4*D)/(D-A)) ≥ 0
proof-
have Re D / Re (D - A) ≥ 0
using ⟨Re (A * D) < 0⟩ ⟨is-real A⟩ ⟨is-real D⟩
using chordal-circle-det-positive[of Re D Re A]
by (simp add: field-simps)
thus ?thesis
using ⟨is-real A⟩ ⟨is-real D⟩ ⟨A ≠ D⟩
by (subst Re-divide-real, auto) (metis mult-nonneg-nonpos zero-le-divide-iff
zero-le-mult-iff zero-le-numeral)
qed
have 4 * (D - A) - 4 * D = 4 * -A
by (simp add: field-simps)
hence 4 - 4 * D / (D - A) = -4 * A / (D - A)
using ⟨A ≠ D⟩
by (smt ab-semigroup-mult-class.mult-ac(1) diff-divide-eq-iff eq-iff-diff-eq-0 mult-minus1
mult-minus1-right mult-numeral-1-right neg-numeral-def right-diff-distrib-numeral

```

times-divide-eq-right)
hence $4 - 4 * D / (D - A) = 4 * A / (A - D)$
by (metis (hide-lams, no-types) minus-diff-eq minus-divide-left minus-divide-right
minus-mult-left neg-numeral-def)
hence $4 - (\text{cor}(\sqrt{\text{Re}((4*D)/(D-A))}))^2 = \text{cor}(\text{Re}(4 / (A - D))) * A$
using ⟨is-real A⟩ ⟨is-real D⟩ ⟨A ≠ D⟩ ⟨Re(4 * D / (D - A)) ≥ 0⟩
by (subst cor-squared, subst real-sqrt-power[symmetric], simp) (simp add:
Re-divide-real complex-of-real-Re of-real-numeral)

moreover
have $- (\text{cor}(\sqrt{\text{Re}((4*D)/(D-A))}))^2 = \text{cor}(\text{Re}(4 / (A - D))) * D$
using ⟨is-real A⟩ ⟨is-real D⟩ ⟨A ≠ D⟩ ⟨Re((4*D)/(D-A)) ≥ 0⟩
by (subst cor-squared, subst real-sqrt-power[symmetric], simp) (simp add:
Re-divide-real complex-of-real-Re of-real-numeral minus-divide-right)

ultimately
show circline-mat-eq (mk-circline-rep A 0 0 D) (chordal-circle-rep zero-homo-rep
r)
using ⟨is-real A⟩ ⟨is-real D⟩ ⟨A ≠ 0 ∨ D ≠ 0⟩ ⟨r = sqrt(Re((4*D)/(D-A)))⟩
by (simp add: chordal-circle-rep-def mk-circline-rep-def Abs-circline-mat-inverse)
(rule-tac x=Re(4/(A-D)) in exI, auto)
qed

lemma chordal-circle':
assumes $B \neq 0$ $(A, B, C, D) \in \{H. \text{hermitean } H \wedge H \neq \text{mat-zero}\}$ $\text{Re}(\text{mat-det}(A, B, C, D)) \leq 0$
 $C * a^2 + (D - A) * a - B = 0$ $r = \sqrt{((4 - \text{Re}(-4 * a/B)) * A) / (1 + \text{Re}(a * \text{cnj } a))}$
shows mk-circline A B C D = chordal-circle (of-complex a) r
using assms
proof transfer
fix A B C D a :: complex **and** r :: real

let ?k = $-4 * a / B$

assume *: $(A, B, C, D) \in \{H. \text{hermitean } H \wedge H \neq \text{mat-zero}\}$ **and** **: $B \neq 0$
 $C * a^2 + (D - A) * a - B = 0$ **and** rr: $r = \sqrt{((4 - \text{Re}(\text{?k} * A)) / (1 + \text{Re}(a * \text{cnj } a)))}$ **and** det: $\text{Re}(\text{mat-det}(A, B, C, D)) \leq 0$

have is-real A is-real D C = cnj B
using * hermitean-elems
by auto

from ** **have** a12: let dsc = $\sqrt{\text{Re}((D-A)^2 + 4 * (B * \text{cnj } B))}$
in a = $(A - D + \text{cor } dsc) / (2 * C) \vee a = (A - D - \text{cor } dsc) / (2 * C)$
proof-
have $\text{Re}((D-A)^2 + 4 * (B * \text{cnj } B)) \geq 0$
using ⟨is-real A⟩ ⟨is-real D⟩

```

    by (subst complex-mult-cnj-cmod) (simp add: power2-eq-square)
    hence csqrt ((D - A)2 - 4 * C * -B) = cor (sqrt (Re ((D - A)2 + 4 * (B
* cnj B))))
      using csqrt-real[of ((D - A)2 + 4 * (B * cnj B))] ⟨is-real A⟩ ⟨is-real D⟩ ⟨C
= cnj B⟩
        by (auto simp add: power2-eq-square field-simps)
        thus ?thesis
          using complex-quadratic-equation-full[of C a D - A -B] ⟨C * a2 + (D - A)
* a - B = 0⟩ ⟨B ≠ 0⟩ ⟨C = cnj B⟩
            by (simp add: Let-def)
qed

have is-real ?k
  using a12 ⟨C = cnj B⟩ ⟨is-real A⟩ ⟨is-real D⟩
  by (auto simp add: Let-def div-reals)
have a ≠ 0
  using **
  by auto
hence Re ?k ≠ 0
  using ⟨is-real (-4*a / B)⟩ ⟨B ≠ 0⟩
  by (metis complex-surj complex-zero-def mult-eq-0-iff nonzero-divide-eq-eq zero-neq-neg-numeral)

moreover
have -4 * a = cor (Re ?k) * B
  using complex-of-real-Re[OF ⟨is-real (-4*a / B)⟩] ⟨B ≠ 0⟩
  by simp
moreover
have is-real (a/B)
  using ⟨is-real ?k⟩
  by (metis Im-mult-real complex-Im-neg-numeral complex-Re-neg-numeral mult-eq-0-iff
times-divide-eq-right zero-neq-neg-numeral)
hence is-real (B * cnj a)
  by (smt mult.commute complex-In-mult-cnj-zero complex-cnj-divide complex-cnj-zero-iff
eq-cnj-iff-real eq-divide-eq times-divide-eq-right)
hence B * cnj a = cnj B * a
  using eq-cnj-iff-real[of B * cnj a]
  by (simp add: complex-cnj)
hence -4 * cnj a = cor (Re ?k) * C
  using ⟨C = cnj B⟩
  using complex-of-real-Re[OF ⟨is-real ?k⟩] ⟨B ≠ 0⟩
  by (simp, simp add: field-simps)
moreover
have 1 + a * cnj a ≠ 0
  by (subst complex-mult-cnj-cmod) (smt cor-add of-real-0 of-real-1 of-real-eq-iff
realpow-square-minus-le)
have r2 = (4 - Re (?k * A)) / (1 + Re (a * cnj a))
proof-
  have Re (a / B * A) ≥ -1
    using a12 chordal-circle-radius-positive[of A B C D] * ⟨B ≠ 0⟩ det

```

```

by (auto simp add: Let-def field-simps)
from mult-right-mono-neg[OF this, of -4]
have 4 - Re (?k * A) ≥ 0
  using Re-mult-real[of -4 a / B * A]
  by (simp add: field-simps)
moreover
have 1 + Re (a * cnj a) > 0
  by (subst complex-mult-cnj-cmod) (smt Re-complex-of-real ‹a ≠ 0› norm-eq-zero
zero-less-power2)
ultimately
have (4 - Re (?k * A)) / (1 + Re (a * cnj a)) ≥ 0
  by (metis divide-nonneg-pos)
thus ?thesis
  using rr
  by simp
qed
hence r2 = Re ((4 - ?k * A) / (1 + a * cnj a))
  using ‹is-real ?k› ‹is-real A› ‹1 + a * cnj a ≠ 0›
  by (subst Re-divide-real, auto)
hence (cor r)2 = (4 - ?k * A) / (1 + a * cnj a)
  using ‹is-real ?k› ‹is-real A›
  using mult-reals[of ?k A]
  by (simp add: cor-squared) (subst complex-of-real-Re, subst div-reals, auto)
hence 4 - (cor r)2 * (a * cnj a + 1) = cor (Re ?k) * A
  using complex-of-real-Re[OF ‹is-real (-4*a/B)›]
  using ‹1 + a * cnj a ≠ 0›
  by (simp add: field-simps)
moreover
have ?k = cnj ?k
  using ‹is-real ?k›
  by (subst eq-cnj-iff-real) simp
have ?k2 = cor ((cmod ?k)2)
  using cor-cmod-real[OF ‹is-real ?k›]
  unfolding power2-eq-square
  by (subst cor-mult) (metis minus-mult-minus)
hence ?k2 = ?k * cnj ?k
  using complex-mult-cnj-cmod[of ?k]
  by simp
hence ***: a * cnj a = (cor ((Re ?k)2) * B * C) / 16
  using complex-of-real-Re[OF ‹is-real (-4*a/B)›] ‹C = cnj B› ‹is-real (-4*a/B)›
B ≠ 0
  by (simp add: complex-cnj)
from ** have cor ((Re ?k)2) * B * C - 4 * cor (Re ?k) * (D-A) - 16 = 0
  using complex-of-real-Re[OF ‹is-real ?k›]
  by (simp add: power2-eq-square, simp add: field-simps, algebra)
hence ?k * (D-A) = 4 * (cor ((Re ?k)2) * B * C / 16 - 1)
  by (subst (asm) complex-of-real-Re[OF ‹is-real ?k›]) algebra
hence ?k * (D-A) = 4 * (a * cnj a - 1)

```

```

by (subst (asm) ***[symmetric]) simp

hence  $4 * a * \text{cnj } a - (\text{cor } r)^2 * (a * \text{cnj } a + 1) = \text{cor } (\text{Re } ?k) * D$ 
  using  $\langle 4 - (\text{cor } r)^2 * (a * \text{cnj } a + 1) = \text{cor } (\text{Re } ?k) * A \rangle$ 
  using complex-of-real-Re[OF is-real  $(-4*a/B)$ ]
  by simp algebra
ultimately
show circline-mat-eq (mk-circline-rep A B C D) (chordal-circle-rep (of-complex-coords a) r)
  using *  $\langle a \neq 0 \rangle$ 
  by (simp add: mk-circline-rep-def Abs-circline-mat-inverse) (rule-tac x=Re  $(-4*a / B)$  in exI, simp)
qed

lift-definition o-circline-point-0h :: ocircline is circline-point-0h-rep
done

lemma of-ocircline-o-circline-point-0h [simp]: of-ocircline o-circline-point-0h = circline-point-0h
  by (metis circline-point-0h-def o-circline-point-0h-def of-ocircline.abs-eq)

lemma ocircline-set-0h:
assumes ocircline-set  $H = \{0_h\}$ 
shows  $H = o\text{-circline-point-0h} \vee H = \text{opposite-ocircline } (o\text{-circline-point-0h})$ 
proof-
have of-ocircline  $H = \text{circline-point-0h}$ 
  using assms
  using circline-set-ocircline-set[of  $H$ , symmetric]
  using unique-circline-type-zero-0h' card-eq1-circline-type-zero[of of-ocircline  $H$ ]
  by blast
thus ?thesis
  by (metis inj-of-ocircline of-ocircline-o-circline-point-0h)
qed

```

11.13 Disc automorphisms

```

lemma circline-set-fix-iff-circline-fix:
assumes circline-set  $H' \neq \{\}$ 
shows (moebius-pt M) '(circline-set H) = circline-set  $H' \longleftrightarrow$  moebius-circline M H =  $H'$ 
using assms
by (subst moebius-circline-set, auto) (rule inj-circline-set[of - H'], auto)

lemma ocircline-set-fix-iff-ocircline-fix:
assumes ocircline-set  $H' \neq \{\}$ 
shows (moebius-pt M) '(ocircline-set H) = ocircline-set  $H' \longleftrightarrow$ 
moebius-ocircline M H =  $H' \vee$  moebius-ocircline M H = opposite-ocircline H'
using assms inj-ocircline-set[of - H']
by (subst moebius-ocircline-set, auto)

```

```

definition Unitary11-gen-rep where
  Unitary11-gen-rep M  $\longleftrightarrow$  unitary11-gen (Rep-moebius-mat M)

lift-definition Unitary11-gen :: moebius  $\Rightarrow$  bool is Unitary11-gen-rep
apply (auto simp add: Unitary11-gen-rep-def)
apply (simp add: unitary11-gen-mult-sm)
apply (simp add: unitary11-gen-div-sm)
done

lemma unit-circle-fix-iff-Unitary11-gen:
  shows moebius-circline M unit-circle = unit-circle  $\longleftrightarrow$  Unitary11-gen M (is ?lhs
= ?rhs)
proof
  assume ?lhs
  thus ?rhs
  proof (transfer)
    fix M
    assume circline-mat-eq (moebius-circline-rep M unit-circle-rep) unit-circle-rep
    then obtain k where k  $\neq$  0 (1, 0, 0, -1) = cor k *sm congruence (mat-inv
(Rep-moebius-mat M)) (1, 0, 0, -1)
      by auto
    hence (1/cor k, 0, 0, -1/cor k) = congruence (mat-inv (Rep-moebius-mat
M)) (1, 0, 0, -1)
      using mult-sm-inv-l[of cor k congruence (mat-inv (Rep-moebius-mat M)) (1,
0, 0, -1)]
      by simp
    hence congruence (Rep-moebius-mat M) (1/cor k, 0, 0, -1/cor k) = (1, 0,
0, -1)
      using Rep-moebius-mat[of M] mat-det-inv[of Rep-moebius-mat M]
      using congruence-inv[of mat-inv (Rep-moebius-mat M) (1, 0, 0, -1) (1/cor
k, 0, 0, -1/cor k)]
      by simp
    hence congruence (Rep-moebius-mat M) (1, 0, 0, -1) = cor k *sm (1, 0, 0,
-1)
      using congruence-scale-m[of Rep-moebius-mat M 1/cor k (1, 0, 0, -1)]
      using mult-sm-inv-l[of 1/cor k congruence (Rep-moebius-mat M) (1, 0, 0,
-1) (1, 0, 0, -1)] (k  $\neq$  0)
      by simp
    thus Unitary11-gen-rep M
      using <k  $\neq$  0>
      unfolding Unitary11-gen-rep-def unitary11-gen-def
      by simp
  qed
next
  assume ?rhs
  thus ?lhs
  proof (transfer)

```

```

fix M
assume Unitary11-gen-rep M
hence unitary11-gen (mat-inv (Rep-moebius-mat M))
  using Rep-moebius-mat[of M]
  using unitary11-gen-mat-inv
  by (simp add: Unitary11-gen-rep-def)
thus circline-mat-eq (moebius-circline-rep M unit-circle-rep) unit-circle-rep
  unfolding unitary11-gen-real
  by auto (rule-tac x=1/k in exI, simp)
qed
qed

lemma unit-circle-set-fix-iff-Unitary11-gen:
  shows (moebius-pt M ` (circline-set unit-circle) = (circline-set unit-circle))  $\longleftrightarrow$ 
    Unitary11-gen M (is ?lhs  $\longleftrightarrow$  ?rhs)
  using unit-circle-fix-iff-Unitary11-gen[of M] circline-set-fix-iff-circline-fix[of unit-circle
    M unit-circle]
  using one-on-unit-circle
  by auto

definition Unitary11-gen-direct-rep where
  Unitary11-gen-direct-rep M  $\longleftrightarrow$ 
    (let (A, B, C, D) = Rep-moebius-mat M
      in unitary11-gen (A, B, C, D)  $\wedge$  (B = 0  $\vee$  Re ((A*D)/(B*C)) > 1))

lift-definition Unitary11-gen-direct :: moebius  $\Rightarrow$  bool is Unitary11-gen-direct-rep
proof-
  fix M M'
  let ?M = Rep-moebius-mat M and ?M' = Rep-moebius-mat M'
  assume moebius-mat-eq M M'
  then obtain k where *: k  $\neq$  0 Rep-moebius-mat M' = k *sm Rep-moebius-mat
    M
  by auto
  hence **: unitary11-gen (Rep-moebius-mat M)  $\longleftrightarrow$  unitary11-gen (Rep-moebius-mat
    M')
  using unitary11-gen-mult-sm[of k ?M] unitary11-gen-div-sm[of k ?M]
  by auto
  obtain A B C D where MM: (A, B, C, D) = Rep-moebius-mat M
    by (cases Rep-moebius-mat M) auto
  obtain A' B' C' D' where MM': (A', B', C', D') = Rep-moebius-mat M'
    by (cases Rep-moebius-mat M') auto

  show Unitary11-gen-direct-rep M = Unitary11-gen-direct-rep M'
    using * ** MM MM'
    unfolding Unitary11-gen-direct-rep-def Let-def
    by auto
qed

lemma ounit-circle-fix-iff-Unitary11-gen-direct:

```

```

shows moebius-ocircline M ounit-circle = ounit-circle  $\longleftrightarrow$  Unitary11-gen-direct
M (is ?lhs  $\longleftrightarrow$  ?rhs)
proof
assume *: ?lhs
have moebius-circline M unit-circle = unit-circle
apply (subst moebius-circline-ocircline[of M unit-circle])
apply (subst of-circline-unit-circline)
apply (subst *)
by simp

hence Unitary11-gen M
by (simp add: unit-circle-fix-iff-Unitary11-gen)
thus ?rhs
using *
proof (transfer)
fix M
let ?M = Rep-moebius-mat M
let ?H = (1, 0, 0, -1)
obtain A B C D where MM: (A, B, C, D) = ?M
by (cases ?M) auto
assume Unitary11-gen-rep M ocircline-mat-eq (moebius-circline-rep M unit-circle-rep)
unit-circle-rep
then obtain k where 0 < k ?H = cor k *sm congruence (mat-inv ?M) ?H
by auto
hence congruence ?M ?H = cor k *sm ?H
using congruence-inv[of mat-inv ?M ?H (1/cor k) *sm ?H] Rep-moebius-mat[of
M]
using mult-sm-inv-l[of cor k congruence (mat-inv ?M) ?H ?H]
using mult-sm-inv-l[of 1/cor k congruence ?M ?H]
using congruence-scale-m[of ?M 1/cor k ?H]
by (auto simp add: mat-det-inv)
then obtain a b k' where k' ≠ 0 ?M = k' *sm (a, b, cnj b, cnj a) sgn (Re
(mat-det (a, b, cnj b, cnj a))) = 1
using unitary11-sgn-det-orientation'[of ?M k] ⟨k > 0⟩
by auto
moreover
have mat-det (a, b, cnj b, cnj a) ≠ 0
using ⟨sgn (Re (mat-det (a, b, cnj b, cnj a))) = 1⟩
by (metis complex-Re-zero sgn-zero zero-neq-one)
ultimately
show Unitary11-gen-direct-rep M
using unitary11-sgn-det[of k' a b ?M A B C D]
using MM[symmetric] ⟨k > 0⟩ ⟨Unitary11-gen-rep M⟩
by (simp add: Unitary11-gen-rep-def Unitary11-gen-direct-rep-def sgn-1-pos
split: split-if-asm)
qed
next
assume ?rhs
thus ?lhs

```

```

proof (transfer)
fix M
let ?M = Rep-moebius-mat M
obtain A B C D where MM: (A, B, C, D) = ?M
  by (cases ?M) auto
assume Unitary11-gen-direct-rep M
hence unitary11-gen ?M B = 0 ∨ 1 < Re (A * D / (B * C))
  using MM[symmetric]
  by (auto simp add: Unitary11-gen-direct-rep-def)
have sgn (if B = 0 then 1 else sgn (Re (A * D / (B * C)) - 1)) = 1
  using ‹B = 0 ∨ 1 < Re (A * D / (B * C))›
  by auto
then obtain k' where k' > 0 congruence (Rep-moebius-mat M) (1, 0, 0, -1)
= cor k' *sm (1, 0, 0, -1)
  using unitary11-orientation[OF ‹unitary11-gen ?M› MM[symmetric]]
  by (auto simp add: sgn-1-pos)
thus ocircline-mat-eq (moebius-circline-rep M unit-circle-rep) unit-circle-rep
  using congruence-inv[of ?M (1, 0, 0, -1) cor k' *sm (1, 0, 0, -1)]
Rep-moebius-mat[of M]
  using congruence-scale-m[of mat-inv ?M cor k' (1, 0, 0, -1)]
  by auto
qed
qed

```

Blaschke factor

```

definition blaschke-rep where
blaschke-rep a = Abs-moebius-mat (1, -a, -cnj a, 1)

```

```

lemma blaschke-rep-Rep1:
assumes cmod a ≠ 1
shows Rep-moebius-mat (blaschke-rep a) = (1, -a, -cnj a, 1)
using assms
by (simp add: blaschke-rep-def Abs-moebius-mat-inverse)

```

```

lemma blaschke-rep-Rep2:
assumes a * cnj a ≠ 1
shows Rep-moebius-mat (blaschke-rep a) = (1, -a, -cnj a, 1)
using assms
by (simp add: blaschke-rep-def Abs-moebius-mat-inverse)

```

```

lift-definition blaschke :: complex ⇒ moebius is blaschke-rep
by (simp del: moebius-mat-eq-def)

```

```

lemma blaschke-a-to-zero:
assumes cmod a ≠ 1
shows moebius-pt (blaschke a) (of-complex a) = 0h
proof-
from assms have a * cnj a ≠ 1
  by simp

```

```

thus ?thesis
  by (transfer) (simp add: blaschke-rep-Rep2, rule-tac x=1/(1 - a*cnj a) in
exI, simp add: field-simps)
qed

lemma blaschke-inv-a-inf:
  assumes cmod a ≠ 1
  shows moebius-pt (blaschke a) (inversion-homo (of-complex a)) = ∞h
proof-
  from assms have a * cnj a ≠ 1
    by simp
  thus ?thesis
    unfolding inversion-homo-def
    by (transfer) (simp add: blaschke-rep-Rep2 vec-cnj-def, rule-tac x=1/(1 -
a*cnj a) in exI, simp add: field-simps)
qed

lemma blaschke-Unitary11-gen-rep:
  assumes a * cnj a ≠ 1
  shows Unitary11-gen-rep (blaschke-rep a)
proof-
  have is-real (1 - a*cnj a)
    by auto
  moreover
  hence cor (Re (1 - a*cnj a)) = 1 - a*cnj a
    by (rule complex-of-real-Re)
  moreover
  have Re (a*cnj a) ≠ 1
    using ⟨is-real (1 - a*cnj a)⟩ assms
    by (metis complex-In-mult-cnj-zero complex-of-real-Re of-real-1)
  ultimately
  show ?thesis
    using assms
    using blaschke-rep-Rep2
    by (auto simp add: blaschke-rep-def Unitary11-gen-rep-def unitary11-gen-real
mat-adj-def mat-cnj-def complex-cnj field-simps simp del: complex-Re-mult) (rule-tac
x=Re (1 - a*cnj a) in exI, simp del: complex-Re-mult)
qed

lemma blaschke-unitary11-gen-direct-rep:
  assumes Re (a * cnj a) < 1
  shows Unitary11-gen-direct-rep (blaschke-rep a)
proof-
  have a * cnj a ≠ 1
    using assms
    by (cases a, simp)
  show ?thesis
  proof (cases a = 0)
    case True

```

```

thus ?thesis
  using blaschke-Unitary11-gen-rep[of a]
  by (simp add: Unitary11-gen-direct-rep-def Unitary11-gen-rep-def blaschke-rep-def)
next
  case False
  hence Re (a * cnj a) > 0
    by (subst complex-mult-cnj-cmod) (metis Re-complex-of-real zero-less-norm-iff
zero-less-power)
  thus ?thesis
    using assms ‹a * cnj a ≠ 1› ‹a ≠ 0›
    using blaschke-Unitary11-gen-rep[of a] blaschke-rep-Rep2[of a] Re-divide-real[of
a*cnj a 1]
    by (auto simp add: Unitary11-gen-direct-rep-def blaschke-rep-def Unitary11-gen-rep-def
simp del: complex-Re-mult)
  qed
qed

lemma blaschke-Unitary11-gen:
  assumes a * cnj a ≠ 1
  shows Unitary11-gen (blaschke a)
using assms
by (transfer) (rule blaschke-Unitary11-gen-rep)

lemma blaschke-Unitary11-gen-direct:
  assumes Re (a * cnj a) < 1
  shows Unitary11-gen-direct (blaschke a)
using assms
by transfer (simp add: blaschke-unitary11-gen-direct-rep)

lemma blaschke-unit-circle-fix:
  assumes cmod a ≠ 1
  shows moebius-circeline (blaschke a) unit-circle = unit-circle
using assms
using blaschke-Unitary11-gen unit-circle-fix-iff-Unitary11-gen
by simp

lemma blaschke-ounit-circle-fix:
  assumes cmod a < 1
  shows moebius-ocircline (blaschke a) ounit-circle = ounit-circle
proof-
  have Re (a * cnj a) < 1
    using assms
    by (metis complex-mod-sqrt-Re-mult-cnj real-sqrt-lt-1-iff)
  thus ?thesis
    using assms
    using blaschke-Unitary11-gen-direct ounit-circle-fix-iff-Unitary11-gen-direct
    by simp
qed

```

```

lemma [simp]: hermitean (1, 0, 0, -1)
by (auto simp add: hermitean-def mat-adj-def mat-cnj-def)

definition is-disc-aut where is-disc-aut f  $\longleftrightarrow$  bij-betw f unit-disc unit-disc

lemma is-disc-aut-iff-unit-disc-fix:
  shows is-disc-aut (moebius-pt M)  $\longleftrightarrow$  (moebius-pt M) ‘ unit-disc = unit-disc
  using bij-moebius-pt[of M]
  unfolding is-disc-aut-def is-moebius-def
  unfolding bij-betw-def
  by auto (metis injD inj-onI)

lemma comp-inv-l:
  assumes f o inv g = h bij g
  shows f = h o g
  using assms
  by (metis bij-def o-inv-o-cancel)

lemma in-unit-disc-cmod-lt-1:
  assumes of-complex a ∈ unit-disc
  shows cmod a < 1
  using assms
  unfolding unit-disc-def disc-def
  apply auto
  proof (transfer)
    fix a
    assume in-ocircline-rep unit-circle-rep (of-complex-coords a)
    hence Re a * Re a + Im a * Im a < 1
      by (simp add: in-ocircline-rep-def Let-def vec-cnj-def)
    hence (cmod a)2 < 1
      unfolding cmod-def
      by (simp, simp add: power2-eq-square)
    thus cmod a < 1
      by (metis less-1-mult not-less-iff-gr-or-eq one-power2 power2-eq-square)
  qed

```

11.14 Angle between circlines

```

fun mat-det-12 :: complex-mat  $\Rightarrow$  complex-mat  $\Rightarrow$  complex where
  mat-det-12 (A1, B1, C1, D1) (A2, B2, C2, D2) = A1*D2 + A2*D1 - B1*C2
  - B2*C1

lemma mat-det-12-mm-l [simp]: mat-det-12 (M *mm A) (M *mm B) = mat-det
  M * mat-det-12 A B
  by (cases M, cases A, cases B) (simp add: field-simps)

lemma mat-det-12-mm-r [simp]: mat-det-12 (A *mm M) (B *mm M) = mat-det
  M * mat-det-12 A B

```

```

by (cases M, cases A, cases B) (simp add: field-simps)

lemma mat-det-12-sm-l [simp]: mat-det-12 (k *sm A) B = k * mat-det-12 A B
by (cases A, cases B) (simp add: field-simps)

lemma mat-det-12-sm-r [simp]: mat-det-12 A (k *sm B) = k * mat-det-12 A B
by (cases A, cases B) (simp add: field-simps)

lemma mat-det-12-congruence [simp]:
  mat-det-12 (congruence M A) (congruence M B) = (cor ((cmod (mat-det M))2))
  * mat-det-12 A B
by ((subst mult-mm-assoc[symmetric])+, subst mat-det-12-mm-l, subst mat-det-12-mm-r,
  subst mat-det-adj) (auto simp add: field-simps complex-mult-cnj-cmod)

lemma mat-det-congruence [simp]:
  mat-det (congruence M A) = (cor ((cmod (mat-det M))2)) * mat-det A
by (simp add: mat-det-adj complex-mult-cnj-cmod field-simps)

definition cos-angle-rep where
cos-angle-rep H1 H2 =
(let H1 = Rep-circline-mat H1;
 H2 = Rep-circline-mat H2 in
 - Re (mat-det-12 H1 H2) / (2 * (sqrt (Re (mat-det H1 * mat-det H2)))))

lemma [simp]: is-real (mat-det (Rep-circline-mat H))
  using Rep-circline-mat[of H]
  by (simp add: mat-det-hermitean-real)

lift-definition cos-angle :: ocircline ⇒ ocircline ⇒ real is cos-angle-rep
by (auto simp add: cos-angle-rep-def Let-def real-sqrt-mult)

lemma ang-vec-opposite-opposite':
  assumes a1 ≠ E a2 ≠ E
  shows (E - a1) (E - a2) = (a1 - E) (a2 - E)
  using ang-vec-opposite-opposite[of E - a1 E - a2] assms
  by (simp add: field-simps del: ang-vec-def)

lemma cos-ang-circ-simp:
  assumes E ≠ μ1 E ≠ μ2
  shows cos (ang-circ E μ1 μ2 p1 p2) = sgn-bool (p1 = p2) * cos (arg (E - μ2)
  - arg (E - μ1))
  using assms
  using cos-periodic-pi2[of arg (E - μ2) - arg (E - μ1)]
  using cos-periodic-pi3[of arg (E - μ2) - arg (E - μ1)]
  using ang-circ-simp[OF assms, of p1 p2]
  by auto (auto simp add: field-simps)

lemma Re-sgn:
  assumes is-real A A ≠ 0

```

```

shows  $\text{Re}(\text{sgn } A) = \text{sgn-bool}(\text{Re } A > 0)$ 
using assms
by (cases A) simp

lemma Re-mult-real3:
assumes is-real z1 is-real z2 is-real z3
shows  $\text{Re}(z1 * z2 * z3) = \text{Re } z1 * \text{Re } z2 * \text{Re } z3$ 
using assms
by (metis Re-mult-real mult-reals)

lemma [simp]:  $\text{sgn}(\sqrt{x}) = \text{sgn } x$ 
by (smt real-sqrt-eq-zero-cancel-iff real-sqrt-lt-0-iff sgn-real-def)

lemma sgn-divide:
fixes x y :: real
shows  $\text{sgn}(x / y) = \text{sgn } x / \text{sgn } y$ 
by (metis divide-inverse inverse-sgn real-scaleR-def sgn-scaleR)

lemma real-circle-sgn-r:
assumes is-circle H (a, r) = euclidean-circle H
shows  $\text{sgn } r = -\text{circline-type } H$ 
using assms
proof transfer
fix H a r
obtain A B C D where HH: Rep-circline-mat H = (A, B, C, D)
  by (cases Rep-circline-mat H) auto
hence is-real A is-real D
  using hermitean-elems Rep-circline-mat[of H]
  by auto
assume  $\neg \text{circline-A0rep } H (a, r) = \text{euclidean-circle-rep } H$ 
hence A ≠ 0
  using (¬ circline-A0rep H) HH
  by (simp add: circline-A0rep-def)
hence  $\text{Re } A * \text{Re } A > 0$ 
  using (is-real A)
  by (metis complex-Im-zero complex-Re-zero complex-equality not-real-square-gt-zero)

thus  $\text{sgn } r = -\text{circline-type-rep } H$ 
  using HH (a, r) = euclidean-circle-rep H (is-real A) (is-real D) (A ≠ 0)
  by (simp add: euclidean-circle-rep-def circline-type-rep-def Re-divide-real sgn-minus[symmetric]
    sgn-divide)
qed

lemma
assumes
  is-circle (of-ocircline H1) is-circle (of-ocircline H2)
  circline-type (of-ocircline H1) < 0 circline-type (of-ocircline H2) < 0
  (a1, r1) = euclidean-circle (of-ocircline H1) (a2, r2) = euclidean-circle (of-ocircline H2)

```

$of\text{-complex } E \in ocircleline\text{-set } H1 \cap ocircleline\text{-set } H2$
shows $\cos\text{-angle } H1 H2 = \cos(\text{ang-circ } E a1 a2)$ ($pos\text{-oriented } H1$) ($pos\text{-oriented } H2$)
proof–
let $?p1 = pos\text{-oriented } H1$ **and** $?p2 = pos\text{-oriented } H2$
have $E \in circle a1 r1 E \in circle a2 r2$
using $classic\text{-circle}[of of\text{-ocircle } H1 a1 r1]$ $classic\text{-circle}[of of\text{-ocircle } H2 a2 r2]$
using $assms$ *of-complex-inj*
by *auto*
hence $*: cdist E a1 = r1 cdist E a2 = r2$
unfolding *circle-def*
by (*simp-all add: norm-minus-commute*)
have $r1 > 0 r2 > 0$
using $assms(1\text{--}6)$ $real\text{-circle-sgn-r}[of of\text{-ocircle } H1 a1 r1]$ $real\text{-circle-sgn-r}[of of\text{-ocircle } H2 a2 r2]$
by *auto* (*metis neg-0-less-iff-less sgn-1-pos sgn-sgn*) +
hence $E \neq a1 E \neq a2$
using $\langle cdist E a1 = r1 \rangle \langle cdist E a2 = r2 \rangle$
by *auto*

let $?k = sgn\text{-bool } (?p1 = ?p2)$
let $?xx = ?k * (r1^2 + r2^2 - (cdist a2 a1)^2) / (2 * r1 * r2)$

have $\cos(\text{ang-circ } E a1 a2 ?p1 ?p2) = ?xx$
using $law\text{-of}\text{-cosines}[of a2 a1 E] * \langle r1 > 0 \rangle \langle r2 > 0 \rangle$ $\cos\text{-ang-circ-simp}[OF \langle E \neq a1 \rangle \langle E \neq a2 \rangle]$
by (*subst (asm) ang-vec-opposite-opposite' [OF ⟨E ≠ a1⟩ [symmetric] ⟨E ≠ a2⟩ [symmetric], symmetric]*) *simp*
moreover
have $\cos\text{-angle } H1 H2 = ?xx$
using $\langle r1 > 0 \rangle \langle r2 > 0 \rangle$
using $\langle (a1, r1) = euclidean\text{-circle } (of\text{-ocircle } H1) \rangle \langle (a2, r2) = euclidean\text{-circle } (of\text{-ocircle } H2) \rangle$
using $\langle is\text{-circle } (of\text{-ocircle } H1) \rangle \langle is\text{-circle } (of\text{-ocircle } H2) \rangle$
using $\langle circline\text{-type } (of\text{-ocircle } H1) < 0 \rangle \langle circline\text{-type } (of\text{-ocircle } H2) < 0 \rangle$
proof transfer
fix $a1 r1 H1 H2 a2 r2$
obtain $A1 B1 C1 D1$ **where** $HH1: Rep\text{-circline-mat } H1 = (A1, B1, C1, D1)$
by (*cases Rep-circline-mat H1*) *auto*
obtain $A2 B2 C2 D2$ **where** $HH2: Rep\text{-circline-mat } H2 = (A2, B2, C2, D2)$
by (*cases Rep-circline-mat H2*) *auto*
have $*: is\text{-real } A1 is\text{-real } A2 is\text{-real } D1 is\text{-real } D2$ $\text{cnj } B1 = C1 \text{ cnj } B2 = C2$
using $Rep\text{-circline-mat}[of H1]$ $Rep\text{-circline-mat}[of H2]$ $hermitean\text{-elems}[of A1 B1 C1 D1]$ $hermitean\text{-elems}[of A2 B2 C2 D2]$ $HH1 HH2$
by *auto*
have $\text{cnj } A1 = A1 \text{ cnj } A2 = A2$
using $\langle is\text{-real } A1 \rangle \langle is\text{-real } A2 \rangle$
by (*case-tac[!] A1, case-tac[!] A2, auto*)

```

assume  $\neg \text{circline-}A0\text{-rep } (\text{id } H1) \neg \text{circline-}A0\text{-rep } (\text{id } H2)$ 
hence  $A1 \neq 0 A2 \neq 0$ 
using HH1 HH2
by (auto simp add: circline- $A0$ -rep-def)
hence  $\text{Re } A1 \neq 0 \text{ Re } A2 \neq 0$ 
using ⟨is-real A1⟩ ⟨is-real A2⟩
by (metis complex- $\text{Im-zero}$  complex- $\text{Re-zero}$  complex-equality)+

assume circline-type- $\text{rep } (\text{id } H1) < 0$  circline-type- $\text{rep } (\text{id } H2) < 0$ 
assume  $(a1, r1) = \text{euclidean-circle-}\text{rep } (\text{id } H1) (a2, r2) = \text{euclidean-circle-}\text{rep }$ 
 $(\text{id } H2)$ 
assume  $r1 > 0 r2 > 0$ 

let ?D12 = mat-det-12 (Rep-circline-mat H1) (Rep-circline-mat H2) and ?D1
= mat-det (Rep-circline-mat H1) and ?D2 = mat-det (Rep-circline-mat H2)
let ?x1 = (cdist a2 a1)2 - r12 - r22 and ?x2 = 2*r1*r2
let ?x = ?x1 / ?x2
have *:  $\text{Re } (?D12) / (2 * (\sqrt{(\text{Re } (?D1 * ?D2))})) = \text{Re } (\text{sgn } A1) * \text{Re } (\text{sgn } A2) * ?x$ 
proof-
let ?M1 = (A1, B1, C1, D1) and ?M2 = (A2, B2, C2, D2)
let ?d1 = B1 * C1 - A1 * D1 and ?d2 = B2 * C2 - A2 * D2
have  $\text{Re } ?d1 > 0 \text{ Re } ?d2 > 0$ 
using HH1 HH2 ⟨circline-type- $\text{rep } (\text{id } H1) < 0$ ⟩ ⟨circline-type- $\text{rep } (\text{id } H2) < 0$ ⟩
by (auto simp add: circline-type- $\text{rep-def}$ )
hence **:  $\text{Re } (?d1 / (A1 * A2)) > 0 \text{ Re } (?d2 / (A2 * A1)) > 0$ 
using ⟨is-real A1⟩ ⟨is-real A2⟩ ⟨A1 ≠ 0⟩ ⟨A2 ≠ 0⟩
by – (simp add: Re-divide-real, metis Re-divide-real complex- $\text{Re-mult}$ 
divide-pos-pos eq-divide-imp mult-eq-0-iff not-real-square-gt-zero)+
have ***: is-real (?d1 / (A1 * A2)) ∧ is-real (?d2 / (A2 * A1))
using ⟨is-real A1⟩ ⟨is-real A2⟩ ⟨A1 ≠ 0⟩ ⟨A2 ≠ 0⟩ ⟨cnj B1 = C1[symmetric]
cnj B2 = C2[symmetric]⟩ ⟨is-real D1⟩ ⟨is-real D2⟩
by (subst div-reals, simp, simp, simp)+

have cor ?x = mat-det-12 ?M1 ?M2 / (2 * sgn A1 * sgn A2 * cor (sqrt (Re
?d1) * sqrt (Re ?d2)))
proof-
have A1*A2*cor ?x1 = mat-det-12 ?M1 ?M2
proof-
have 1: A1*A2*(cor ((cdist a2 a1)2)) = ((B2*A1 - A2*B1)*(C2*A1
- C1*A2)) / (A1*A2)
using ⟨(a1, r1) = euclidean-circle- $\text{rep } (\text{id } H1)$ ⟩ ⟨(a2, r2) = euclidean-circle- $\text{rep } (\text{id } H2)$ ⟩
unfolding cdist-def cmod-square
using HH1 HH2 * ⟨A1 ≠ 0⟩ ⟨A2 ≠ 0⟩ ⟨cnj A1 = A1⟩ ⟨cnj A2 = A2⟩
apply (subst complex-of-real-Re)
apply (simp add: complex-mult-cnj-cmod power2-eq-square)

```

```

apply (simp add: euclidean-circle-rep-def complex-cnj power2-eq-square
field-simps)
done
have 2:  $A1*A2*cor(-r1^2) = A2*D1 - B1*C1*A2/A1$ 
using  $\langle a1, r1 \rangle = euclidean-circle-rep(id H1)$ 
using  $HH1 *** *** \langle A1 \neq 0 \rangle$ 
apply (simp add: euclidean-circle-rep-def power2-eq-square)
apply (subst complex-of-real-Re, simp)
apply (simp add: field-simps)
done
have 3:  $A1*A2*cor(-r2^2) = A1*D2 - B2*C2*A1/A2$ 
using  $\langle a2, r2 \rangle = euclidean-circle-rep(id H2)$ 
using  $HH2 *** *** \langle A2 \neq 0 \rangle$ 
apply (simp add: euclidean-circle-rep-def power2-eq-square)
apply (subst complex-of-real-Re, simp)
apply (simp add: field-simps)
done
have  $A1*A2*cor((cdist a2 a1)^2) + A1*A2*cor(-r1^2) + A1*A2*cor(-r2^2)$ 
= mat-det-12 ?M1 ?M2
using  $\langle A1 \neq 0 \rangle \langle A2 \neq 0 \rangle$ 
by (subst 1, subst 2, subst 3) (simp add: field-simps)
thus ?thesis
by (simp add: field-simps)
qed

moreover

have  $A1 * A2 * cor(\langle x2 \rangle) = 2 * sgn A1 * sgn A2 * cor(\sqrt(Re ?d1) * \sqrt(Re ?d2))$ 
proof-
have 1:  $\sqrt(Re(?d1 / (A1 * A1))) = \sqrt(Re ?d1) / |Re A1|$ 
using  $\langle A1 \neq 0 \rangle \langle is-real A1 \rangle$ 
by (subst Re-divide-real, simp, simp, subst real-sqrt-divide, simp)

have 2:  $\sqrt(Re(?d2 / (A2 * A2))) = \sqrt(Re ?d2) / |Re A2|$ 
using  $\langle A2 \neq 0 \rangle \langle is-real A2 \rangle$ 
by (subst Re-divide-real, simp, simp, subst real-sqrt-divide, simp)

have  $sgn A1 = A1 / cor |Re A1|$ 
using  $\langle is-real A1 \rangle$ 
unfolding sgn-eq
by (cases A1, simp)

moreover
have  $sgn A2 = A2 / cor |Re A2|$ 
using  $\langle is-real A2 \rangle$ 
unfolding sgn-eq
by (cases A2, simp)

ultimately
show ?thesis
using  $\langle a1, r1 \rangle = euclidean-circle-rep(id H1) \langle a2, r2 \rangle = euclidean-circle-rep$ 

```

```

(id H2)@ HH1 HH2
  using *** ⟨is-real A1⟩ ⟨is-real A2⟩
    by (simp add: euclidean-circle-rep-def) (subst 1, subst 2, simp add:
of-real-numeral)
qed

ultimately

have (A1 * A2 * cor ?x1) / (A1 * A2 * (cor ?x2)) =
  mat-det-12 ?M1 ?M2 / (2 * sgn A1 * sgn A2 * cor (sqrt (Re ?d1) *
sqrt (Re ?d2)))
  by simp
thus ?thesis
  using ⟨A1 ≠ 0⟩ ⟨A2 ≠ 0⟩
  by simp
qed
hence cor ?x * sgn A1 * sgn A2 = mat-det-12 ?M1 ?M2 / (2 * cor (sqrt
(Re ?d1) * sqrt (Re ?d2)))
  using ⟨A1 ≠ 0⟩ ⟨A2 ≠ 0⟩
  by (simp add: sgn-zero-iff)
moreover
have Re (cor ?x * sgn A1 * sgn A2) = Re (sgn A1) * Re (sgn A2) * ?x
proof-
  have is-real (cor ?x) is-real (sgn A1) is-real (sgn A2)
    using ⟨is-real A1⟩ ⟨is-real A2⟩ Im-complex-of-real[of ?x]
    by auto
  thus ?thesis
    using Re-complex-of-real[of ?x]
    by (subst Re-mult-real3, auto simp add: field-simps)
qed
moreover
have *: sqrt (Re ?D1) * sqrt (Re ?D2) = sqrt (Re ?d1) * sqrt (Re ?d2)
  using HH1 HH2
  by (subst real-sqrt-mult[symmetric])+ (simp add: field-simps)
have 2 * (sqrt (Re (?D1 * ?D2))) ≠ 0
  using ⟨Re ?d1 > 0⟩ ⟨Re ?d2 > 0⟩ HH1 HH2 ⟨is-real A1⟩ ⟨is-real A2⟩ ⟨is-real
D1⟩ ⟨is-real D2⟩
  using Rep-circline-mat[of H1] mat-det-hermitean-real[of Rep-circline-mat
H1]
  by (subst Re-mult-real, auto)
hence **: Re (?D12 / (2 * cor (sqrt (Re (?D1 * ?D2)))))) = Re (?D12) /
(2 * (sqrt (Re (?D1 * ?D2)))))
  using ⟨Re ?d1 > 0⟩ ⟨Re ?d2 > 0⟩ HH1 HH2 ⟨is-real A1⟩ ⟨is-real A2⟩ ⟨is-real
D1⟩ ⟨is-real D2⟩
  by (subst Re-divide-real) (auto simp add: Im-complex-of-real)
have Re (mat-det-12 ?M1 ?M2 / (2 * cor (sqrt (Re ?d1) * sqrt (Re ?d2)))) =
= Re (?D12) / (2 * (sqrt (Re (?D1 * ?D2)))))
  using HH1 HH2 Rep-circline-mat[of H1] mat-det-hermitean-real[of Rep-circline-mat
H1]

```

```

    by (subst **[symmetric], subst Re-mult-real, simp, subst real-sqrt-mult, subst
*, simp)
ultimately
show ?thesis
by simp
qed
have **: pos-oriented-rep H1  $\longleftrightarrow$  Re A1 > 0 pos-oriented-rep H2  $\longleftrightarrow$  Re A2
> 0
using ⟨Re A1 ≠ 0⟩ HH1 ⟨Re A2 ≠ 0⟩ HH2
by (auto simp add: pos-oriented-rep-def)
show cos-angle-rep H1 H2 = sgn-bool (pos-oriented-rep H1 = pos-oriented-rep
H2) * (r12 + r22 - (cdist a2 a1)2) / (2 * r1 * r2)
unfolding cos-angle-rep-def Let-def
using ⟨r1 > 0⟩ ⟨r2 > 0⟩
by (subst divide-minus-left, subst *, subst Re-sgn[OF ⟨is-real A1⟩ ⟨A1 ≠ 0⟩],
subst Re-sgn[OF ⟨is-real A2⟩ ⟨A2 ≠ 0⟩], subst **, subst **, simp add: field-simps)
qed
ultimately
show ?thesis
by simp
qed

lemma [simp]: sqrt a * sqrt a = |a|
by (subst real-sqrt-mult[symmetric]) simp

lemma cos-angle H1 H2 = cos-angle (moebius-ocircline M H1) (moebius-ocircline
M H2)
proof transfer
fix H1 H2 M
show cos-angle-rep H1 H2 = cos-angle-rep (moebius-circline-rep M H1) (moebius-circline-rep
M H2)
unfolding cos-angle-rep-def Let-def moebius-circline-rep-Rep mat-det-12-congruence
mat-det-congruence
using Rep-moebius-mat[of M] mat-det-inv[of Rep-moebius-mat M]
by (auto simp add: power2-eq-square real-sqrt-mult field-simps)
qed

lemma
assumes mat-det (A, B, C, D) ≠ 0
shows moebius-circline (mk-moebius A B C D) imag-unit-circle = imag-unit-circle
 $\longleftrightarrow$ 
unitary-gen (A, B, C, D) (is ?lhs = ?rhs)
proof
assume ?lhs
thus ?rhs
using assms
proof transfer

```

```

fix A B C D :: complex
let ?M = (A, B, C, D) and ?E = (1, 0, 0, 1)
assume circline-mat-eq (moebius-circline-rep (mk-moebius-rep A B C D) imag-unit-circle-rep)
imag-unit-circle-rep mat-det ?M ≠ 0
then obtain k where k ≠ 0 ?E = cor k *sm congruence (mat-inv ?M) ?E
by (auto simp add: mk-moebius-rep-Rep)
hence unitary-gen (mat-inv ?M)
using mult-sm-inv-l[of cor k congruence (mat-inv ?M) ?E ?E]
unfolding unitary-gen-def
by (rule-tac x=1/cor k in exI, simp del: mat-inv.simps, metis eye-def
mat-eye-r)
thus unitary-gen ?M
using unitary-gen-inv[of mat-inv ?M] ⟨mat-det ?M ≠ 0⟩
by (simp add: mat-inv-inv del: mat-inv.simps)
qed
next
assume ?rhs
thus ?lhs
using assms
proof transfer
fix A B C D :: complex
let ?M = (A, B, C, D) and ?E = (1, 0, 0, 1)
assume unitary-gen ?M mat-det ?M ≠ 0
hence unitary-gen (mat-inv ?M)
using unitary-gen-inv[of ?M]
by simp
then obtain k where k ≠ 0 mat-adj (mat-inv ?M) *mm (mat-inv ?M) = cor
k *sm eye
using unitary-gen-real[of mat-inv ?M] mat-det-inv[of ?M]
by auto
hence *: ?E = (1 / cor k) *sm (mat-adj (mat-inv ?M) *mm (mat-inv ?M))
using mult-sm-inv-l[of cor k eye mat-adj (mat-inv ?M) *mm (mat-inv ?M)]
by simp
show circline-mat-eq (moebius-circline-rep (mk-moebius-rep A B C D) imag-unit-circle-rep)
imag-unit-circle-rep
using ⟨mat-det ?M ≠ 0⟩ ⟨k ≠ 0⟩
by (simp add: mk-moebius-rep-Rep del: mat-inv.simps) (rule-tac x=1/k in
exI, subst *, simp del: mat-inv.simps, metis eye-def mat-eye-r)
qed
qed
end

```